







FACULTEIT DIERGENEESKUNDE  
approved by EAEVE



Laboratorium voor Immunologie  
Vakgroep Virologie, Parasitologie en Immunologie  
Faculteit Diergeneeskunde  
Universiteit Gent

# Insights into the epidemiology of enteropathogens of young pigs raised in Cuban piggeries

**Pedro Yoelvys de la Fé Rodríguez**

Promotor:

**Prof. Dr. Bruno Maria Goddeeris**

Co-promotoren:

**Prof. Dr. Eric Cox en Prof. Dr. Luis O. Maroto Martin**

Proefschrift voorgelegd aan de Faculteit Diergeneeskunde tot het behalen van de  
graad van Doctor in de Diergeneeskundige Wetenschappen – Universiteit Gent, 2012



## Table of contents

List of abbreviations		7
Introduction		9
<b>Chapter 1</b>	<i>General introduction</i>	13
	<b>Aims of the study</b>	45
<b>Chapter 2</b>	<i>Association of toxigenic E. coli and several other pathogens with porcine pre- and post-weaning diarrhea in Villa Clara province, Cuba</i>	
	2.1 Abstract	49
	2.2 Introduction	50
	2.3 Materials and methods	50
	2.4 Results	54
	2.5 Discussion	57
	2.6 Acknowledgments	60
<b>Chapter 3</b>	<i>Antibiotic resistance and genetic relatedness among pathogenic E. coli isolated from intestinal contents of diarrheic piglets in Villa Clara province, Cuba</i>	
	3.1 Abstract	63
	3.2 Introduction	64
	3.3 Materials and methods	64
	3.4 Results	66
	3.5 Discussion	68
	3.6 Acknowledgments	70
<b>Chapter 4</b>	<i>High prevalence of F4<sup>+</sup> and F18<sup>+</sup> E. coli in Cuban piggeries as determined by serological survey</i>	
	4.1 Abstract	73
	4.2 Introduction	74
	4.3 Materials and methods	74
	4.4 Results	78
	4.5 Discussion	86
	4.6 Acknowledgments	88

<b>Chapter 5</b>	<i>Screening commercial pigs in Villa Clara province, Cuba, for mucin 4 polymorphisms and susceptibility/resistance to F18<sup>+</sup> E. coli</i>	
5.1	Abstract	91
5.2	Introduction	92
5.3	Materials and methods	92
5.4	Results and Discussion	94
5.5	Acknowledgments	97
<b>Chapter 6</b>	<i>General discussion and future perspectives</i>	
6.1	General discussion	101
6.2	Future perspectives	105
Summary		107
Samenvatting		111
Resumen		115
References		119
<i>Curriculum vitae</i>		143
Word of thanks		147

**List of abbreviations**

ABTS	2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid)
AIDA-I	adhesin involved in diffuse adherence
<i>B. coli</i>	<i>Balantidium coli</i>
<i>C. parvum</i>	<i>Cryptosporidium parvum</i>
<i>C. perfringens</i>	<i>Clostridium perfringens</i>
CPE	<i>C. perfringens</i> enterotoxin
DAS-ELISA	double-antibody-sandwich enzyme-linked immunosorbent assay
<i>E. coli</i>	<i>Escherichia coli</i>
EAST-I	enteroaggregative <i>E. coli</i> heat stable enterotoxin
EDTA	Ethylenediaminetetraacetic acid
ELISA	enzyme-linked immunosorbent assay
EM	electron microscopy
EPEC	enteropathogenic <i>E. coli</i>
ETEC	enterotoxigenic <i>E. coli</i>
FAT	fluorescent antibody technique
F18R	F18 receptor
F4R	F4 receptor
<i>I. suis</i>	<i>Isospora suis</i>
IC	immunochemistry
IF	immunofluorescence
IFAT	indirect fluorescence antibody technique
IHC	immunohistochemistry
Int	intimin
ISH	<i>In-situ</i> hybridization
kDa	kilo Dalton
LT	heat-labile enterotoxin
MAb	monoclonal antibody
OD	optical density
PAA	porcine attaching and effacing-associated factor
PAGE	polyacrylamide gel electrophoresis
pAPN	porcine aminopeptidase N
PCR	polymerase chain reaction

PCR-RFLP	PCR-restriction fragment length polymorphism
PEDV	porcine epidemic diarrhea virus
PRCV	porcine respiratory coronavirus
PRRSV	porcine respiratory and reproductive syndrome virus
RT-PCR	reverse transcriptase PCR
ST	heat-stable enterotoxin
TGE	transmissible gastroenteritis
TGEV	transmissible gastroenteritis virus
VN	virus neutralization
VTEC	verocytotoxigenic <i>E. coli</i>



## Introduction

Porcine pre- and post-weaning diarrhea negatively impact the economic feasibility of the swine industry due to mortality, costs of medication, and growth retardation (Francis, 1999; Katsuda *et al.*, 2006; Ushida *et al.*, 2009). Housing and management conditions (e.g. hygiene, comfort temperature, intake of maternal antibodies, feeding), vaccination against enteric pathogens, and surveillance of antibiotic resistance are crucial aspects to consider during prevention of porcine diarrhea (Fairbrother *et al.*, 2005; Fairbrother, 2006; Straw *et al.*, 2006).

Diarrhea of young pigs is frequently caused or complicated by rotavirus, transmissible gastroenteritis virus (TGEV), porcine epidemic diarrhea virus (PEDV), enterotoxigenic *Escherichia coli* (ETEC), toxigenic *Clostridium perfringens*, and *Coccidia* (Wieler *et al.*, 2001; Straw *et al.*, 2006; Katsuda *et al.*, 2006). Therefore, the differential identification of infectious agents is necessary to evaluate diarrhea epidemiology in a swine herd (Collins *et al.*, 1989; Niestrath *et al.*, 2002; Nuñez *et al.*, 2003). However, reports of surveys aimed at studying the mixed condition of piglet's infectious diarrhea have been scarce worldwide (Wieler *et al.*, 2001; Adesiyun *et al.*, 2001; Yaeger *et al.*, 2002; Katsuda *et al.*, 2006); many studies on porcine diarrhea have covered only single pathogens (Quilez *et al.*, 1996; Osek, 1999; Barreiros *et al.*, 2003), and although in a TGEV/PEDV prevalence survey, also pathogenic bacteria were identified, results were not shown (Chae *et al.*, 2000). When performing experiments related with the porcine digestive tract, the differential identification of enteropathogens should be carried out as enteropathogens can significantly influence the results (Jensen *et al.*, 2006; Niestrath *et al.*, 2010). Furthermore there is urgent need to better identify synergisms or antagonisms among these pathogens (Baba and Gaafar, 1985; Choi *et al.*, 2003).

ETEC are the most common cause of diarrhea in suckling and recently weaned pigs (Katsuda *et al.*, 2006). Cheng *et al.* (2006), Zhang *et al.* (2007), Madoroba *et al.* (2009), and Vidotto *et al.* (2009) found that F4 or F18 fimbriae were the major fimbrial antigens expressed by pathogenic *E. coli* associated with swine diarrhea. These fimbriae mediate the adhesion of ETEC to receptors present on the surface of enterocytes, favoring gut colonization. The subsequent production of enterotoxins by these bacteria leads to the secretion of electrolytes and water across the mucosa, resulting in a watery diarrhea. Additionally, F18 favors the adhesion of verocytotoxigenic *E. coli* (VTEC) which cause edema disease through a systemic vascular damage provoked by the verocytotoxin STx2e (Fairbrother, 2006; Fairbrother and Gyles, 2006; Oanh *et al.*, 2010).

Swine production is very important in Cuba as pork is the most consumed meat. In 2005 1,980,000 pigs were slaughtered and five years later already 3,266,600 according to the National Office for Statistics (ONE, 2011).

In Cuban piggeries diarrhea is common and strongly reduces the survival rate of young pigs leading to considerable economic losses: diarrheic diseases are responsible for 31% and 37% of the total mortality during the pre- and post-weaning periods, respectively (Cabrera and García, 2009). For instance, in the whole country 506,400 suckling piglets died in 2009 (12.2% mortality; ONE, 2011), and the 31% mortality provoked by diarrhea in this age group represent 156,984 deaths (Cabrera and García, 2009).

It is contradictory that in Cuba specific and updated epidemiological information related to swine enteropathogens is scarce. The prevention and control of diarrhea in Cuban piggeries are not always well conducted due to lack of infrastructure in the Provincial Veterinary Diagnostic Laboratories to properly perform the identification of enteropathogens (Cabrera *et al.*, 2010).

Pedroso and Talavera (1983) showed in 1983 the presence of F4<sup>+</sup> and F5<sup>+</sup> *E. coli* in feces of piglets in the Havana province, whereas Blanco *et al.* (2006) could not isolate F4<sup>+</sup>, F5<sup>+</sup> or F41<sup>+</sup> *E. coli* from diarrheic piglets in 2002 in the Villa Clara province. Most of their *E. coli* isolates (61%) were F18<sup>+</sup>. The prevalence of *Cryptosporidium parvum* and *Isospora suis* was studied more than 20 years ago in the Havana province in diarrheic piglets and was 2.1% and 44.7%, respectively (Cabrera and García, 1985; Koudela *et al.*, 1989). The first outbreak of epidemic transmissible gastroenteritis (TGE) in Cuba was reported in Havana in 2003 (Barrera *et al.*, 2005).

The present thesis contributes to the epidemiological characterization of enteropathogens associated with porcine pre- and post-weaning diarrhea in Cuba, with special emphasis on pathogenic *E. coli*. All epidemiological data obtained and discussed herein can be used for the implementation of accurate preventive and therapeutic strategies to control porcine enteropathogens in Cuba.

## Chapter 1

---

*General introduction*



### **Cuban context**

In 1493, during his second trip to America, Christopher Columbus carried eight Iberian-trunk pigs to Cuba on request of the queen Isabella I of Castile and León (Laguna, 1998). Nevertheless, for more than four centuries during the colonial and neo-colonial periods, the production of cane sugar, alcohol, tobacco, and leather were primordial in Cuba and pork was not an important production. By 1959, 68% of swines were bred extensively. Approximately, 16% of the herds were intensively managed by breeders and the remainder was a family-type production. From a genetic point of view, 22% of swine were pure Criollo, and 69% were Criollo's crossbreds. Only 9% of swine herds had specialized breeds, mainly Hampshire and Duroc. These genetic and management statuses did not allow an efficient swine production leading to a high import dependency from the North American market before 1959 (Rico, 2005).

From the early 60's, as part of the changes that occurred in Cuba due to the 1959 Revolution, swine production benefited from the following strategies:

- i- Creation of the swine production department in the Ministry of Agriculture and elaboration of a national policy for pig health and management.
- ii- Foreign consultancy and staff training.
- iii- Building of farms for intensive swine production and introduction of pure swine breeds into breeding programs monitored by swine genetic centers located all over the country, which ensured specialized swine production.

From 1971 to 1989, artificial insemination started to be applied and swine production was re-organized under a pyramidal structure including genetic centers, multiplication piggeries (which obtained replacement stock animals) and commercial piggeries. Also, swine sector benefited from investments in facilities and from importations (mainly food and medicines) at preferential prices agreed with the Council of Mutual Economic Assistance (COMECON). During this period, about 84% of swine production occurred by the specialized state sector and the remaining 16% by private producers and agricultural cooperatives (Rico, 2005).


From the 90's onwards, Cuban economy was negatively impacted by the dissolution of COMECON due to the collapse of the communism in Eastern Europe and the Soviet Union, which were at that moment the main trade partners and sources for technology transfer. Then, as happened with most institutions, the services of the Cuban Veterinary Medicine Institute as well as swine production reduced

their activity. Specifically, the diagnostic effectiveness decreased and ongoing research on animal health slackened, resulting in insufficient diagnosis of viral and bacterial agents. As a result, information regarding swine enteric pathogens have been limited in Cuba:

- Pedroso and Talavera (1983) detected F4<sup>+</sup> and F5<sup>+</sup> *E. coli* by immunofluorescence in feces of piglets in the Havana province.
- The prevalence of *C. parvum* in diarrheic piglets from Havana was 2.5% in 1985 (Cabrera and Garcia, 1985).
- 44.7% of diarrheic piglets were infected with *I. suis* in a piggery from Havana in 1989 (Koudela *et al.*, 1989).
- Fuentes *et al.* (2001) reported a 2.8% prevalence of F4<sup>+</sup> *E. coli* in 212 diarrheic piglets sampled in 1998 in the Camagüey province.
- The first outbreak of TGEV was reported in 2003 in Havana and the disease was later reproduced experimentally (Barrera *et al.*, 2005).
- Blanco *et al.* (2006) genotyped *E. coli* isolates from diarrheic piglets in 2002 and found that most of them (61%) carried the F18 encoding gene and that F4 encoding gene was absent.

The Cuban Institute for Swine Research recently reported that gastroenteric diseases cause 31% and 37% of mortality in suckling and weaned piglets, respectively (Cabrera and García, 2009). In suckling piglets, another causes of death, most of them closely related with diarrhea, such as crushing by the sow (14%), malnutrition (8%), congenital diseases (e.g. myoclonia; 6%), low weight at birth (4%), and hypoglycemia (3%) also require special attention. In weaned piglets, respiratory diseases (9%), malnutrition (5%), and classic swine fever (5%) are common causes of death (GRUPOR, 2008; Cabrera and García, 2009). Therefore, in 2008 the vice-director of the Cuban National Institute of Veterinary Medicine recommended an improvement on swine management and on the specific diagnosis of enteric diseases in order to decrease mortality by diarrhea in young pigs (Ricardo, 2008).

Nowadays, policies of the Cuban Ministry of Agriculture are to improve and increase swine production as pork is the most consumed meat in the country. In 2008 the Cuban swine herd was estimated to be 1,878,600 pigs (ONE, 2011), with the national company GRUPOR ruling the swine production (Fig. 1; Rico, 2005).



**Figure 1.** Organization of the swine production in Cuba.

**Table 1.** Range of age at which the most common swine enteropathogens can cause disease.

	Age of pigs				
	24 hours	5 days	3 weeks	5 weeks	10 weeks
Enteropathogens					
TGEV	-----	-----	-----	-----	-----
PEDV	-----	-----	-----	-----	-----
Rotavirus	-----	-----	-----	-----	-----
<i>E. coli</i>	-----	-----	-----	-----	-----
<i>Clostridium</i>	-----	-----	-----	-----	-----
<i>Salmonella</i>	-----	-----	-----	-----	-----
<i>Isospora suis</i>	-----	-----	-----	-----	-----
<i>Strongyloides</i>	-----	-----	-----	-----	-----

**Multi-infectious etiology of diarrhea in young pigs**

The clinical pictures of porcine pre- or post-weaning diarrhea often do not point to a specific etiology because of variations in clinical signs and epidemiology that can be produced by the effect of one or more disease agents. A control program of diarrhea could fail if one agent playing a primary role in the pathogenesis remains sub-diagnosed. An explosive onset and rapid spread of diarrhea is usually associated with a viral etiology. An insidious onset, slow spread, and gradual increase in severity over time tend to be seen with bacterial or parasitic diseases (Straw *et al.*, 2006).

A useful tool for veterinary practitioners regarding presumptive diagnosis of infectious diarrhea is the age at which piglets are affected. But this is rather an indicator of which pathogens could be responsible, and is not as precise as a specific etiologic identification test (Table 1; Straw *et al.*, 2006).

The prevalence and repetitiveness of combined infections of enteropathogens reported worldwide lead to the introduction of the term “*mixed-type*” during epidemiological analysis of diarrhea. However, most prevalence studies on swine enteropathogens focused on single pathogens (Quilez *et al.*, 1996; Barreiros *et al.*, 2003) and checking for a multi-infectious etiology has often been neglected (Chae *et al.*, 2000). Studies aimed at the differential identification of gastroenteric infections have displayed swine diarrhea as a complex syndrome in which several pathogens can be associated (Table 3), explaining a difficult control of pre- and post-weaning diarrhea in a piggery where enteropathogen identification has not been well performed. A few surveys undertaken to differentially identify enteric pathogens in suckling and weaned pigs with diarrhea indicate a high diversity of mixed infections which might lead to or complicate diarrheal outbreaks in a piggery (Table 4). However, worldwide available data regarding combined infections of enteropathogens remain difficult to compare as different tests were used for the same pathogen or diagnoses did not cover all potential enteropathogens.

The main characteristics for differential diagnosis of swine gastroenteric infections were summarized in Table 2 (adapted from Thomson, 2006).



**Table 2.** Insights for the differentiation of some common enteric diseases of young pigs.

Pathogens	Age	Signs	Gross lesions	Histological lesions	Laboratory methods
TGEV and PEDV	All ages	Watery diarrhea. Rapid dehydration. Vomiting	Thin-walled pale intestine, sparse content	Severe villous atrophy	PCR, IHC, ISH, or IF of intestinal contents. Serology
Rotavirus	1 day to 7 weeks old. Most frequent at 2-3 weeks	Watery-pasty diarrhea, sub-clinical. Varying dehydration	Fluid ingesta, pale intestines. Sparse stomach contents	Moderate villous atrophy	Virus detection: PAGE, PCR, ELISA, Tissue IHC
<i>E. coli</i> (ETEC, EPEC)	Neonatal: 1-4 days. Post-weaning: 1-3 weeks after weaning.	Watery, yellowish diarrhea. Sudden death. Dehydration.	Fluid ingesta, small intestinal congestion, watery content. Stomach full of milk.	Mucosal congestion, edema. Bacterial attachment to gut epithelia.	Culture, serotypes of isolates, PCR. Tissue IHC. Agglutination test.
<i>C. perfringens</i> type C	1-14 days (rarely older)	Hemorrhagic watery diarrhea. Sudden death	Hemorrhagic intestines, mucosal necrosis	Mucosal necrosis, associated with Gram <sup>+</sup> rods	<i>C. perfringens</i> toxins ELISA's on intestinal content. Histopathology
<i>Salmonella</i> spp.	All ages after weaning	Variable, watery muco-hemorrhagy. Most infections subclinical	Fibrinous or hemorrhagic, ulcers, intestinal lesions	Ulcers, neutrophil infiltration, fibrinous thrombi	Culture, serotyping, phage type. Antibody detection.
<i>Cryptosporidium</i> spp.	3 days to weaning	Mild-moderate yellowish diarrhea. Varying degrees of dehydration	Fluid ingesta	None or mild villous atrophy. Oocysts in the epithelium	Mucosal smear for cryptosporidial oocysts. Histopathology
<i>I. suis</i>	5-21 days	Watery/yellowish diarrhea. Dehydration	Fluid ingesta, necrosis of intestinal mucosa	Villous atrophy, fibroncrotic enteritis, intracellular coccidians.	Stained mucosal smear. Histopathology. Identification of <i>Coccidia</i>

**Table 3.** Reports on differential identification and occurrence of enteropathogens in suckling (S) and weaned (W) pigs suffering from diarrhea worldwide.

Pigs	Pathogenic viruses (%)					Pathogenic bacteria (%)			
	Rotavirus	Coronavirus	Sapovirus	Adenovirus	PRRSV	ETEC	Salmonella	C. perfringens	C. difficile* E. durans
S	8	59.2 (TGEV)	NT	0.53	NT	19.6	NI	0.40 (type C)	NT
	17	NT	NT	NT	NT	18.2	NT	NT	NT
	5.1	3.4	NT	NT	NT	7.1	NI	NT	NT
	42	6 (TGEV)	NT	NT	15	9**	NI	6 <sup>NE</sup>	55
	69.3	0	5.3	NT	NT	13.3	0	0 (type C)	NT
	20.7	NT	NT	NT	NT	9.9	NT	0 (type C)	NT
W	13.3	NT	NT	NT	NT	10	NT	43.3 (type A), 0 (C)	23.3
	0	51.6	NT	NT	NT	42.7	NI	NT	NT
S&W	59.3	0	16.2	NT	NT	54.7	0	0 (type C)	NT
	4	13.4	NT	NT	NT	17.6	NI	NT	NT

Pigs	Parasites (%)					Negatives (%)	Country	Reference
	I. suis	Eimeria spp.	C. parvum	B. coli	Helminths			
S	15.3	0	NT	NT	NT	9.2	Canada	Morin et al., 1983
	53.8	0	NT	NT	NT	NI	Australia	Driesen et al., 1993
	28.8	NI	NI	NI	0	NI	Germany	Wieler et al., 2001
	NT	NT	NT	NT	NT	NI	U.S.A.	Yaeger et al., 2002
	18.7 (Coccidia)	0	NI	NI	NI	17	Japan	Katsuda et al., 2006
	18.8 (Coccidia)	NT	NT	6.9	NT	15.8	Romania	Costinar et al., 2010
W	0	NI	NI	NI	NI	NI	Brazil	Cruz et al., 2010
	1.2	NI	NI	NI	0	NI	Germany	Wieler et al., 2001
S&W	18.6 (Coccidia)	16.2	NI	NI	NI	9.5	Japan	Katsuda et al., 2006
	20.9	0.7	1.4	0.7	0	57.7	Germany	Wieler et al., 2001

NT, not tested; NI, no information; \*, toxin detection; \*\*, no determination of virulence factors and serotypes; <sup>NE</sup>, necrotic enteritis.

**Table 4.** Occurrence (%) of enteric mixed infections in suckling (S) and weaned (W) pigs with diarrhea reported worldwide.

"Mixed types"	Tested	%	S/W	Country	Reference
Rotavirus + PEDV	157 98	45.2 9.1	S	South Korea Czech Republic	Song <i>et al.</i> , 2006 Czanderlova <i>et al.</i> , 2010
Rotavirus + TGEV	749 98	0.8 4.5	S	Canada Czech Republic	Morin <i>et al.</i> , 1983 Czanderlova <i>et al.</i> , 2010
Rotavirus + TGEV + PEDV	98	22.7	S	Czech Republic	Czanderlova <i>et al.</i> , 2010
Rotavirus + PRRSV	100	1	S	U.S.A.	Yaeger <i>et al.</i> , 2002
Rotavirus + Sapovirus	153 116	1.3 1.7	S W	Japan	Katsuda <i>et al.</i> , 2006
Rotavirus + Sapovirus + ETEC	153 116	2 2.6	S W		
Rotavirus + TGEV + ETEC	749	0.66	S	Canada	Morin <i>et al.</i> , 1983
PRRSV + <i>Clostridium difficile</i>	100	3	S	U.S.A.	Yaeger <i>et al.</i> , 2002
Rotavirus + <i>Clostridium difficile</i>	100	6			
TGEV + ETEC	749	4.3	S	Canada	Morin <i>et al.</i> , 1983
Rotavirus + ETEC	749	1.07	S	Canada	Morin <i>et al.</i> , 1983
	1054	1.0		Australia	Driesen <i>et al.</i> , 1993
	100	1.0		U.S.A.	Yaeger <i>et al.</i> , 2002
	153 116	6.5 19.0	W	Japan	Katsuda <i>et al.</i> , 2006
Rotavirus + Sapovirus + ETEC + <i>Coccidia</i>	116	6.0	W	Japan	Katsuda <i>et al.</i> , 2006
Rotavirus + Sapovirus + ETEC + <i>C. parvum</i>	116	0.9	W		
Sapovirus + ETEC + <i>C. parvum</i>	116	0.9	W		
Rotavirus + Sapovirus + <i>Coccidia</i>	153 116	0.7 0.9	S W		
Rotavirus + Sapovirus + <i>C. parvum</i>	116	0.9	W		
Rotavirus + TGEV + <i>Coccidia</i>	749	0.13	S	Canada	Morin <i>et al.</i> , 1983
Rotavirus + ETEC + <i>Coccidia</i>	1054	0.6	S	Australia	Driesen <i>et al.</i> , 1993
	101	2.9		Romania	Costinar <i>et al.</i> , 2010
	153 116	1.3 3.4	W	Japan	Katsuda <i>et al.</i> , 2006
Rotavirus + ETEC + <i>Coccidia</i> + <i>C. parvum</i>	116	2.6	W	Japan	Katsuda <i>et al.</i> , 2006
Rotavirus + ETEC + <i>Coccidia</i> + <i>B. coli</i>	101	0.9	S	Romania	Costinar <i>et al.</i> , 2010
Rotavirus + <i>Coccidia</i>	9*	55.6	S	U.S.A.	Roberts and Walker, 1982
	749	1.2		Canada	Morin <i>et al.</i> , 1983
	1054	6.7		Australia	Driesen <i>et al.</i> , 1993
	101	8.9		Romania	Costinar <i>et al.</i> , 2010
	153 116	7.2 2.6	W	Japan	Katsuda <i>et al.</i> , 2006
Rotavirus + <i>Coccidia</i> + <i>B. coli</i>	101	6.9	S	Romania	Costinar <i>et al.</i> , 2010
Rotavirus + <i>C. parvum</i>	116	2.6	W	Japan	Katsuda <i>et al.</i> , 2006
Sapovirus + <i>Coccidia</i>	153	0.7	S		
TGEV + <i>Coccidia</i>	749	2.8	S	Canada	Morin <i>et al.</i> , 1983
ETEC + <i>Clostridium difficile</i>	100	1.0	S	U.S.A.	Yaeger <i>et al.</i> , 2002

**Table 4** (continuation). Occurrence (%) of enteric mixed infections in suckling (S) and weaned (W) pigs with diarrhea reported worldwide.

"Mixed types"	Tested	%	S/W	Country	Reference
ETEC + <i>Coccidia</i>	749	0.8	S	Canada	Morin <i>et al.</i> , 1983
	1054	10.7		Australia	Driesen <i>et al.</i> , 1993
	153	2.6		Japan	Katsuda <i>et al.</i> , 2006
	116	0.9	W		
ETEC + <i>C. parvum</i>	116	2.6	W	Japan	Katsuda <i>et al.</i> , 2006
Occurrence of multiple infections	749	11.7	S	Canada	Morin <i>et al.</i> , 1983
	1054	19.1		Australia	Driesen <i>et al.</i> , 1993
	100	12		U.S.A.	Yaeger <i>et al.</i> , 2002
	101	27.7		Romania	Costinar <i>et al.</i> , 2010
	153	22.2			
	116	47.4	W	Japan	Katsuda <i>et al.</i> , 2006
	92	9.0		Hungary	Nagy <i>et al.</i> , 1996
	287	9.1	S&W	Germany	Wieler <i>et al.</i> , 2001

In this table and in references to this table *Coccidia* refers to *I. suis* and or *Eimeria* spp.

As shown in Table 4, combined infections of Rotavirus-ETEC, Rotavirus-*Coccidia*, ETEC-*Coccidia*, and Rotavirus-ETEC-*Coccidia* have been commonly reported. Despite a low occurrence, the associations Rotavirus-ETEC-*Coccidia*-*C. parvum*, TGEV-Rotavirus-ETEC, and TGEV-Rotavirus-*Coccidia*, are important to consider because of the proven pathogenic effect of the involved agents.

Katsuda *et al.* (2006) in Japan and Costinar *et al.* (2010) in Romania reported that 22.2% and 27.7% of suckling diarrheic piglets, respectively, carried mixed infections. Therefore, mixed infections have to be accurately identified during diarrhea outbreaks to plan and implement an efficient disease surveillance, prevention and control in every piggery, geographic area, or management system.

Knowledge on the role of individual enteropathogens during pathogenesis and course of combined enteric infections is limited. Practically, no controlled experiments have been designed to study the interaction between the diverse enteropathogens; however, interesting findings have been reported and discussed by some authors:

- Baba and Gaafar (1985) demonstrated that piglets which have been infected with *S. typhimurium* and subsequently with *I. suis*, showed significantly smaller ( $p < 0.05$ ) *S. typhimurium* counts in the feces, caecal contents and mesenteric lymph nodes than those infected with *S. typhimurium* alone. They mentioned as possible causes the cleansing effect and low intestinal pH due to coccidial diarrhea, but the exact mechanism remains to be clarified.

- Lecce *et al.* (1982) suspected that rotavirus provokes a damage of the epithelium that can alter binding sites on enterocytes, favoring gut colonization by ETEC. Similar observations were made by Melin *et al.* (2004).
- Necropsy seventeen days post-*Ascaris* infection revealed lower hepatic fibrosis and a lesser degree of liver eosinophilia in *Ascaris* infected pigs pre-inoculated with *Salmonella* than those receiving *Ascaris* inoculation only (Wade and Gaafar, 1981). Similar results were seen in pigs recovered from TGE and later infected with *Ascaris* (Gaafar *et al.*, 1973).
- Janke *et al.* (1988) concluded that rotaviral infection in colostrum-deprived piglets is inhibited by enteroviruses infection.
- Choi *et al.* (2003), after using plant lectins for studying the histochemistry of the jejunal mucosa of pigs infected with *I. suis*, argue that variations on the glycoconjugates composition due to isosporosis could favor *E. coli* colonization.
- Enemark *et al.* (2003a) found that mixed infections of rotavirus and *Cryptosporidium* cause a dramatic aggravation of diarrhea and clinical signs; piglets mono-infected with Rotavirus or *Cryptosporidium* showed no or very mild clinical signs of illness.
- *C. perfringens* type C may colonize lesions initiated by *Coccidia*, rotavirus, TGEV and PEDV (Songer and Uzal, 2005).
- Zintl *et al.* (2007) found that *Cryptosporidium* infections are not commonly combined with *Salmonella* infections in pigs.
- Jung *et al.* (2008) proposed that the severe diarrhea in PEDV-rotavirus A co-infected piglets may be more associated with the immunity level of the host rather than to any synergistic effect of rotavirus on PEDV enteritis. They suggested that concurrent infection with porcine rotavirus A does not synergistically enhance intestinal villous atrophy.
- Kim *et al.* (2010a) reported that 124/182 (68%) rotavirus A infections were mixed with other enteric pathogens.

### **Major enteropathogens causing diarrhea in young pigs**

#### ***Escherichia coli***

*E. coli* are an important cause of diarrhea in suckling and recently weaned pigs, and are responsible for significant economical losses worldwide. Most porcine ETEC and VTEC strains produce fimbriae, which enable them to colonize the epithelial surface of the porcine small intestine. Pathology is the result of the production of toxins. ETEC strains produce heat-stable (STa or STb) or heat-labile (LT) which induce fluid loss resulting in diarrhea (Nataro and Kaper, 1998).

VTEC produce the verocytotoxin STx2e which is responsible for edema disease. Some *E. coli* strains can produce both enterotoxins and the verocytotoxin STx2e which is responsible for edema disease, and are appropriately referred to as ETEC/VTEC (Blanco *et al.*, 2006).

Attachment to specific receptors is essential for colonization and pathogenesis. Fimbriae mediate the attachment of *E. coli* to the intestinal mucosal surface. F4 and F18 fimbriae have been commonly identified in *E. coli* associated with PWD. But, whereas F18 fimbriae are almost exclusively associated with PWD, F4 fimbriae are also a predominant colonization factor implicated in neonatal diarrhea. Of F4, 3 serotypes have been described, namely F4ab, F4ac and F4ad, of which F4ac is the most prevalent (Choi and Chae, 1999). Of F18, two serotypes have been identified: F18ab and F18ac. F18ab is more expressed by VTEC strains and F18ac by ETEC strains. Other fimbriae such as F5, F6, and F41 are mainly expressed by porcine ETEC isolated from newborn piglets with diarrhea. Nevertheless, F4 and to a lesser extent F18, continue to be the major fimbrial antigen types in ETEC and VTEC identified in diagnostic laboratories in the U.S.A. (Moon *et al.*, 1999), China (Cheng *et al.*, 2006), Brazil (Vidotto *et al.*, 2009), Zimbabwe (Madoroba *et al.*, 2009), and Vietnam (Oanh *et al.*, 2010).

Some candidate receptors have been suggested for F4<sup>+</sup> *E. coli* in the gut epithelium of pigs (F4R). Erickson *et al.* (1992; 1994) proposed the intestinal mucin-like sialoglycoproteins (F4acR; IMTGP), Grange and Mouricout (1996) reported an intestinal transferrin (F4abR) and Grange *et al.* (1999) an intestinal glycosphingolipid (F4adR). Recently, Rasschaert (2008) identified porcine aminopeptidase-N (pAPN) as a receptor for F4ac. Looking at binding to glycosphingolipids, Coddens *et al.* (2011) found that F4ab binds galactosylceramide, sulfatide, sulf-lactosylceramide and globotriaosylceramide present in epithelial cells of the porcine intestine, whereas F4ac only binds galactosylceramide.

Not all pigs have receptors for adhesion of F4<sup>+</sup> and F18<sup>+</sup> *E. coli*. Some pigs are receptor negative. Testing *in vitro* adhesion of the three F4 variants (ab, ac, and ad) to brush border membranes of enterocytes, isolated enterocytes or isolated villi, six phenotypes could be identified: phenotype A binds all three F4 variants, phenotype B F4ab and F4ac, phenotype C F4ab and F4ad, phenotype D F4ad, phenotype E none of the variants, and phenotype F F4ab only. Based on that classification, prevalence studies have been done in different swine herds: in Midwestern United States and The Netherlands 30% and 50% of pigs showed the phenotype E, respectively (Sellwood *et al.*, 1975, Baker *et al.*, 1997; Bijlsma *et al.*, 1985). Conversely, 80% of the Belgian pigs were classified in the F4 susceptible phenotype A, and only 4% in the resistant phenotype E (Cox and Houvenaghel, 1987).

The F4ab/ac receptor loci are closely linked and linked to the transferrin locus on chromosome 13. They behave as a single autosomal dominant gene (Guérin *et al.*, 1993). Python *et al.* (2002)

concluded that the receptor for F4ac binds F4ab bacteria as well, and that it is controlled by one gene localized between S0068 and Sw1030 on chromosome 13. Jørgensen *et al.* (2004) detected that a mutation G→C in intron 7 of *mucin 4* gene was strongly associated with the ETEC F4ab/ac adhesive phenotype. They developed a PCR-RFLP which detected polymorphisms in a *mucin 4* gene fragment by its digestion with *Xba*I enzyme, and allows genotyping pigs for resistance or susceptibility to adhesion of *E. coli* mediated by F4ab/ac fimbriae. Three different pig genotypes can be observed: resistant (no adhesion) homozygote carrying indigestible alleles (RR) and susceptible heterozygote (SR) as well as homozygote (SS). Conversely, Rasschaert *et al.* (2007) could not confirm this good correlation when comparing results of the DNA-based marker test with results of the *in vitro* villous adhesion assay. A reason for the difference in both studies could be the use of a different *in vitro* adhesion assay as Jørgensen *et al.* (2004) used single enterocytes whereas Rasschaert *et al.* (2007) used isolated villi.

Looking at binding of F18 to glycosphingolipids, Coddens *et al.* (2009) could identify the receptors for F18. They observed a high specific interaction of F18 *E. coli* with glycosphingolipids having blood group A/B/H determinants on type 1 core chains, as well as the blood group A type 4 heptaglycosylceramide.

The *in vitro* villous adhesion tests also allow discrimination between F18R<sup>+</sup> and F18R<sup>-</sup> pigs. Absence or presence of the F18R is genetically determined and susceptibility being dominant over resistance (Bertschinger *et al.*, 1993). The gene controlling expression of the F18R was mapped to the halothane linkage group on pig chromosome 6 (Meijerink *et al.*, 1997). This locus contained two candidate genes, *FUT1* and *FUT2*, both encoding α2-fucosyltransferases. Expression analysis of these two genes in the porcine small intestine revealed that only the *FUT1* gene was expressed in all examined pigs (Meijerink *et al.*, 2000). This gene is localized in chromosome 6q11 and the enzyme alpha (1,2)-fucosyltransferase 1 transfers fucose to lipid, carbohydrate, and protein backbones present in the intestine. Sequencing of this gene showed a polymorphism (G or A) at nucleotide 307 resulting in an amino acid substitution at position 103 (Ala/Thr) of the enzyme. Presence of the A nucleotide on both alleles (*FUT1A/A* genotype) led to significantly reduced activity of the enzyme corresponding to the F18-fimbriated *E. coli* resistant genotype, whereas susceptible pigs had either the heterozygous *FUT1G/A* or the homozygous *FUT1G/G* genotype. These findings have led to the development of a PCR-RFLP using *Cfo*I enzyme that allows the classification of pigs into susceptible or resistant to F18<sup>+</sup> *E. coli* adhesion (Meijerink *et al.*, 1997).

Frydendahl *et al.* (2003) found a high correlation between F18R<sup>+</sup> genotypes and susceptibility to F18<sup>+</sup> *E. coli*; however, pigs carrying the resistant F18R genotype were not entirely protected against intestinal colonization. Coddens *et al.* (2007) also found a significant positive but weak correlation

( $r=0.307$ ,  $p<0.05$ ) between the *FUT1* genotype and the F18R phenotype. Indeed they reported a weak adhesion of F18<sup>+</sup> *E. coli* to the villous brush border in some genetically F18R<sup>-</sup> pigs and absence of adhesion in some F18R<sup>+</sup> pigs.

Heat-labile enterotoxins (LT) produced by ETEC have a high molecular weight (86 kDa) and two major serogroups, which do not cross-react immunologically, have been identified: LT-I and LT-II. Only LT-I is neutralized by anti-cholera toxin. LT produced by porcine *E. coli* (LTp) belongs to the LT-I group. The toxin complex is constituted by one A subunit (biologically active) and five B subunits, which bind mainly to gangliosides and some intestinal glycoproteins (Teneberg *et al.*, 1994). The A subunit stimulates the production of c-AMP by activating adenylate cyclase located on the basolateral membrane of polarized intestinal epithelial cells, resulting in a secretory diarrhea due to a decrease in Na<sup>+</sup> absorption by villus cells and activation of Cl<sup>-</sup> secretion by crypt cells. As for cholera toxin, additional mechanisms influencing intestinal motility and ion secretion have been attributed to the mode of action of LT, i.e. increase of prostaglandin-E release, modeling of the enteric nervous system via vasoactive intestinal polypeptides and a mild intestinal inflammatory response involving interleukin-6 (Sears and Kaper, 1996; Nataro and Kaper, 1998; Fairbrother and Gyles, 2006).

The heat-stable enterotoxins (STa and STb) are antigenically and genetically unrelated monomeric toxins that contain multiple cysteine residues, whose disulfide bonds account for the heat stability. STa is a small (2 kDa), non-immunogenic, protein most effective in piglets younger than 2 weeks of age. STa acts on guanylate cyclase so increasing the concentration of c-GMP in epithelial cells (Vaandrager *et al.*, 1994), leading to inhibition of Na<sup>+</sup> absorption and stimulation of Cl<sup>-</sup> secretion with water passively following resulting in a secretory diarrhea. Two variants have been described for STa: STp has been identified in porcine, bovine and human ETEC, while ST<sub>h</sub> is only produced by ETEC strains infecting humans (Fairbrother and Gyles, 2006). STb is also a small (5 kDa), poorly immunogenic, protein that strongly binds to acidic glycosphingolipids, including sulfatide (or 3'-sulfolactosylceramide) and several gangliosides (Dubreuil, 1997; Rousset *et al.*, 1998).

Table 5 shows results of diverse surveys aimed at detecting virulence encoding genes in *E. coli* isolated from feces of young pigs worldwide.

In addition to the natural existence of diverse *E. coli* virulence factors which lead host colonization and disease, the increase in antimicrobial resistance which is occurring in ETEC and VTEC strains, make swine colibacillosis difficult to control. Due to the young age in which piglets become infected with these pathogens, many breeders easily and systematically administer antibiotics resulting in a high selection pressure towards resistance. Boerlin *et al.* (2005), Travis *et al.* (2006) and Wang *et al.* (2010) reported clear associations between antibiotic resistance and virulence genes in swine pathogenic *E. coli*. For instance, genes coding for STa/STb and tetracycline



resistance were found to be co-located on a self-conjugative plasmid (pTC, 120-kb) which is widely distributed among porcine ETEC (Olasz *et al.*, 2005).

As good management, hygiene and vaccination in swine farms, also antibiotics have helped in preventing and controlling swine colibacillosis, but it also provoked the appearance and selection of resistant and multidrug-resistant *E. coli* which tend to persist in time and space (Maidhof *et al.*, 2002; Blake *et al.*, 2003; Moredo *et al.* 2007; Dewulf *et al.*, 2007; Akwar *et al.* 2008; Vieira *et al.*, 2009; Bibbal *et al.*, 2009). In New Zealand, Nulsen *et al.* (2008) found a higher antibiotic resistance to tetracycline (60% v/s 5%), streptomycin (25% v/s 3%), and cotrimoxazole (11% v/s 0%) in *E. coli* isolated on conventional farms (where is a higher antibiotic pressure) than on organic farms. Maynard *et al.* (2003) concluded that the genes behind phenotypic antibiotic resistance are not static and their prevalence is determined by various selection forces such as the use of specific antimicrobials.

Investigation of virulence factors, antimicrobial resistance, and genetic profiles are essential to deeply study the epidemiology of ETEC associated with swine diarrhea. ETEC carrying the same virulence profiles and with similar antibiogram sensitivity are likely to show a highly similar pulsotype by PAGE (Lee *et al.*, 2009; Bibbal *et al.*, 2009). Such mixed studies are becoming an important epidemiological tool. Thorsteinsdottir *et al.* (2010) found the same resistance profile and pulsotype among *E. coli* isolated from broiler meat and slaughterhouse workers. They stated that isolates sharing the same genetic profile and resistance patterns can arise on different farms. Conversely, Rosengren *et al.* (2009) and Smith *et al.* (2010) did not find clear associations between antimicrobial resistance and virulence profile in *E. coli* isolated from healthy versus diseased pigs. Surveillance studies of swine colibacillosis should be applied to every geographic area or even at farm or production system level as a high variety of antibiotics, resistance genes, virulence genes, and their co-location onto conjugative plasmids or pathogenicity islands in *E. coli* lead to diverse associations or clones (Hendriksen *et al.*, 2008; Harada *et al.*, 2008; Wang *et al.*, 2010; Smith *et al.*, 2010). Additionally, the co-selection of *E. coli* resistant to some antibiotics (i.e. kanamycin) by the use of other antibiotics (i.e. tetracycline) is contributing to increase antibiotic resistance in swine farms. Therefore risk assessment has to be performed for every antibiotic or every chemical group in order to better police the selection and persistence of resistant *E. coli* (Harada *et al.*, 2008).

**Table 5.** Recent studies reporting virulence encoding genes in *E. coli* isolated from feces of young pigs worldwide.

Syndrome & age group	n	% from tested isolates													Country/Reference
		VF+	F4	F5	F6	F41	F18	Int	STa	STb	LT	STx2e	EAST1		
ND															
(<21 d.)	200	63	33.5	10.5	0	0	0	NT	58	52.5	35	NT	NT	Vietnam/Do et al., 2006	
	13	46.2 <sub>F</sub> , 100 <sub>T</sub>	0	0	23	0	23	NT	53.8	61.5	7.7	7.7	NT	Cuba/Blanco et al., 2006	
	220	56 <sub>F</sub> , 74 <sub>T</sub>	38	3	3	3	9	3	13	49	42	4	65	Slovakia/Vu-Khac et al., 2007	
	30	17 <sub>F</sub> , 26.7 <sub>T</sub>	0	0	0	0	0	0	17	0	0	0	0	26	Zimbabwe/Madoroba et al., 2009
(4-21 d.)	196	32 <sub>T</sub>	9.7	7.7	0.5	7.7	8.8	NT	9.2	15.8	16.3	8.8	NT		
ND&PWD															
(0-35 d.)	83	79.3 <sub>F</sub> , 90 <sub>T</sub>	45.8	18.4	26.3	5.3	26.3	10.5	80.7	44.6	45.8	15.7	NT	Japan/Katsuda et al., 2006	
(2-72 d.)	562	34	2.3	2	2.5	2.8	5.3	7.1	7.3	4.8	7.8	5.2	13.9	South Korea/Kim et al., 2010b	
PWD															
(4-6 w.)	372	29 <sub>F</sub> , 80.1 <sub>T</sub>	19.1 <sup>A</sup>	2.1 <sup>A</sup>	1.1 <sup>A</sup>	2.9 <sup>A</sup>	2.7	NT	12.4	4.3	57.3	20.4	NT	Poland/Osek, 1999	
Healthy	46	3.5 <sub>F</sub> , 8.7 <sub>T</sub>	0 <sup>A</sup>	4.3 <sup>A</sup>	0 <sup>A</sup>	0 <sup>A</sup>	2.2	NT	2.2	0	2.2	4.3	NT		
(4-10 w)	215	50.2 <sub>F</sub> , 7.2 <sub>T</sub>	9.8 <sup>A</sup>	10.7 <sup>A</sup>	15.8 <sup>A</sup>	9.8 <sup>A</sup>	25.6 <sup>A</sup>	NT	74.4	14	2.3	8.8	NT	China/Chen et al., 2004	
(>30 d.)	23	82.6 <sub>F</sub> , 100 <sub>T</sub>	0	0	0	0	82.6	NT	65.2	73.9	4.3	78.2	NT	Cuba/Blanco et al., 2006	
-	101	60 <sub>F</sub> , 77 <sub>T</sub>	19	0.9	5	0.9	35	0.9	26	46	20	5	64	Slovakia/Vu-Khac et al., 2006	
-	304	58 <sub>F</sub>	37.1	0.3	0	0.3	19.7	0.7 <sup>D</sup>	15.8	41.8	33.2	9.9	20	U.S.A./Zhang et al., 2007	
-	100	100	44	30	25	32	38	NT	40	47	71	3	NT	Brazil/Vidotto et al., 2009	
PWD&ED															
-	230	40.9	10	1.7	4.3	0.8	18.3	NT	27.5	15.2	8.7	15.2	NT	South Korea/Kwon et al., 2002	
(4-8 w.)	219	85.4 <sub>F</sub> , 87.2 <sub>T</sub>	44.7	0	0.9	0	39.3	1.4	26.5	7.6	61.6	16.4	65.8	Denmark/Frydendahl, 2002	
-	240	100	3.7	0	0	0	26.2	28.3	14.5	9.1	10.8	35	NT	China/Cheng et al., 2006	

ND, neonatal diarrhea; PWD, post-weaning diarrhea; ED, edema disease; n, tested *E. coli* isolates; VF, virulence factor; <sub>T</sub> and <sub>F</sub>, positive for at least one of the tested toxin or adhesin genes respectively; <sup>A</sup>, identified by the agglutination test; <sup>D</sup>, Zhang et al. (2007) also reported AIDA-I (15.5%) and PAA (34.5%); NT, not tested; d., days old; w., weeks old.

***Clostridium perfringens***

*C. perfringens* is a Gram-positive anaerobic bacterium that is able to form spores. It is widespread in the environment (e.g. in soil and sewage) and is commonly found in the intestine of animals. *C. perfringens* strains are classified into five toxinotypes (A, B, C, D, and E) according to the production of alfa ( $\alpha$ ), beta ( $\beta$ ), epsilon ( $\epsilon$ ) and or iota ( $\iota$ ) toxins which are crucial in the pathogenesis of clostridiosis (Petit *et al.*, 1999); besides, they can also produce a pore-forming enterotoxin called *C. perfringens* enterotoxin (CPE; McClane, 1996) and a cytotoxic  $\beta$ 2-toxin (Garmory *et al.*, 2000) which are usually tested for subtyping.

Alfa and  $\beta$ -toxigenic strains have been associated with swine diarrhea worldwide (Morin *et al.*, 1983; Niestrath *et al.*, 2002; Yaeger *et al.*, 2002; Das *et al.*, 2009; Cruz *et al.*, 2010).

*C. perfringens* type-C produce  $\alpha$  and  $\beta$  toxins and cause hemorrhagic, often fatal, necrotic enteritis in young piglets. The disease is most frequent in 3-day-old piglets, but it can appear in the first 12 hours of life (Songer and Meer, 1996; Songer and Uzal, 2005). This strain is rarely found in the intestine of healthy piglets. It is important to remark that piglet-piglet transmission occurs, and spores persist in the environment as they are resistant to heat, disinfectants, and ultraviolet light. The main source of infection in a piggery is the intestine of the sow (Songer, 1996). Type C clostridiosis can occur epidemically in non-vaccinated herds, and prevalence in affected litters can reach 100% with mortality close to 100% (Songer and Uzal, 2005).

The basis of the pathogenicity of *C. perfringens* type A strains ( $\alpha$ -toxigenic) frequently isolated from pigs with enteritis has not been clearly established but necrotic intestinal lesions have been experimentally induced by inoculation of *C. perfringens* type A culture supernatant (Songer, 1996). The enteropathogenicity of these strains might result from high levels of  $\alpha$ -toxin production, from molecular variants that are more stable to protease digestion or are more active, from different host sensitivity to  $\alpha$ -toxin (Ginter *et al.*, 1996), or due to the side production of  $\beta$ 2-toxin (Garmory *et al.*, 2000; Hendriksen *et al.*, 2006) or the *C. perfringens* enterotoxin.

The  $\alpha$ -toxin is a phospholipase C sphingomyelinase that hydrolyzes phospholipids (e.g. lecithin) and promotes membrane disorganization, resulting in blood vessel contraction, increased vascular permeability, platelet aggregation and myocardial dysfunction, all of which contribute to local and systemic clinical manifestations (Bunting *et al.*, 1997; Naylor *et al.*, 1998). The  $\beta$ 1- and  $\beta$ 2-toxins induce hemorrhagic necrosis of the intestinal mucosa (Gibert *et al.*, 1997). Although these toxins are cytotoxic (Gibert *et al.*, 1997), their mode of action has not yet been completely elucidated. The fact that  $\beta$ 1-toxin displays a significant homology at the amino acid level with  $\alpha$ -toxin, and leucocidin of *Staphylococcus*

*aureus* which form multimers and pores in eukaryotic cell membranes, suggests that  $\beta$ 1-toxin has a similar mode of action (Hunter *et al.*, 1993). Gurtner *et al.* (2010) confirmed that  $\beta$ 1-toxin causes disruption of the actin cytoskeleton of endothelial cells. Also, *C. perfringens* secretes a variety of hydrolytic enzymes that degrade extracellular substrates and components resulting from cell lysis. It is possible that these enzymes act synergistically with membrane-damaging toxins during cell disruption (Petit *et al.*, 1999). Zeng *et al.* (2011) developed and recommended the application of recombinant fusion toxoids as good vaccine candidate against the  $\alpha$ ,  $\beta$ 1, and  $\beta$ 2 clostridial toxins.

Testing for *C. perfringens* toxins by enzyme-linked immunosorbent assay (ELISA) is a reliable test (Naylor *et al.*, 1997; Niestrath *et al.*, 2002), i.e. the commercially available Bio-X ELISA kit (Bio-X Diagnostics, Marche-en-Famenne, Belgium) that detects the  $\alpha$ -,  $\beta$ - and  $\epsilon$ -toxins in intestinal contents or culture supernatants. Genotyping of *C. perfringens* has simplified the routine diagnosis. Multiplex-PCR for detecting fragments of genes encoding toxins enables the diagnostician to screen larger numbers of samples with higher accuracy and greatly reduces the amount of false-negative results (Das *et al.*, 2009; Baker *et al.*, 2010).

In summary, the clinics and the differential diagnosis with other enteropathogens is very important to consider when implicating *C. perfringens* as primary agent. Toxinotype A is widespread in the intestines of pigs worldwide (Table 6) and its association with diarrhea has to be carefully analyzed in every outbreak. Type C strains frequently cause hemorrhagic necrotic enteritis in young pigs (Petit *et al.*, 1999).

**Table 6.** Reports of surveys assessing association of *C. perfringens* with swine diarrhea worldwide.

Age groups	Tested	%	Associated toxins					Assay	Country/Reference
			$\alpha$	$\beta$ 1	$\beta$ 2	$\epsilon$	CPE		
1-15 d.	749	0.4	<sup>NE</sup> +	<sup>NE</sup> +				Clinics, histopathology general bacteriologic procedures	Canada/Morin <i>et al.</i> , 1983
1-3 d.	13	100 <sup>M</sup>	+					Based on anamnesis, differential diagnose, histopathology, general bacteriologic procedures, inoculation bioassay and reverse passive latex agglutination assay	U.S.A./Collins <i>et al.</i> , 1989
Suckers	33	87.9	+	-	+	-	-	General bacteriologic procedures, PCR	U.S.A./Garmory <i>et al.</i> , 2000
	33	12	+	+	+	-	-		
	7 <sup>ND</sup>	0	-	-	-	-	-		
Suckers	10 <sup>DN</sup>	50 <sup>LS</sup>	+	-	NT	-	NT	DAS-ELISA	Germany/Niestrath <i>et al.</i> , 2002
1-7 d.	100	6	<sup>C</sup> +	<sup>C</sup> +				Necrotizing intestinal lesions in association with large, gram-positive bacilli lining necrotic villus remnants. Dense grow of <i>C. perfringens</i>	U.S.A./Yaeger <i>et al.</i> , 2002
Suckers	220 <sup>A</sup>	90.9	+	-	+	-	-	PCR (during this survey <i>C.p.</i> isolates were only tested for <i>cpb2</i> gene which codes for $\beta$ 2 toxin)	U.S.A./Bueschel <i>et al.</i> , 2003
	36 <sup>C</sup>	97.2	+	+	+	-	-		
	9 <sup>C.p. ND</sup>	11.1	-	-	+	-	-		
Suckers	153	0	-	-	NT	-	NT	General bacteriologic procedures, PCR	Japan/Katsuda <i>et al.</i> , 2006
Weaners	116	0	-	-	NT	-	NT		
12-14 m.	146	2.1 <sup>T</sup>	+	-	+	-	-	General bacteriologic procedures, PCR	Malaysia/Das <i>et al.</i> , 2009
2-3 m.	96	2.1 <sup>T</sup>	+	-	+	-	-		
1-7 d.	30	33.3	+	-	+	-	-	General bacteriologic procedures, PCR	Brazil/Cruz <i>et al.</i> , 2010
Suckers	333	89.8	+	-	NT	-	-	General bacteriologic procedures, PCR	U.S.A./Baker <i>et al.</i> , 2010

<sup>NE</sup>, based on findings of necrotic enteritis which is associated with *C. perfringens* C; <sup>M</sup>, morbidity close to 100% and 13 piglets (8 sick and 5 healthy) were selected for *C. perfringens* study; <sup>\*</sup>, in 79% of *C. perfringens* A; <sup>ND</sup>, non diarrheic control piglets, <sup>LS</sup>, mixed with *I. suis*, and 1/5 positive piglets had necrotic enteritis; <sup>C</sup>, *C. perfringens* type C; <sup>A</sup>, *C. perfringens* type A; <sup>C.p. ND</sup>, *C. perfringens* isolated from non diarrheic piglets; <sup>T</sup>, *C. perfringens* was isolated from pigs which died after tetracycline resistant acute diarrhea; <sup>DN</sup>, diarrheic and non-diarrheic pigs.

**Transmissible gastroenteritis virus and porcine epidemic diarrhea virus**

TGEV belongs to the *Coronaviridae* family. The virus contains a single-stranded genomic RNA and only one serotype is known (Kemeny, 1976). Three major structural proteins described for coronaviruses are the spike glycoprotein (S; 180–200 kDa), the membrane protein (M; 21–30 kDa), and the nucleoprotein (N; 45–50 kDa) (Spaan *et al.*, 1988). The S protein is the most interesting protein from an antigenic and immunogenic point of view (Delmas *et al.*, 1986; Jiménez *et al.*, 1986; Torres *et al.*, 1995). Four antigenic sites (C, B, D, and A) were mapped on the S protein starting from the N-terminal end and antibodies against them can be found in the serum of TGEV-infected pigs (Correa *et al.*, 1990). The spike glycoprotein initiates infection by binding to the enterocytes via pAPN, which has been identified as a coronavirus receptor (Delmas *et al.*, 1992; Hansen *et al.*, 1998). PEDV, another coronavirus, also binds specifically to pAPN and this binding can be inhibited by anti-pAPN antibodies (Oh *et al.*, 2003; Li *et al.*, 2007). TGEV shows also sialic acid binding activity, perhaps providing a second binding site that may account for the enteropathogenicity of the different strains (Schwegmann-Wessels *et al.*, 2002; Schwegmann-Wessels and Herrler, 2006).

Three swine coronaviruses are pathogenically or antigenically related: TGEV, PEDV, and porcine respiratory coronavirus (PRCV). TGEV and PRCV cross-react serologically and are very closely related only differing in a deletion mutation of 224–227 amino acids in the Spike protein S of PRCV in comparison with TGEV. PRCV cannot be distinguished from enteropathogenic strains of TGEV by a virus neutralization test (Callebaut *et al.*, 1988). Indeed, infection with PRCV induces the production of antibodies able to neutralize both TGEV and PRCV at the same titer (Pensaert *et al.*, 1986). Previous PRCV infections in piglets and or sows, or concurrent TGEV/PRCV infections influence and change the pathogenesis of TGEV, thereby reducing the severity of disease (Cox *et al.*, 1993; Kim *et al.*, 2000). As a consequence, the presence of PRCV in Europe reduced the incidence and severity of epidemic TGE (Pensaert *et al.*, 1993). Therefore, a low prevalence (0.9%) of TGEV infection compared with PEDV in South Korea could be due to the high prevalence of PRCV (Chae *et al.*, 2000). Also in the United States and Japan, a decrease in TGE incidence has been reported in areas with a high prevalence of anti-PRCV antibodies (Yaeger *et al.*, 2002; Miyazaki *et al.*, 2010). In TGEV- and PRCV-seronegative herds, however, TGE remains a major cause of sickness and death in piglets (Barrera *et al.*, 2005; Saif and Sestak, 2006).

Characteristics of TGE acute form are a short incubation period, diarrhea, vomiting, and dehydration. Mortality approaches 100% in newborn piglets, but decreases with age. Sows infected shortly after parturition may be severely affected by diarrhea, hypogalactia, and agalactia (Djurickovic *et al.*, 1969; Saif and Sestak, 2006). The first outbreaks of TGE reported in February 2003 in Cuba were a classic

example of this form. On affected farms, 100% of recently farrowed sows and their litters had diarrhea. The clinical signs in newborns included very liquid and fetid, yellowish feces and vomiting, leading to serious dehydration. At the onset of the disease, sows showed a lack of appetite followed by vomiting and agalactia, but all recovered. The weaned and fattening pigs of these farms presented severe clinical signs, although only 8% lethality was reached. The disease spread rapidly to other farms over the island. In the Havana province, 15 outbreaks affecting 23.201 animals, caused 10.547 deaths, and 5.256 animals had to be slaughtered (Barrera *et al.*, 2005; IMV, 2003).

The clinical signs of TGE are usually milder when TGEV is introduced into seropositive farms, or when TGEV infects less susceptible animals, such as sows or finisher pigs in seronegative farms. Endemic TGE is limited to seropositive herds and diarrhea can occur in pigs from 6 days old until 2 weeks after weaning, and the mortality varies from 10-20% or even less. During this presentation TGE is difficult to differentiate from rotavirus or enteric colibacillosis (Saif and Sestak, 2006). TGEV infection therefore occasionally goes undiagnosed. Risk for TGE was greater in herds with more than 50 breeding pigs than in smaller ones. ( $p < 0.01$ ; Cubero *et al.*, 1993).

Traditional methods to confirm TGEV infection are based on clinical signs, microscopic lesions, virus isolation, fluorescent antibody test, electron microscopy or immunohistochemistry, and an enzyme immunofiltration assay (Phillips and Westerman, 1991; Saif and Sestak, 2006).

The appearance of the closely related PRCV complicated the diagnosis. However, the absence of 2 antigenic sites (B and C) in the S protein of PRCV as a consequence of the deletion of 224–227 amino acids, was the basis for its differentiation from TGEV (Rasschaert *et al.*, 1990) and a number of ELISA's based on MAb's directed against the antigenic determinant in the S protein that is deleted in PRCV have been developed to differentiate TGEV and PRCV (Jabrane *et al.*, 1992; Sestak *et al.*, 1999; López *et al.*, 2009). PEDV is not serologically related to TGEV or PRCV. Jung *et al.* (2003) recommended the multiplex RT-PCR as a monitoring and diagnostic tool to confirm the presence of PEDV and TGEV in formalin-fixed, paraffin-embedded tissues.

Apart from the epidemic acute outbreaks, where the TGEV prevalence and mortality is close to 100%, many studies focused on the prevalence of antibodies or fecal shedding of TGEV and PEDV worldwide (Table 7).

## Rotavirus

Rotaviruses are RNA viruses of the *Reoviridae* family, which replicate mainly in the small intestine and spread mainly via the fecal-oral route. The rotaviral genome consists of 11 segments of double stranded RNA coding for six viral structural and six non-structural proteins (Matthijnssens *et al.*, 2008). The viral particles are composed of a capsid which contains three layers of viral proteins (VP) described as the outer (glycoprotein VP7 and non-glycosylated protease-sensitive VP4), intermediate (VP6), and inner (VP2) layers. VP1-3 are the core proteins (Yuan *et al.*, 2006).

Rotavirus A is common in piggeries and is more prevalent in piglets from 1 to 3 weeks old and soon after weaning (Atii *et al.*, 1990). In pigs positive for rotavirus A, Halaihel *et al.* (2010) found that diarrhea is more likely to occur in the ones younger than 8 weeks old. The prevalence of infection and disease is favored by predisposing factors such as immunological quality and quantity of colostrum intake, nutrition and the immune status of the sows, poor sanitary conditions in pens and around the piggeries, and high population density (Steel and Torres-Medina, 1984; Atti *et al.*, 1990; Zijlstra *et al.*, 1999; Barreiros *et al.*, 2003). Additionally, the continuous exposure of pigs to rotaviruses is favored by their resistance in the environment as they maintain infective for 32 months at 10°C in stool specimens (Ramos *et al.*, 2000).

Porcine rotaviruses are antigenically diverse. The two outer capsid proteins, VP7 (G genotype) and VP4 (P genotype), independently elicit serotype-specific neutralizing immune responses that are believed to play an important role in protection against recurrent infections (Santos and Hoshino, 2005). Based on the differences in nucleic acid sequences of the outer capsid VP7 and VP4 encoding genes, 23 G and 31 P genotypes of rotavirus A have been described (Abe *et al.*, 2009; Ursu *et al.*, 2009). In pigs, 10 G types (1-6, 8-10 and 11) and 7 P types (5-8, 13, 9 and 23) of rotavirus A have been associated with diarrhea (Martella *et al.*, 2001; Barreiros *et al.*, 2003). Group A rotavirus infection has been recognized to occur in both enzootic and epizootic forms of swine diarrhea, resulting in serious economic losses in the suckling and weaning piglet population of commercial piggeries (Martella *et al.*, 2007; Kim *et al.*, 2010a).

The rotaviral replication in the villous epithelial cells cause malabsorption due to loss of absorptive cells and villous atrophy, which is the more accepted mechanism of rotavirus induced diarrhea in pigs (Greenberg and Estes, 2009). Rotavirus also induces an intestinal inflammatory response that may contribute to a secretory-type diarrhea (Zijlstra *et al.*, 1999), evokes intestinal fluid and electrolyte secretion by activation of the nervous system in the intestinal wall as evidenced by the use of four enteric nervous system inhibitor drugs (Lundgren *et al.*, 2000). The rotaviral non-structural protein 4



induces diarrhea in a similar way as *E. coli* STa by activating guanylate cyclase (Ball *et al.*, 1996; Kavanagh *et al.*, 2010).

Colostrum-deprived piglets inoculated with rotavirus 24 hours after birth develop profuse diarrhea with high mortality (63%). Interestingly, when these piglets were re-grouped with their colostrum-fed litter mates, the later got infected and developed diarrhea with a mortality of only 8% and decreased weight gains. Piglets recovered from rotavirus can excrete the virus up to 21 days post-infection and the severity of the infection is age dependent, and piglets inoculated at 5-14 days old developed diarrhea but suffered a low mortality rate (Svensmark *et al.*, 1989a). The highest rotavirus prevalence occurs most frequently in litters from primiparous sows (Svensmark *et al.*, 1989b). Sows are easily infected with rotavirus by contact with an infected litter, but they do not show signs of diarrhea, meaning that they are an important source of infection to their offspring (Svensmark *et al.*, 1989a). Moreover this contact leads to a better transfer of anti-rotavirus immunoglobulins through colostrum in multiparous sows.

Field studies diverge concerning the pathogenicity of porcine rotaviruses. Rotavirus A was identified in feces of 43 out of 96 (44.8%) piglets suffering from acute gastroenteritis, while none of 41 non-diarrheic piglets were positive ( $P < 0.01$ ) in Nigeria (Atii *et al.*, 1990). Rotavirus A was the most prevalent enteropathogen associated with diarrhea in piglets from Japan (Katsuda *et al.*, 2006). In contrast, in Argentina, Parra *et al.* (2008) reported that only 5/30 1-45-day-old pigs positive to Rotavirus had diarrhea. These statements mean that in every geographic area or management system rotavirus can behave different, for example, in South Korea rotavirus A shedding in pig feces increases during the summer season (Kim *et al.*, 2010a).

The clinical signs of rotaviral enteritis are not pathognomonic. Worldwide, electron microscopy, immune electron microscopy, immunohistochemistry, immunofluorescence, ELISA, virus isolation, latex agglutination, dot blot hybridization, RNA electrophoretotyping, and RT-PCR have been employed to confirm rotaviral diagnosis (Yuan *et al.*, 2006). Electron microscopy is a highly specific method (Rodák *et al.*, 2009). ELISA is a reliable technique for the detection of rotaviral antigens in fecal samples or intestinal contents and is specially standardized and available for the diagnosis of porcine rotavirus A (Ingezim Rota DAS 1.1.RT.K.2 DAS-ELISA) with similar results as electron microscopy (Goyal *et al.*, 1987; Rodák *et al.*, 2004). Electrophoretotyping of viral RNA isolated from feces is a useful technique, but it has to be confirmed by serological or nucleic acid-based methods (Mattion *et al.*, 1989; Rácz *et al.*, 2000). A variety of serological test have been performed and the ones that study the immunoglobulin isotypes are of help when detecting active or recent infections (Yuan and Saif, 2002). Multiplex RT-PCR offers a

more sensitive and reliable method for G and P genotyping of group A rotavirus (Martella *et al.*, 2001; Barreiros *et al.*, 2003). Chizhikov *et al.* (2002) recommended the oligonucleotide microarray hybridization for the identification of the G genotypes of all rotavirus strains combining RT-PCR and DNA-DNA hybridization. On swine stool specimens, Kang *et al.* (2007) suggested the use of immunochromatographic assays that can separately and accurately detect porcine rotaviruses, TGEV and PEDV.

In pigs, multiple rotavirus serogroups and serotypes have been detected worldwide (Table 8).

**Table 7.** Occurrence of TGEV and PEDV detected by assays applied to serum and feces of pigs worldwide.

Groups & age	Tested	%		Assay	Country/Reference
		TGEV	PEDV		
Serum					
-	665	0	-	VN	U.S.A./Woods <i>et al.</i> , 1990
Sows	-	0.6	-	Competitive ELISA	Great Britain/Brown and Paton, 1991
-	229	89.9	-	ELISA	U.S.A./Phillips and Westerman, 1991
Breeding pigs	6000	1.27	-	ELISA	Spain/Cubero <i>et al.</i> , 1993
Pigs in positive farms	-	5-60	-	ELISA	Spain/Cubero <i>et al.</i> , 1993
Pigs in abattoirs	5.337	0	-	ELISA	South Africa/Williams <i>et al.</i> , 1994
>7 months feral swine	117	0	-	IFAT	U.S.A./Saliki <i>et al.</i> , 1998
Wild boars	134	1	-	IFAT	Czech Republic/Sedlak <i>et al.</i> , 2008
Wild boars	178	0	0	ELISA	Slovenia/Vengust <i>et al.</i> , 2006
-	263	12.5	-	ELISA	Japan/Miyazaki <i>et al.</i> , 2010
Diarrheic feces					
1-15 d.	749	59.2	-	FAT and EM	Canada/Morin <i>et al.</i> , 1983
0-21 d.	1258	0.9	50.4	RT-PCR	South Korea/Chae <i>et al.</i> , 2000
1-7 d.	33	0			
8-14 d.	50	4			
15-21 d.	19	10.5		EM	Germany/Wieler <i>et al.</i> , 2001
22-28 d.	16	0			
36-42 <sup>w</sup> d.	31	51.6			
1-7 d.	100	6	-	FAT and IHC	U.S.A./Yaeger <i>et al.</i> , 2002
1-14 d.	157	2.5	13.5	RT-PCR	South Korea/Song <i>et al.</i> , 2006
1-21 d.	153	0	0	RT-PCR	Japan/Katsuda <i>et al.</i> , 2006
22-35 <sup>w</sup> d.	116	0	0	RT-PCR	Japan/Katsuda <i>et al.</i> , 2006
2-28 d.	68	52	41.8	IC	Czech Republic/Czanderlova <i>et al.</i> , 2010

<sup>w</sup>, weaned; d., days old.

**Table 8.** Worldwide occurrence (%) of Rotavirus in feces of diarrheic pigs during last decade.

Age group	Tested	%	A	B	C	Assay	Country/Reference
1-60 d. <sup>DN</sup>	165	35.3	100	NT	NT	PAGE, ELISA, RT-PCR	Brazil/Rácz <i>et al.</i> , 2000
1-7 d.	100	42	100	NT	NT	ELISA	U.S.A./Yaeger <i>et al.</i> , 2002
1-7 d.	33	0	NT	NT	NT	EM	Germany/Wieler <i>et al.</i> , 2001
8-14 d.	50	2					
15-21 d.	19	5.3					
22-28 d.	16	25					
36-42 d.	31	0					
< 7 d.	19	53	100	NT	NT	PAGE	Brazil/Barreiros <i>et al.</i> , 2003
8-21 d.	20	60					
> 21 d.	60	62					
1-7 d.	60	81.7	60-70*	NI	NI	RT-PCR	Japan/Katsuda <i>et al.</i> , 2006
8-14 d.	44	61.4					
15-21 d.	46	60.9					
22-28 d.	40	77.5					
29-35 d.	46	43.5					
1-14 d.	157	10.8	100	NT	NT	RT-PCR	South Korea/Song <i>et al.</i> , 2006
Piglets	175	22.3	100	NT	NT	ELISA in fecal samples	Thailand/Khamrin <i>et al.</i> , 2007
1-3 m.	102	-	71.5	NT	31.3	RT-PCR	Italy/Martella <i>et al.</i> , 2007
	86**		81.3		25.5	EM, RT-PCR	
< 45 d. <sup>DN</sup>	905	3.3	100	NT	NT	PAGE, ELISA, RT-PCR	Argentina/Parra <i>et al.</i> , 2008
< 3 w.		50	100	NT	NT	RT-PCR	Slovenia/Steyer <i>et al.</i> , 2008
3-10 w.	-	35.7					
> 10 w.		46.2					
Piglets	476	18.9	100	NT	NT	ELISA (fecal VP6) PCR	Czech Republic/Rodák <i>et al.</i> , 2009
	21***	76.2	NT	14.3	76.2		
0-4 w. <sup>DN</sup>	45	26.7	100	NT	NT	RT-PCR	Spain/Halaihel <i>et al.</i> , 2010
4-8 w.	19	47					
8-16 w.	83	14.5					
16-24 w.	63	6.3					
3-70 d.	475	38.3	100	NT	NT	RT-PCR, nested PCR	South Korea/Kim <i>et al.</i> , 2010a
1-7 d.	30	13.3	NI	NI	NI	PAGE	Brazil/Cruz <i>et al.</i> , 2010
2-28 d.	98	30.6	100	NT	NT	IC	Czech Republic/Czanderlova <i>et al.</i> , 2010

A, B and C, groups of Rotavirus (%); <sup>DN</sup>, diarrheic and non-diarrheic pigs. \*, from the positives; \*\*, positives by EM; \*\*\*, samples EM<sup>+</sup>/VP6 ELISA<sup>+</sup>; NI, no information; NT, not tested; d., days old; w., weeks old; m., months old.

***Cryptosporidium***

The contribution of *Cryptosporidium* to piglet diarrhea should not be ruled out as experimental infections have revealed its potential pathogenicity (Suárez-Luengas *et al.*, 2007). In two series of experimental infections to compare the infectivity of the *Cryptosporidium* pig genotype I (*C. suis*) and *C. parvum* in piglets, Enemark *et al.* (2003ab) reported that infection with *C. suis* caused no or very mild clinical signs whereas *C. parvum* provoked diarrhea. The species or genotypes of *Cryptosporidium* can be distinguished on the basis of the small subunit ribosomal RNA gene sequence (Xiao *et al.*, 1999).

It has been suggested that the withdrawal of protective maternal antibodies and the stress of weaning renders weaners particularly susceptible to infection (Hamnes *et al.*, 2007). As referred by Sanford (1987), Izumiyama *et al.* (2001) and Ryan *et al.* (2003), the higher positivity of pigs to *Cryptosporidium* occurs in the post-weaning period and Chen and Huang (2007) reported that the youngest pig excreting oocysts was 3 days old, and the oldest pig was 3.5 years old. Sows could maintain the parasite life cycle in piggeries, even in the absence of piglets, because stress associated with pregnancy and hormonal changes may increase susceptibility to infection (Zintl *et al.*, 2007). Recently Chen and Huang (2007) showed that *Cryptosporidium* isolated from pigs in eastern China belonged to the *C. parvum* 'mouse' genotype indicating that rodents as sows play a role in swine cryptosporidiosis epidemiology as reservoirs.

The specific mechanism by which *C. parvum* induces diarrhea has not been identified. Many molecules like CSL, a 1.300-kDa conserved apical complex glycoprotein, are crucial for the adherence of sporozoites and merozoites to the gut epithelium. *Cryptosporidium* gets internalized into the enterocytes and locates in a membrane-bound compartment on the apical surface. Therefore, direct damage of the brush border membrane, which provokes structural villous atrophy, and parasite derived molecules cause malabsorption and a sequence of biochemical/immunological changes like increase of secretogenic prostaglandins leading to diarrhea (Okhuysen and Chappell, 2002). Worldwide many studies have associated this *Apicomplexa* with swine diarrhea. Chen and Huang (2007) found in China a positive correlation between oocyst positivity and diarrhea, and *Cryptosporidium* prevalence was not significantly different between pre and post-weaned piglets. Hamnes *et al.* (2007) reported in Norway a significantly higher prevalence of diarrhea among the *Cryptosporidium* positive litters than among the negative ones. In Serbia it was reported that all *Cryptosporidium* positive nursing piglets had diarrhea, whereas in post-weaned piglets and adults *Cryptosporidium* infection was asymptomatic (Mišić *et al.*, 2003). In Mexico, Nevarez *et al.* (1997) implicated *Cryptosporidium* as infectious cause of enteritis in three 10-week-old weaned piglets from a crowded herd with bad sanitation and subjected to massive

treatment with several antibiotics. Conversely, in 20 out of 48 diarrheic pigs, *C. parvum* was not determined to be the only cause of diarrhea, suggesting that it may act in concert with other agents (i.e. *E. coli*, *I. suis*, *Salmonella*, adenovirus) to induce or exacerbate clinical disease and suggesting a strong possibility of subclinical infection (Sanford, 1987). Similarly, Quilez *et al.* (1996), Ryan *et al.* (2003), Guselle *et al.* (2003) and Vítovec *et al.* (2006) did not associate *Cryptosporidium* infection with swine diarrhea. Enemark *et al.* (2003a) found that a mixed infection of rotavirus and *Cryptosporidium* caused a dramatic aggravation of diarrhea. Vítovec *et al.* (2006), Maddox-Hyttel *et al.* (2006), and Johnson *et al.* (2008a) reported that *Cryptosporidium* occurs more frequently in weaned pigs than in sucking piglets.

Table 9 shows the fecal occurrence of *Cryptosporidium* in pigs raised worldwide during the last decade and several diagnostic techniques, which have been applied for its identification.

### ***Isospora***

The coccidian protozoa *Isospora* is one of the most common enteropathogens amongst suckling piglets in intensive pig production units, and is an important enteropathogen to be surveyed and controlled as the infection is associated with diarrhea (Harleman and Meyer, 1984; Vitovec and Koudela, 1990; Niestrath *et al.*, 2002).

Piglets get infected by the ingestion of infectious oocysts from which the sporozoites emerge and invade the epithelial cells of the small intestine. Later, the course of the life cycle phases merogony (trophozoites, meronts, and merozoites), gametogony (macro and microgametocytes) and sporogony (immature oocyst) contribute to the destruction of enterocytes and to the colonization of the intestinal epithelium. Around four to five days after infection, piglets frequently have pasty to watery yellowish or grayish diarrhea for 3-7 days, dehydration, and weight loss (Mundt *et al.*, 2006). Clinical signs of isosporosis mostly appear in suckling piglets in the first three weeks of life with high morbidity and low mortality (Mundt *et al.*, 2003).

Lindsay *et al.* (1992) and Otten *et al.* (1996) reported that isosporosis can occur in any farm, independently of their size or management system. Niestrath *et al.* (2002) reported that the rearing conditions and hygienic status on swine farms influence the prevalence of coccidian infections in piglets.

Traditionally, sows have been thought to be the main source of *Isospora* in a farm, but Karamon *et al.* (2007) in Poland demonstrated no correlation between the excretion of oocysts by the sows and the infection of piglets born from them. The farrowing pens contaminated with oocysts excreted by previous litters is considered to be the main source of infection, showing the importance of good sanitation (oocysts are resistant to most of disinfectants) and the use of self-cleaning floors in farrowing

facilities for isosporosis prevention (Lindsay *et al.*, 1989; Niestrath *et al.*, 2002; Karamon *et al.*, 2007). In that context, Niestrath *et al.* (2002) commonly found *I. suis* in farms which employed straw bedding.

The general laboratory methods which included staining and or visualization of oocyst morphology by light microscopy have been employed to identify *Isospora* but their sensitivity is negatively influenced by the high fat content of piglets feces and because oocysts are not always excreted or are excreted intermittently. Dauschies *et al.*, 2001 reported that autofluorescence microscopy is more sensitive than bright field microscopy for the detection of *I. suis* oocysts after flotation or in direct smears. Recently a PCR–RFLP assay based on the rDNA ITS-1 region has been developed to identify *Isospora* (Samarasinghe *et al.*, 2007; Johnson *et al.*, 2008b). Table 10 shows the assays mostly employed for the identification of porcine *Eimeriidae* and their prevalence worldwide.

**Table 9.** Worldwide reported fecal occurrence (%) of *Cryptosporidium* in pigs during last decade.

Groups	Tested	%	<i>C. suis</i>	<i>C. Gen II</i>	<i>C. parvum</i>	<i>C. muris</i>	Assay	Country/Reference
1 m.	213	32.3			x		Formol-ether concentration	Japan/Izumiyama <i>et al.</i> , 2001
3 m.	19	42.1			x		Immunofluorescent staining (A)	
6 m.	252	0.4			x			
Suckers and weaners	287	1.4			x		Carbol-fuchsin staining	Germany/Wieler <i>et al.</i> , 2001
Adults	589	10.5			x		Acid-fast staining	South Korea/Yu and Seo, 2004
< 5 w.	3368	5.7					Aniline-carbol-methyl violet staining	Czech Republic/Vítovec <i>et al.</i> , 2006
> 6 w.	835	24.1					Sheather's flotation, PCR	
Sows	135	0					Immunofluorescent staining	Denmark/Maddox-Hyttel <i>et al.</i> , 2006
< 1 m.	488	6						
1-4 w.	504	71						
Sows	245	4						
1-7 d.	60	0					Flotation <sup>b</sup>	Japan/Katsuda <i>et al.</i> , 2006
8-14 d.	44	0						
15-21 d.	46	0						
22-28 d.	40	2.5						
29-35 d.	46	28.3						
4-10 w.	127	15						
10-24 w.	121	7.4						
6-12 m. gilts	15	6.7					Auramine phenol staining	Ireland/Zintl <i>et al.</i> , 2007
4 y. sows	75	13.3			x		Sheather sugar flotation, PCR	
3-4 y. boars	4	0						
1-2 <sup>w</sup> m.	75	30.7	x	x				
2-6 m.	42	11.9	x	x			Formol-ether concentration, PCR	Spain/Suárez-Luengas <i>et al.</i> , 2007
Sows	25	16	x					
Suckers	183	10.4			x		Acid-fast staining, PCR	China/Chen and Huang, 2007
Weaners	317	12.9						
Fatteners	536	13.8						
Boars	91	8.8						
11d. - 3 w.	123	10.6	x				PCR	Australia/Johnson <i>et al.</i> , 2008a
4 w. - 6 m.	156	32.7	x	x				
Sows	10	0						

**Table 9** (continuation). Worldwide reported fecal occurrence (%) of *Cryptosporidium* in pigs during last decade.

Groups	Tested	%	<i>C. suis</i>	<i>C. Gen II</i>	<i>C. parvum</i>	<i>C. muris</i>	Assay	Country/Reference
Suckers	116	21.8	x				Aniline-carbol-methyl violet staining, PCR	Czech Republic/Kvác <i>et al.</i> , 2009
Weaners	131	29	x	x		x		
Fatteners	123	17	x	x		x		
Sows	40	2.5						
<6 m.	105	11.4						
>6 m.	133	6.7						
0-3 w.	20	20	x*	x*	x*		Formol-ether concentration and acid-fast staining	Turkey/Uysal <i>et al.</i> , 2009
4-7 w.	36	38.9					Immunofluorescent staining, PCR	UK/Featherstone <i>et al.</i> , 2010**
8-13 w.	93	48.4						
14-24 w.	119	40						
>24 w.	39	30.8						

<sup>w</sup>, weaned pigs; d., days old; w., weeks old; m., months old; y., years old; \*, information from the overall sample; <sup>D</sup>, all piglets had diarrhea; (A), all pigs were asymptomatic; \*\*, *Cryptosporidium* pig genotype II was detected in 25 out of 39 (64.1%) sequenced isolates. *C. parvum* was identified in eight (20.5%) isolates, and *C. suis* was identified in six (15.4%) isolates.



**Table 10.** Worldwide reported fecal occurrence (%) of *Eimeriidae* in pigs during last decade.

Group & age	Tested	<i>I. suis</i>	<i>Eimeria</i> spp.	<i>Eimeriidae</i>	Assay	Country/Reference
1-7 d.	41	7.3	NI		Merthiolate iodine formalin concentration	Germany/Wieler <i>et al.</i> , 2001
8-14 d.	102	39.2	NI			
15-21 d.	40	32.5	NI			
22-28 d.	18	11.1	NI			
29-35 d.	4	25	NI			
36-42 d.	82	1.2	NI			
10-20 d.	300 <sup>U</sup>	33	NI		Culture on 2.5% potassium dichromate, Telemann sedimentation method	Belgium/Leten <i>et al.</i> , 2002
Diarrheic	264 <sup>U</sup>	49.2	NI		Sedimentation-Flotation, McMaster	Germany/Niestrath <i>et al.</i> , 2002
Non-diarrheic	81 <sup>U</sup>	22	NI			
4-10 w.	265 <sup>NT</sup>			5.4	NaCl Flotation	China/Weng <i>et al.</i> , 2005
11-17 w.	259			18.9		
18-23 w.	231			20.3		
Sows	268			51.1		
Boars	85			61.1		
2-30 w.	133 <sup>N</sup>	76.7	NI		NaCl/sugar Flotation	Germany, Austria, Switzerland /Mundt <i>et al.</i> , 2005
1-7 d.	60 <sup>D</sup>			0	Flotation	Japan/Katsuda <i>et al.</i> , 2006
8-14 d.	44 <sup>D</sup>			36.4		
15-21 d.	46 <sup>D</sup>			26.1		
22-28 d.	40 <sup>D</sup>			10		
29-35 d.	46 <sup>D</sup>			26.1		
1-7 w.	514	10	7		Sodium acetate-acetic acid-formalin (SAF) concentration.	Germany/Damriyasa and Bauer, 2006
5-28 d.	571 <sup>L,U</sup>	31.7	1.4		NaCl/sugar Flotation, McMaster, culture on 2.5% potassium dichromate	Poland/Karamon <i>et al.</i> , 2007
5-28 d.	171 <sup>M,U</sup>	18.1	0.6			
5-28 d.	38 <sup>S,U</sup>	13.2	28.9			
Sows	159 <sup>L</sup>	8.2	4.4			
Sows	98 <sup>M</sup>	5.1	5.1			
Sows	10 <sup>S</sup>	0	40			
10 d. - 3 w.	123	6.4	NI		PCR- RFLP	Australia/Johnson <i>et al.</i> , 2008b
4 w. - 6 m.	156	16.3	NI			
Sows	10	0	NI			

**Table 10** (continuation). Worldwide reported fecal occurrence (%) of *Eimeriidae* in pigs during last decade.

Group & age	Tested	<i>I. suis</i>	<i>Eimeria</i> spp.	<i>Eimeriidae</i>	Assay	Country/Reference
7-15 d.	-	32.8	NI		Sheather's sugar Flotation, Giemsa and Ziehl-Nielsen staining	Serbia/Savic <i>et al.</i> , 2010
d. 13	-	44.3	NI			
Diarrheic	-	37.4	NI			
Non-diarrheic	-	15	NI			
10-18 d. <sup>U</sup>	580	37	NI		NaCl/glucose Flotation	Belgium/Laitat <i>et al.</i> , 2010
Suckers	118	65.6 <sup>D</sup>	NI		NaCl/glucose Flotation, McMaster	Greece/Skampardonis <i>et al.</i> , 2010
1-7 d.	30 <sup>D</sup>	0	NI		Flotation	Brazil/Cruz <i>et al.</i> , 2010
Suckers	709 <sup>U</sup>	26.4	NI		Cornell-Wisconsin flotation	Canada/ Aliaga-Leyton <i>et al.</i> , 2011

<sup>L</sup>, M and S, from large (>100 sows), medium (25-100 sows) or small (<25 sows) farms; <sup>U</sup>, litters; <sup>D</sup>, piglets were tested during all suckling period; <sup>N</sup>, never treated herds; <sup>D</sup>, all tested piglets had diarrhea; <sup>NT</sup>, no regular regime of anti-parasite treatment.  
Studies reported as *Eimeriidae* did not make the differential identification of *Eimeria* spp. and *I. suis* oocysts.

## **Aims of the study**

---



## Aims of the study

Porcine pre- and post-weaning diarrhea is a multi-factorial disease responsible for tremendous economic losses to the pig industry worldwide, and several infectious agents play a major role in the pathogenesis (Wieler *et al.*, 2001; Katsuda *et al.*, 2006).

In Cuba, swine veterinary practitioners have to face a high occurrence and a yearly increased incidence of porcine diarrhea, as well as limitations in the specific identification of its etiology, leading to uncontrolled administration of antibiotics, insufficient prevention programs, and scarce information on its epidemiology. In 2008, the Cuban Institute for Swine Research reported gastroenteric diseases (an unspecific terminology) to be the cause of 31% and 37% of the total piglet's mortality during the pre- and post-weaning periods, respectively (Cabrera and García, 2009).

Therefore, in order to contribute to the epidemiological characterization of enteropathogens of young pigs in Cuba, this PhD thesis focused on

- (Chapter 2) the identification of enteropathogens currently associated with pre- and post-weaning diarrhea,
- (Chapter 3) the antibiotic resistance and genetic relatedness of pathogenic *E. coli*,
- (Chapter 4) the seroprevalence of specific antibodies against *E. coli* fimbriae and
- (Chapter 5) the genetic susceptibility of the commercial breed currently raised in Cuban piggeries to enteric colonization by fimbriated *E. coli*.

With all these data and results we want to propose a strategy to remediate the problem of pre- and post-weaning diarrhea in Cuba.



## Chapter 2

---

*Association of toxigenic E. coli and several other pathogens with porcine pre- and post-weaning diarrhea in Villa Clara province, Cuba.*

Based on:

de la Fé Rodríguez PY, Maroto Martín LO, Cruz Muñoz E, Butaye P, Goddeeris BM, Cox E, 2011.

Several enteropathogens are associated with porcine pre-weaning and post-weaning diarrhea in Villa Clara province, Cuba.

Submitted to Tropical Animal Health and Production (manuscript number TROP-2977).







### 2.1 Abstract

Intestinal contents of suckling (n = 45) and recently weaned (n = 45) diarrheic pigs from Villa Clara province, Cuba, were tested for the presence of important viral, bacterial, and parasitic enteropathogens. At least one enteropathogen was detected in 64.4% and in 42.2% of suckling and weaned pigs, respectively. ETEC was significantly the most frequent pathogen in suckling piglets (28.9%;  $p < 0.05$ ) as well as in weaned pigs (22.2%), and most of isolates were either STa<sup>+</sup>/STb<sup>+</sup> or F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup>. The overall occurrence of the rest of pathogens was 10% for TGEV and *C. parvum*, 6.7% for rotavirus A and *I. suis*, 5.6% for  $\alpha$ -toxigenic *C. perfringens*, 3.3% for VTEC, and 2.2% for *Salmonella enterica* subspecies *enterica* serotype Newport. TGEV and  $\alpha$ -toxigenic *C. perfringens* were only identified in suckling piglets, while *S. Newport* and VTEC were only detected in weaned pigs. PEDV,  $\beta$ -toxigenic *C. perfringens*, *Eimeria* spp., and helminths were not identified during this study. Twelve out of 48 enteropathogen positive pigs (25%) carried mixed infections. ETEC was present in 10 out of 12 mixed infections, and TGEV infections were never combined. The results obtained herein demonstrate that several enteropathogens, either alone or as part of a combined infection, are associated with porcine pre- and post-weaning diarrhea in Villa Clara province, Cuba. It is necessary to implement actions aimed on the surveillance, prevention, and control of swine enteropathogens in Cuban piggeries.

## 2.2 Introduction

Many infectious agents, alone or in combination, can play an important role in the pathogenesis of porcine pre- or post-weaning diarrhea, which are multi-factorial diseases negatively affecting performance of young pigs and swine production efficiency (Katsuda *et al.*, 2006). Rotaviruses, TGEV, PEDV, ETEC, toxigenic *C. perfringens*, and *Coccidia* have been identified as the most associated pathogens with piglet's diarrhea worldwide (Chae *et al.*, 2000; Wieler *et al.*, 2001; Barrera *et al.*, 2005; Straw *et al.*, 2006; Katsuda *et al.*, 2006).

In 2008, the Cuban Institute for Swine Research reported that gastroenteric diseases caused 31% and 37% of the total piglet's mortality during the pre- and post-weaning periods, respectively (Cabrera and García, 2009). However, there is scarce information on the epidemiology of porcine enteropathogens because in most of the diarrhea outbreaks, diagnosis is restricted to clinical and macro-pathological examinations (Cabrera and García, 2009). Also, limited research with a focus on porcine enteropathogens has been performed during the last 20 years in Cuban piggeries (Fuentes *et al.*, 2001; Barrera *et al.*, 2005; Blanco *et al.*, 2006).

Therefore, in order to contribute to the epidemiological characterization of porcine infectious diarrhea in Cuba, the present study undertook the differential identification of enteropathogens that could be associated with porcine pre- or post-weaning diarrhea in the Villa Clara province.

## 2.3 Materials and methods

### *Survey design*

Along the period from May to June 2008, the department of Veterinary services of the Company of Pork Production (GRUPOR) of Villa Clara, which is the province that produces most pork in Cuba (González *et al.*, 2007), assigned 90 diarrheic piglets for being surveyed for enteropathogens. Piglets were distributed in 6 of the 8 largest indoor piggeries free of hog cholera outbreaks in the province, otherwise access and sampling is forbidden due to biosecurity rules. The herd of the 6 piggeries together comprised 5252 sows and 7455 suckling piglets; the number of newly weaned piglets was not controlled. All piggeries followed a similar continuous flow management (farrow-to-finish) without straw bedding, they had similar facilities, they used the same antibiotics, and they had shared breeding stock animals routinely. Pregnant sows were housed individually (Fig. 1), and at the end of gestation they were moved up to elevated farrowing crates with completely perforated plastic flooring (Fig. 2 and 3). After weaning, piglets were housed in groups of 10 to 20 in elevated pens with perforated floors (Fig. 4).

On the appointed day to sample in every piggery, the responsible veterinarian was asked to select during the early morning inspection 6-9 litters (2 to 25-day-old) and 6-9 newly weaned pens

(26 to 48-day-old) showing signs of watery diarrhea to complete 15 sampling places spread over the farrowing and weaning houses. Then, one piglet per litter or per pen was randomly caught, and selected and euthanatized if sign of diarrhea was present in the perineum. Before euthanasia, blood was collected and serum was obtained as described in Chapter 4. Piglets and sows were not under antibiotic therapy or vaccinated against enteropathogens.



**Figure 1.** Individual housing of pregnant sows in a Cuban piggyery.



**Figure 2.** Elevated farrowing crates with completely perforated plastic flooring currently used in Cuban piggyeries.



**Figure 3.** Inside and outside of a farrowing installation in a Cuban piggyery.



**Figure 4.** Weaned pigs raised in elevated pens with perforated floors in a Cuban piggyery.

Specimens (i.e. intestinal contents, serum, blood) were maintained in an insulated transport container filled with refrigerant gels until arrival to the laboratory, and Stuart Transport Medium modified by Ringertz was employed for keeping bacteria alive. For the identification of pathogens for which commercial diagnostic kits were planned to be used, intestinal contents and sera were

aliquoted and stored at -20°C until the kit application to all samples. Intestinal contents started to be processed quickly for bacterial culture and general parasitological methods.

### *Bacteriological identification*

In order to isolate *E. coli*, intestinal contents were inoculated on McConkey agar for 18 h at 37°C in aerobic atmosphere. Thereafter 3-5 lactose-fermenting colonies per sample were separately re-inoculated on McConkey agar. Then, for biochemical identification one colony per plate was inoculated on Luria agar for 18 h at 37°C, and was later identified as *E. coli* if it was able to ferment lactose and glucose with gas production (tested on Kligler Iron agar), to produce indole from tryptophan [visualized employing the Kovac's reagent: 50 g/l of 4-(dimethylamino)benzaldehyde + 75% of 11 M 1-butanol + 25% of concentrated 12 M HCl], and also if did not hydrolyze aesculin (tested on Bile Aesculin agar; Moyaert *et al.*, 2006; Bruggeman *et al.*, 2008) and if did not utilize citrate (tested on Simmons citrate agar). Intestinal contents were also plated onto 5% sheep blood agar and a maximum number of 3 hemolytic colonies were further processed as described above. Finally, one hemolytic *E. coli* isolate per pig was selected for being tested for fimbriae (F4, F5, F6, F18, and F41) and toxins (STa, STb, LT, and STx2e) encoding genes.

### *Identification of genes encoding virulence factors in E. coli by multiplex-PCR*

From an overnight culture on Brain Heart Infusion agar at 37°C, 3-4 colonies were suspended in 100 µl sterile distilled water, and heated for 15 min at 90°C in warm water bath. Next, the bacterial lysate was chilled on ice for 5 min and centrifuged at 14,000×g for 1 min. Aliquots of the supernatant were stored at -20°C and used as template DNA. The reaction mixture (20 µl) was composed by 2 µl of template DNA, 0.5 µM of each primer (Table 1; Bruggeman *et al.*, 2008), 0.2 mM of each dNTP (Pharmacia), 2.5 U of Taq polymerase (Ampli Taq Gold, Perkin Elmer), 1X of Ampli Taq Gold buffer (10 mM Tris pH 8.3, 5 mM KCl), and 5 mM of MgCl<sub>2</sub> (Perkin Elmer). The PCR was made on a thermocycler Gene Amp PCR System 9600 (Perkin Elmer) by pre-denaturation of the mixture at 90°C for 3 min, followed by 30 cycles of amplification including denaturation at 90°C for 1 min, annealing at 55°C for 1 min plus increment of 3 sec/cycle, and primer extension at 70°C for 2 min; the final extension was at 70°C for 10 min. The employed positive controls were *E. coli* B41 (F5<sup>+</sup>/F41<sup>+</sup>/STa<sup>+</sup>), E68 (F4<sup>+</sup>/LT<sup>+</sup>/STb<sup>+</sup>), and 107/86 (F18ab<sup>+</sup>/STx2e<sup>+</sup>). *E. coli* HB101 was the negative control. All PCR products were run at 120 V and 190 mA for 90 min in 3% Nusieve 3.1 agarose gel in 0.5X TBE buffer [10X TBE stock solution: 108 g Tris (Merck) + 55 g Boric Acid (Merck) + 9.3 g EDTA (Merck) + 1000 ml distilled water]. Staining was made with ethidium bromide (Sigma).

**Table 1.** Primer sequences used in the multiplex-PCR for detecting genes encoding virulence factors in *E. coli*.

Target gene	Virulence factor	Oligonucleotide sequence of forward and reverse primers (3'→5')	Amplicon size (bp)
<i>estII</i>	STb	TGCCTATGCATCTACACAAT CTCCAGCAGTACCATCTCTA	113
<i>estIa</i>	STa	CAACTGAATCACTTGACTCTT TTAATAACATCCAGCACAGG	158
<i>fanA</i>	F5	AATACTTGTTTCAGGGAGAAA AACTTTGTGGTTAACTTCCT	230
<i>eltB</i>	LT	GGCGTTACTATCCTCTCTAT TGGTCTCGGTCAGATATGT	272
<i>fedA</i>	F18	TGGTAACGTATCAGCAACTA ACTTACAGTGCTATTTCGACG	313
<i>fasA</i>	F6	GTAACCTCCACCGTTTGTATC AAGTTACTGCCAGTCTATGC	409
<i>faeG</i>	F4	GAATCTGTCCGAGAATATCA GTTGGTACAGGTCTTAATGG	499
<i>fimF41a</i>	F41	AGTATCTGGTTCAGTGATGG CCACTATAAGAGGTTGAAGC	612
<i>Stx2e</i>	STx2e	AATAGTATACGGACAGCGAT TCTGACATTCTGGTTGACTC	733

For identification of *Salmonella* spp., intestinal contents were inoculated on brilliant green agar (BGA) and in 5 ml of Rappaport-Vassiliadis broth for 18 h at 37°C in aerobic atmosphere, and then a re-inoculation on BGA was performed. Non-lactose-fermenting colonies from BGA were further biochemically identified and re-plated on brain heart infusion agar. Thereafter a slide agglutination was carried out using polyvalent (A+B+C1+C2+D+E1+E2+E4+F groups) and monovalent (A, B, C1, and C2 groups) *Salmonella* antisera (Company of biological products C.J. Finlay, Havana, Cuba). Positive isolates on slide agglutination were tested by the multiplex *Salmonella* DNA typing Premi®Test (Veterinary and Agrochemical Research Centre, Brussels, Belgium; Wattiau *et al.*, 2008).

The semi-quantitative detection of *C. perfringens* as well as the identification of the  $\alpha$ - and  $\beta$ -toxins was performed on intestinal contents by a DAS-ELISA kit (BIO K 095, Bio-X Diagnostics Inc., Jemelle, Belgium). A sample was considered positive if it was positive for the bacterium and for at least one toxin.

#### Virus identification

The specific detection of TGEV and PEDV in intestinal contents was carried out by a qualitative chromatographic immunoassay kit (Anigen Rapid TGE/PED Ag test kit, Animal Genetics Inc., Suwon, South Korea). Complementary, sera of suckling piglets were tested for antibodies against TGEV and PRCV by a differentiating blocking ELISA kit (SVANOVIR® TGEV/PRCV-Ab, Svanova Biotech AB, Uppsala, Sweden), which allows the detection of three serological statuses: seronegative piglets for both coronaviruses, seropositive piglets for PRCV and seronegative for TGEV, and seropositive

piglets for TGEV. Rotavirus A was tested in intestinal contents by a DAS-ELISA kit (1.1.RT.K2, Ingenasa, Madrid, Spain).

#### *Parasitological identification*

Intestinal contents were examined for *Eimeriidae* and helminths by light microscopy using the direct smear and Sheather's sugar flotation methods. As previously described by Karamon *et al.* (2007), oocysts sporulation was induced in *Eimeriidae* positive samples to allow the differential identification of *Eimeria* spp. and *I. suis*. *C. parvum* antigen detection was carried out using a DAS-ELISA kit (BIO K 070, Bio-X Diagnostics Inc., Jemelle, Belgium).

#### *Statistical analysis*

The proportions of piglets infected by every enteropathogen in every age group (suckling piglets, newly weaned piglets, 2 to 13, 14 to 25, 26 to 36, and 37 to 48-day-old piglets) as well as in the overall sample were compared by the chi-square test in StatGraphics Plus Version 5.0. Differences were considered significant at  $p < 0.05$ .

## **2.4 Results**

### *Identified enteropathogens*

At least one enteropathogen was identified in 29 out of 45 (64.4%) suckling piglets, and in 19 out of 45 (42.2%) weaned pigs (Table 2). PEDV,  $\beta$ -toxigenic *C. perfringens*, *Eimeria* spp., and helminths were not detected during this survey. The color of scours varied from milky to dark yellowish, and bloody diarrhea was not observed. Pathogenic *E. coli* (ETEC and VTEC) were the most frequent enteropathogens (Table 2), being isolated from intestinal contents of 28.9% of all pigs ( $n=90$ ). The combination of genes encoding STa and STb was commonly detected as most of the pathogenic *E. coli* were either STa<sup>+</sup>/STb<sup>+</sup> (31%) or F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup> (19%; Table 3). One STa<sup>+</sup>/STb<sup>+</sup> and all F18<sup>+</sup> or F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup> isolates were beta-hemolytic.

The only isolated *Salmonella* spp. was *Salmonella enterica* subspecies *enterica* serotype Newport (serogroup C2) in 2 (4.4%) weaned pigs (Table 2), and in one of both cases the infection was combined with ETEC and *C. parvum* (Table 4).

The bacterium *C. perfringens* was detected in 66.7% (30/45) of suckling piglets, but only 16.7% (5/30) of these *Clostridium*-positive piglets were positive for  $\alpha$ -toxin. It was noted that the  $\alpha$ -toxin-positive intestinal contents contained a higher bacterium concentration (average OD<sub>450nm</sub> = 1.679) than the toxin-negative ones (average OD<sub>450nm</sub> = 0.990). In weaned piglets, *C. perfringens* was identified in 26.7% of the pigs, whereas toxins were not detected in this group.



**Table 2.** Percentage of diarrheic piglets infected by enteropathogens and enteric mixed infections per age group between May and June 2008 in the Villa Clara province, Cuba.

Age groups	No. of pigs	Enteropathogens																Enteric mixed infections	Total <sup>a</sup>			
		Rotavirus A		TGEV		ETEC		VTEC		S. Newport		α-toxicogenic C. perfringens		I. suis		C. parvum						
		n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%					
Suckling	2-13-day-old	31	2	6.5	9	29*	7	22.6*	0	0	0	0	3	9.7	2	6.5	1	3.2	4	12.9	20	64.5
	14-25-day-old	14	2	14.3	0	0	6	42.9*	0	0	0	0	2	14.3	2	14.3	2	14.3	3	21.4	9	64.2
	Subtotal	45	4	8.9	9	20	13	28.9*	0	0	0	0	5	11.1	4	8.9	3	6.7	7	15.6	29	64.4
Weaned	26-36-day-old	24	1	4.2	0	0	5	20.8*	3	12.5	2	8.3	0	0	1	4.2	2	8.3	3	12.5	10	41.7
	37-48-day-old	21	1	4.8	0	0	5	23.8*	0	0	0	0	0	0	1	4.8	4	19	2	9.5	9	42.9
	Subtotal	45	2	4.4	0	0	10	22.2*	3	6.7	2	4.4	0	0	2	4.4	6	13.3	5	11.1	19	42.2
Total		90	6	6.7	9	10	23	25.6*	3	3.3	2	2.2	5	5.6	6	6.7	9	10	12	13.3	48	53.3

\* , Enteropathogens frequency which is significantly higher (p<0.05) than the frequency of other enteropathogens for pigs of the same age group.

<sup>a</sup> , Total of enteropathogen positive pigs per age group, pigs carrying mixed infections are counted as one.



**Table 3.** Association and diversity of fimbrial and toxin encoding genes among 26 pathogenic *E. coli* isolated from intestinal contents of suckling (s) and weaned (w) pigs with diarrhea in the Villa Clara province, Cuba.

Toxins Fimbriae	STa		STb		STa/STb		STb/LT		STx2e		None		Total (%)
	s	w	s	w	s	w	s	w	s	w	s	w	
<b>F4</b>	-	-	-	-	3	2	-	-	-	-	-	1	6 (23)
<b>F6</b>	-	1	-	-	-	-	-	-	-	-	-	-	1 (4)
<b>F18</b>	-	-	-	-	-	-	-	2	-	3	-	-	5 (19)
<b>F5/F41</b>	1	-	-	-	-	-	-	-	-	-	-	-	1 (4)
<b>None</b>	1	-	1	2	6	2	1	-	-	-	-	-	13 (50)
Total (%)	3 (12)		3 (12)		13 (50)		3 (12)		3 (12)		1 (4)		26 (100)

Rotavirus A was detected in every age group, with a slightly higher occurrence (14.3%) in 14- to 25-day-old piglets (Table 2). TGEV was identified in 29% of 2- to 13-day-old piglets only. Most of suckling piglets (77.7%) were seropositive for TGEV, which was excreted by 2 out of 5 (40%) TGEV seronegative piglets, as well as by 5 out of 35 (14%) TGEV seropositive ones. Only one piglet was seropositive for PRCV and seronegative for TGEV, and 3 were seronegative for both coronaviruses.

Comparing different age groups, it seemed as if the 14- to 25-day-old piglets were more susceptible to enteropathogens than the others; this age group showed the highest occurrence of rotavirus A, ETEC,  $\alpha$ -toxigenic *C. perfringens*, and *I. suis*, as well as the second highest of *C. parvum*. Also mixed infections were most frequent in this age group. However, it should be noted that the number of tested and therefore also the number of positive animals in this age group was low. Furthermore, the percentage of piglets showing enteric infections was as high in the 2-13-day-old age group.

**Table 4.** Mixed infections by enteropathogens identified in intestinal contents of suckling and weaned pigs suffering from diarrhea in Villa Clara province, Cuba.

Pigs	Age (days)	Mixed infections
Suckling	3	<i>C. parvum</i> + $\alpha$ -toxigenic <i>C. perfringens</i>
	5	rotavirus A + <i>E. coli</i> STb <sup>+</sup>
	11	<i>I. suis</i> + $\alpha$ -toxigenic <i>C. perfringens</i>
	13	rotavirus A + <i>E. coli</i> STa <sup>+</sup> /STb <sup>+</sup>
	14	<i>C. parvum</i> + <i>I. suis</i> + <i>E. coli</i> F4 <sup>+</sup> /STa <sup>+</sup> /STb <sup>+</sup>
	15	<i>I. suis</i> + rotavirus A + <i>E. coli</i> F4 <sup>+</sup> /STa <sup>+</sup> /STb <sup>+</sup>
	16	<i>C. parvum</i> + <i>E. coli</i> STa <sup>+</sup> /STb <sup>+</sup>
Weaned	34	rotavirus A + <i>E. coli</i> F4 <sup>+</sup> /STa <sup>+</sup> /STb <sup>+</sup>
	34	<i>I. suis</i> + <i>E. coli</i> F4 <sup>+</sup> /STa <sup>+</sup> /STb <sup>+</sup>
	36	<i>C. parvum</i> + <i>S. Newport</i> + <i>E. coli</i> STb <sup>+</sup>
	41	<i>I. suis</i> + <i>E. coli</i> STa <sup>+</sup> /STb <sup>+</sup>
	46	<i>C. parvum</i> + <i>E. coli</i> F4 <sup>+</sup>

Infections by multiple enteropathogens were detected in 25% of enteropathogen positive pigs. Mixed infections were slightly more common in suckling piglets (7/12) than in weaned pigs (5/12; Table 4). *I. suis* infections were almost always (5/6) combined with other pathogens; similar results were observed for rotavirus A (4/6) and *C. parvum* (5/9). Alpha-toxigenic *C. perfringens* infections were only mixed with *C. parvum* or *I. suis*. TGEV infections were not mixed, while ETEC was present in 10 out of 12 combined infections.

## 2.5 Discussion

In Cuba, diarrhea markedly influences piglet's mortality (Cabrera and García, 2009). However, knowledge of which enteropathogens could be associated with this syndrome has been limited during the last 20 years. Therefore, the study reported herein updates on the frequency of enteropathogens associated with porcine pre- and post-weaning diarrhea in Cuban piggeries.

Several enteropathogens, either alone or as part of a mixed infection, were identified in 64.4% of suckling piglets and in 42.2% of weaned pigs with diarrhea. ETEC were more frequent ( $p < 0.05$ ) than other enteropathogens in all age groups, except in the 2- to 13-day-old piglets, in which they were as frequent as TGEV. Surveys aimed on the differential identification of enteropathogens in diarrheic piglets reported *I. suis* in Germany (Wieler *et al.*, 2001) and rotavirus in Japan (Katsuda *et al.*, 2006) as the most prevalent agents, the authors attributed a low occurrence of ETEC to vaccination programs and to the use of effective antibiotics. In Cuban piggeries, housing and management conditions are different, vaccination against ETEC has not been applied during the last years (Wong *et al.*, 1995), and there is a high resistance rate of porcine pathogenic *E. coli* to the routinely administered antibiotics (de la Fé Rodríguez *et al.*, 2012).

F4, F5, and F41 encoding genes have not been previously identified in *E. coli* isolates from diarrheic piglets in Cuba (Blanco *et al.*, 2006). Furthermore, our study is the first to demonstrate the enterotoxin encoding genes (STa or STb) in F4<sup>+</sup> and F41<sup>+</sup> ETEC isolates from Cuba as their identification was not performed by Fuentes *et al.* (2001) and Lazo *et al.* (2005). LT was only identified in 3 isolates in the present survey; nevertheless, most of ETEC carried genes encoding ST enterotoxins, which are sufficient for causing diarrhea in neonatal (STa) and newly weaned (STb) pigs (Berberov *et al.*, 2004).

F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup> ETEC as identified in our study, were also isolated from diarrheic piglets in South Korea (Chae *et al.*, 1998; Kwon *et al.*, 2002); in contrast, a strong association between F4 and LT encoding genes has been shown in ETEC associated with swine diarrhea worldwide (Frydendahl, 2002; Zhang *et al.*, 2007). The two F18<sup>+</sup>/LT<sup>+</sup>/STb<sup>+</sup> ETEC isolated from weaned pigs constitute the first report of F18<sup>+</sup> ETEC carrying LT encoding genes in Cuba, corresponding with previous findings in USA (Zhang *et al.*, 2007), and differ from the F18<sup>+</sup>/STx2e<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup> strains previously isolated in Villa

Clara by Blanco *et al.* (2006). Recently, a high seroprevalence of specific antibodies against F4 and F18 fimbriae was demonstrated in Cuban swine (de la Fé Rodríguez *et al.*, 2011), supporting the importance of F4<sup>+</sup> or F18<sup>+</sup> *E coli* in the pathogenesis of swine diarrhea or edema disease over the whole country. ETEC harboring the F6 fimbrial gene are often associated with pre-weaning diarrhea (Blanco *et al.*, 2006), but in the present survey F6 was only linked with post-weaning diarrhea in a low rate like reported by Frydendahl (2002).

*Salmonella* spp. were not as frequently isolated as ETEC; only 2 weaned pigs were infected by *S. Newport*. There is no previous information on the presence of this pathogen in Cuban piggeries. Fever, diarrhea, dehydration, and enteritis have been observed in calves, cows, and horses from which *S. Newport* was isolated (Poppe *et al.*, 2006), but more research on its role in the pathogenesis of piglet's diarrhea is needed (Zhao *et al.*, 2003). Katsuda *et al.* (2006), and other authors (see Tables 3 and 4, Chapter 1), could not isolate *Salmonella* in suckling and newly weaned pigs with diarrhea. Nevertheless, the *Enterobacteriaceae Salmonella* cause diarrhea and enterocolitis in pigs mostly from post-weaning to around 4 months of age, and *Salmonella enterica* subspecies *enterica* serotype Cholerasuis and serotype Typhimurium are the most commonly associated (Griffith *et al.*, 2006). Recently Hur *et al.* (2011) reported that 42 *S. Typhimurium* isolated from diarrheic pigs aged from 2 to 72 days in South Korea carried diverse genes encoding antibiotic resistance and virulence factors. Therefore, it has been recommended that swine production systems have to be routinely surveilled for *Salmonella*, a zoonotic foodborne pathogen (Chiu *et al.*, 2004).

In Cuba, the identification of *C. perfringens* toxins is not performed during the routine analysis of swine diarrhea. The higher concentration of this bacterium detected in  $\alpha$ -toxin-positive intestinal contents when comparing with the  $\alpha$ -toxin-negative ones, suggests the presence of active infectious processes due to  $\alpha$ -toxigenic *C. perfringens* in these suckling piglets. Das *et al.* (2009) found a significant association of *C. perfringens* A with swine diarrhea in India. Here  $\alpha$ -toxigenic *C. perfringens* was the single enteropathogen in 3 out of 5 positive piglets, whereas coinfection with *C. parvum* or *I. suis* was seen in both other piglets; Songer and Uzal (2005) stated that preceding lesions provoked by other enteropathogens can enhance enteric colonization by *Clostridium*. Beta-toxigenic *C. perfringens* was not identified, supporting the absence of reports on necrotic enteritis in the sampled piggeries which might be due to a  $\beta$ -toxin effect.

TGEV always occurred as a single infection of suckling piglets, and the associated diarrhea was not profuse and epidemic like in the classical TGE acute form. A milder form of TGE can be commonly observed (Morin *et al.*, 1983), and its clinical differentiation from coccidial, rotaviral, and ETEC infections is difficult, as clinical signs are less severe than those observed in the classic form. From 2003 on, epidemic TGE outbreaks have been reported in Cuba (Barrera *et al.*, 2005), but the milder behavior found in the present study, and the high amount of seropositive suckling piglets, indicates a

possible change to the endemic form and a high prevalence at herd level as the serological status of newborn piglets is influenced by maternal antibodies. Additionally, antibodies against PRCV were for the first time reported for Cuba during the present survey. So presence of these antibodies could also explain milder TGEV infections since it is well-known that an infection with PRCV induces antibodies able to neutralize and decrease the excretion of TGEV, inhibiting the presentation of epidemic TGE in PRCV positive herds (Cox *et al.*, 1993).

Four out of 6 rotavirus A infections were mixed with ETEC; interestingly, three of them occurred in suckling piglets. Rotaviruses are of importance in the pathogenesis of piglet's diarrhea worldwide (Katsuda *et al.*, 2006; Halaihel *et al.*, 2010), and are often seen in association with other enteropathogens (e.g. ETEC; Melin *et al.*, 2004; Kim *et al.*, 2010a), suggesting that they provoke damage on the epithelium that can alter the binding sites on enterocytes, favoring gut colonization by other enteropathogens (Melin *et al.*, 2004).

In suckling piglets *I. suis* occurred with much lower frequency (8.9%) than previously reported (44.7%) in the Havana province by Koudela *et al.* (1989); the housing improvement carried out in Cuban piggeries during last years could be a factor responsible for this difference. Like in the present survey, in Australia the highest frequency of this parasite was found in suckling piglets (Johnson *et al.*, 2008). Four out of 6 (66.6%) *I. suis* positive pigs were also infected with ETEC, similar results (45.7%) were shown by Chae *et al.* (1998) in South Korea. Choi *et al.* (2003) stated that isosporosis could promote intestinal colonization by ETEC due to an increase of glycoconjugates on the jejunal enterocytes.

The 10% occurrence of *C. parvum* found here is higher than the 2.1% reported in Havana province, Cuba, by Cabrera and García (1985). However, our higher prevalence could be due to the high sensitivity of the identification procedure used here (DAS-ELISA) compared with flotation and staining methods. The prevalence of *C. parvum* in young pigs has ranged from 1.4% to 71% worldwide (Izumiyama *et al.*, 2001; Wieler *et al.*, 2001; Yu and Seo, 2004; Maddox-Hyttel *et al.*, 2006). Experimental infections have confirmed the pathogenicity of *C. parvum* in pigs (Enemark *et al.*, 2003ab), and field studies have displayed a positive correlation of *Cryptosporidium* with swine diarrhea (Mišić *et al.*, 2003; Hamnes *et al.*, 2007). Probably, *C. parvum* acts often in concert with other enteropathogens to induce or exacerbate diarrhea; in that context it is interesting to remark that a mixed infection of rotavirus and *Cryptosporidium* aggravates diarrhea (Enemark *et al.*, 2003a), this mixed condition was not identified in the present study, but 55.5% of *C. parvum* infections were found to be combined with other enteropathogens.

That 25% of enteropathogen positive pigs had mixed infections confirms that in order to control diarrhea on a piggery, a thorough differential diagnosis has to be performed as recommended by Katsuda *et al.* (2006) and Straw *et al.* (2006). However, worldwide many studies assessing the

infectious etiology of swine diarrhea have only analyzed single etiologies (Chae *et al.*, 2000; Hong *et al.*, 2006). The differential identification of swine enteropathogens should be performed during *in vivo* experiments on swine diarrhea in order to accurately know the influence, or not, of certain pathogens on results (Baba and Gaafar, 1985; Choi *et al.*, 2003; Jensen *et al.*, 2006; Jung *et al.*, 2008). For instance, Lecce *et al.* (1982) reported that diarrhea caused by a rotavirus-ETEC mixed infection is more severe than the one caused by rotavirus or ETEC infection alone.

Enteropathogens were not detected in 46.7% of pigs during this study. This could be due to 1/tests sensitivity; 2/non-infectious factors (Straw *et al.*, 2006); 3/enteropathogens not-tested in the present study like other groups of rotaviruses (Kim *et al.*, 2010a), *Clostridium difficile* or PRRSV (Yaeger *et al.*, 2002); 4/not testing of some virulence factors (i.e. EAST-I, AIDA-I, PAA; Lee *et al.*, 2008); 5/insufficient recovery of the intestinal mucosa, such as the villous structure, after pathogen elimination; 6/ testing during the prepatent period of parasites where parasites are not yet detectable by general parasitological methods (Karamon *et al.*, 2007; Johnson *et al.*, 2008).

In conclusion, the study reported herein demonstrates that several enteropathogens are associated with porcine pre- and post-weaning diarrhea in the Villa Clara province, Cuba, of which ETEC is the most frequent. The improvement of swine enteropathogens identification, as well as the implementation of prevention and control programs for piglet's diarrhea which consider all potentially associated enteropathogens, particularly ETEC, might have a positive impact on the performance of young pigs in Cuba.

## **2.6 Acknowledgments**

This study was funded by an Interuniversity Cooperation Program among Vlaams Interuniversitaire Raad (Belgium) and Universidad Central "Marta Abreu" de Las Villas (Cuba), specifically by the sub-project 5: *Improving the quality of graduate and postgraduate education and research programs in plant and animal sciences*. The Cuban Company for Pork Production and the Cuban Institute of Veterinary Medicine are thanked for supporting this survey. Are specially thanked the veterinarians from the sampled piggeries and colleagues Dr. Ania Gonzalez, DVM Einar Artiles, DVM Alberto Rodríguez, and DVM Lester Rodríguez for their help during sampling.

## Chapter 3

---

*Antibiotic resistance and genetic relatedness among pathogenic E. coli isolated from intestinal contents of diarrheic piglets in Villa Clara province, Cuba.*

Based on:

de la Fé Rodríguez PY, Ndemi Kiiru J, Maroto Martin LO, Cruz Muñoz E, Butaye P, Cox E, Goddeeris BM, 2012. Characterization and clonal grouping of pathogenic *Escherichia coli* isolated from intestinal contents of diarrheic piglets in Villa Clara province, Cuba, according to their antibiotic resistance and ERIC-PCR profiles. *Veterinary Microbiology*, 154:425-8.



### 3.1 Abstract

This survey was undertaken to determine the insufficiently known antibiotic resistance and genetic relatedness among pathogenic *E. coli* associated with piglet's diarrhea in Villa Clara province, Cuba. Twenty six *E. coli* belonging to 10 virotypes, and isolated from intestinal contents of 90 piglets suffering from pre- or post-weaning diarrhea in 6 large indoor piggeries, were tested by the Kirby-Bauer disk diffusion method for antibiotic resistance, and by ERIC-PCR (employing primers ERIC1R: 5'-ATG-TAA-GCT-CCT-GGG-GAT-TCA-C-3' and ERIC2F: 5'-AAG-TAA-GTG-ACT-GGG-GTG-AGC-G-3') for genetic relatedness. The higher resistance rates were displayed to antibiotics of old generations which have been traditionally administered under uncontrolled prescriptions in Cuban piggeries. A significantly highest ( $p < 0.05$ ) resistance rate was found to tetracycline (69%); also, the resistance to ampicillin (54%), sulphonamide compounds (50%) and kanamycin (50%) were high, and 65% of isolates were multi-drug resistant. The ERIC-PCR revealed a high degree of polymorphism in the *E. coli* DNA sequences and relatedness among F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup> or F18<sup>+</sup>/LT<sup>+</sup>/STb<sup>+</sup> virotypes isolated from different piggeries, and among STb<sup>+</sup>, STa<sup>+</sup>/STb<sup>+</sup>, F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup> or F18<sup>+</sup>/STx2e<sup>+</sup> isolates from the same piggery. The antibiotic resistance and ERIC-PCR profiling were of help when assessing similarity of isolates: 17 clonal groups were identified. The results of this survey confirm that genetically diverse as well as genetically related pathogenic *E. coli* highly susceptible to many antibiotics (i.e. nalidixic acid, ciprofloxacin, gentamicin, amikacin, chloramphenicol, cephalosporins, amoxicilline-clavulanic acid, and trimethoprim), are associated with piglet's diarrhea in Villa Clara province, Cuba. These findings might be implemented during surveillance and control programs of swine colibacillosis in Cuba.



### 3.2 Introduction

Enteric infections with ETEC or VTEC are important causes of diarrhea or edema disease in pigs, which cause huge economic losses to the swine industry due to mortality, costs of medication, and growth retardation (Francis, 1999; Lazo *et al.*, 2005). Studies aimed on the differential identification of enteropathogens have reported ETEC, alone or as part of mixed infections, as a commonly associated pathogen with piglet's diarrhea worldwide (Wieler *et al.*, 2001; Nagy and Bilkei, 2003; Katsuda *et al.*, 2006; Hong *et al.*, 2006).

The antibiotic resistance profiling and genetic relatedness study among enteropathogenic *E. coli* are important contributions to the epidemiological characterization and control of swine diarrhea (Osek, 2000; Maynard *et al.*, 2003; Hariharan *et al.*, 2004; Lee *et al.*, 2009). However, in Cuba, where diarrheic diseases are responsible for 31% and 37% of the total piglet's mortality during the pre- and post-weaning periods, respectively (Cabrera and García, 2009), the identification of enteropathogens is limited, and the antibiotic resistance of porcine enterobacteria has not been surveyed (Barrera *et al.*, 2005; Lazo *et al.*, 2005; Blanco *et al.*, 2006).

Many DNA-based tests have been applied to reveal genetic relatedness among *Enterobacteriaceae* (e.g. PFGE, REP-PCR, BOX-PCR, and ERIC-PCR; Versalovic *et al.*, 1991; Cesaris *et al.*, 2007; Duan *et al.*, 2009). Enterobacterial repetitive intergenic consensus (ERIC) sequences are highly conserved at the nucleotide sequence level, and they constitute an effective marker when identifying clonal variability of isolates because their chromosomal locations and numbers differ among *Enterobacteriaceae* clones (Versalovic *et al.*, 1991). The ERIC-PCR fingerprinting analysis has been proved to be discriminatory during molecular differentiation of *E. coli* of porcine origin, and provides a better knowledge of their genetic relatedness and clonal origin (Osek, 2000; Namvar and Warriner, 2006; Yuan *et al.*, 2010). Additionally, Dias *et al.* (2009) observed that the circulating clonal composition of pathogenic *E. coli* tested by ERIC-PCR correlates with drug resistance prevalence.

Then, the aim of this study was to determine the antibiotic resistance profile and genetic relatedness among pathogenic *E. coli* isolated from piglets suffering from pre- or post-weaning diarrhea in Villa Clara province, Cuba, as an important contribution to the surveillance and control programs of swine colibacillosis.

### 3.3 Materials and methods

#### *Animals and E. coli isolates*

All pathogenic *E. coli* surveyed herein were isolated in the study discussed in Chapter 2, which was aimed at identifying the frequency of enteropathogens in intestinal contents of pigs suffering

from pre-weaning (n = 45) or post-weaning diarrhea (n = 45) in six large indoor piggeries (>500 sows per piggery) located in Villa Clara province, Cuba.

#### *Antibiotic susceptibility test*

The *E. coli* isolates were tested by the Kirby-Bauer disk diffusion method on Mueller-Hinton agar according to the recommendations of the National Committee for Clinical Laboratory Standards (NCCLS; M2-A7, M100-S10). For this test were selected disks containing antibiotics traditionally administered in Cuban piggeries (i.e. tetracycline-30 µg, sulphonamide compounds-300 µg, ampicillin-10 µg, gentamicin-10 µg, and kanamycin-30 µg; Oxoid), as well as disks containing antibiotics less or practically never administered (i.e. trimethoprim-5 µg, trimethoprim:sulphamethoxazole/1:19-25µg, chloramphenicol-30 µg, amoxicilline-clavulanic acid-30 µg, ciprofloxacin-5 µg, nalidixic acid-30 µg, cefotaxime-30 µg, cefazolin-30 µg, and amikacin-30 µg). The isolates were classified in susceptible, intermediate resistant or resistant according to the inhibition zone diameter showed to each antibiotic and the breakpoints referred by the NCCLS. *E. coli* ATCC 25922 was used as susceptible control, and *Staphylococcus aureus* ATCC 3359 was the resistant control for most of antibiotics. Resistance rates were calculated by dividing the number of resistant and intermediate-resistant isolates by the total number of isolates. Comparison of resistance rates among antibiotics was performed by the chi-square test ( $\chi^2$ ) in StatGraphics Plus Version 5.0 ( $p < 0.05$ ).

#### *Genetic relatedness test*

Genetic relatedness was determined by the ERIC-PCR fingerprinting assay (Versalovic *et al.*, 1991). The ERIC-PCR reaction (25 µl) contained 3 µl of template DNA previously obtained (Chapter 2), 25 pmol of each primer (ERIC1R: 5'-ATG-TAA-GCT-CCT-GGG-GAT-TCA-C-3' and ERIC2F: 5'-AAG-TAA-GTG-ACT-GGG-GTG-AGC-G-3'; Versalovic *et al.*, 1991), 600 µM of dNTPs (Roche Diagnostics GmbH), 1 U of SuperTaq (HT Biotechnology Ltd.), 2.5 µl of 10x SuperTaq buffer, and 2 mM of MgCl<sub>2</sub> (Promega). The PCR amplifications were performed in a thermal cycler PTC-100™ (MJ Research Inc.) by pre-denaturation of the mixture at 95°C for 7 min, followed by 30 cycles of denaturation at 94°C for 1 min, annealing at 52°C for 1 min, and a long extension at 68°C for 8 min, the final extension was at 65°C for 13 min. Eight µl of the PCR products were electrophoretically separated in a 1.5% agarose gel in 1xTAE for 40 min at 80 V plus 1.5 h at 90 V. The DNA bands were stained with ethidium bromide for 30 min and visualized under UV light; the gel image capture was helped by the software Quantity-One (BIO-RAD Laboratories Inc.). The *E. coli* isolates were considered to belong to a single clonal group if they showed a highly similar electrophoretic banding pattern for the ERIC-PCR

products as previously described (Osek, 2000; Dias *et al.*, 2009; Bert *et al.*, 2010), and if they had a similar antibiotic resistance profile.

### 3.4 Results

#### *Antibiotic susceptibility*

Only 4 out of 26 *E. coli* were susceptible to all tested antibiotics. A significantly highest ( $p < 0.05$ ) resistance rate (69%) was detected to tetracycline. The resistance rates to ampicillin (54%), sulphonamide compounds (50%), and kanamycin (50%) were also high. Significantly lower ( $p < 0.05$ ) resistance rates were found for chloramphenicol (12%), nalidixic acid (12%), and gentamicin (4%). All isolates were susceptible to the rest of tested antibiotics (i.e. trimethoprim, trimethoprim:sulphamethoxazole, amoxicilline-clavulanic acid, ciprofloxacin, cefotaxime, cefazolin, and amikacin). Multi-drug resistance was present in 17 out of 26 (65%) isolates: 4 (15%) were resistant to two antibiotics, 3 (12%) to three, 8 (31%) to four, and 1 (4%) to five and six.

F18<sup>+</sup>/LT<sup>+</sup>/STb<sup>+</sup> and F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup> isolates were the most multi-resistant among fimbriated *E. coli*. Chloramphenicol resistance was mostly found in all F18<sup>+</sup>/LT<sup>+</sup>/STb<sup>+</sup> *E. coli*, and VTEC (F18<sup>+</sup>/STx2e<sup>+</sup>) were susceptible to all antibiotics, except to sulphonamide compounds (Fig. 1).

































#### *Genetic relatedness and clonal groups*

The ERIC-PCR revealed that pathogenic *E. coli* associated with porcine pre- and post-weaning diarrhea in Villa Clara province, Cuba, have a high degree of polymorphism in their DNA sequences (Fig. 1), suggesting that most of them are from different clones.

Seventeen clonal groups were distinguished among 26 *E. coli* tested herein. Overall, the genetic profiling corresponded with the antibiotic resistance profile of the isolates and with the virotype where to they belonged (Fig. 1).

The 3 F18<sup>+</sup> VTEC isolated from piggery D belonged to a homogeneous group (XII) as they generated almost an identical ERIC fingerprinting profile and had a similar antibiotic resistance profile. The 2 F18<sup>+</sup>/LT<sup>+</sup>/STb<sup>+</sup> *E. coli*, which were isolated from different piggeries (E and F), had also similar ERIC-PCR and antibiotic resistance, they belonged to group XI.

Four F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup> *E. coli* isolated from suckling and weaned pigs in piggeries E and C appeared to belong to a single clonal group (XV). Although having a similar antibiotic resistance, the banding pattern of the F4<sup>+</sup>/- *E. coli* isolated in piggery C was slightly different from the ones of F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup> *E. coli*. There appeared a genetic diversity among the 8 STa<sup>+</sup>/STb<sup>+</sup> *E. coli*, which were categorized in 6 clonal groups. The 2 STb<sup>+</sup> *E. coli* isolates from piggery A, which did not have antibiotic resistance, had identical genetic profiles.

Virotype	Piggery/pig	ERIC-PCR profile	Antibiotic resistance profile	Group
STa <sup>+</sup>	D/s		Amp-Sul	I
STb <sup>+</sup>	F/w		Tet-Amp-Sul-Kan-Chl-Gen	II
	A/w		None	III
	A/s		None	III
STa <sup>+</sup> /STb <sup>+</sup>	F/s		Tet-Amp-Kan	IV
	D/w		Tet-Kan-Nal	V
	D/w		Tet-Amp-Kan-Nal	V
	D/s		Tet-Kan	VI
	D/s		Kan	VII
				
	B/s		Tet	VIII
	B/s		Tet	VIII
	B/s		Tet-Amp	IX
LT <sup>+</sup> /STb <sup>+</sup>	C/s		Tet-Amp-Sul-Kan	X
F18 <sup>+</sup> /LT <sup>+</sup> /STb <sup>+</sup>	E/w		Tet-Amp-Sul-Chl	XI
				
	F/w		Tet-Sul-Kan-Chl-Nal	XI
F18 <sup>+</sup> /STx2e <sup>+</sup>	D/w		Sul	XII
	D/w		None	XII
	D/w		Sul	XII
F6 <sup>+</sup> /STa <sup>+</sup>	D/w		None	XIII
				
F4 <sup>+</sup>	C/w		Tet-Amp-Sul-Kan	XIV
F4 <sup>+</sup> /STa <sup>+</sup> /STb <sup>+</sup>	E/s		Tet-Amp-Sul-Kan	XV
	E/s		Tet-Amp-Sul	XV
	C/s		Tet-Amp-Sul-Kan	XV
	C/w		Tet-Amp-Sul-Kan	XV
	C/w		Tet-Amp-Sul-Kan	XVI
F5 <sup>+</sup> /F41 <sup>+</sup> /STa <sup>+</sup>	E/s		Tet-Amp	XVII
S. Newport	F/w		Amp-Sul	S-I
	A/w		Sul	S-I
				

**Figure 1.** Clonal groups of pathogenic *E. coli* associated with piglet's diarrhea in Villa Clara province, Cuba, according to their virotype, ERIC-PCR and antibiotic resistance profiles.

S, suckling piglet; W, weaned pig; M, 100bp marker run in the same gel of ERIC-PCR products visualized on its top; Amp, ampicillin; Sul, sulphonamide compounds; Tet, tetracycline; Kan, kanamycin; Chl, chloramphenicol; Gen, gentamicin; Nal, nalidixic acid; S. Newport, *Salmonella enterica* subspecies *enterica* serotype Newport isolated from intestinal contents of weaned pigs and analyzed by the ERIC-PCR and antibiotic susceptibility tests employed for *E. coli*.

### 3.5 Discussion

For the last 20 years, specific reports or epidemiological studies regarding porcine enteropathogens in Cuba have been scarce (Fuentes *et al.*, 2001; Barrera *et al.*, 2005; Lazo *et al.*, 2005; Blanco *et al.*, 2006). A recent survey by de la Fé Rodríguez *et al.* (2011) reported a high serological prevalence of F4<sup>+</sup> and F18<sup>+</sup> *E. coli* in the Cuban swine herd, indicating that these pathogenic bacteria are widespread as potential etiologic agents of colibacillosis; also, ETEC was found to be the most associated enteropathogen with piglet's diarrhea in Villa Clara province, Cuba (25.6% frequency; Chapter 2). Additionally, the present survey determined the antibiotic resistance profile and genetic relatedness among pathogenic *E. coli* associated with porcine pre- and post-weaning diarrhea in Villa Clara province.

A high resistance rate to the traditionally administered antibiotics and a high genetic diversity among pathogenic *E. coli* were observed. The genetic profiling by ERIC-PCR enabled to categorize 26 pathogenic *E. coli* into 17 clusters with a more or less close relationship with the different antibiotic resistance profiles, taking into consideration the virotype of the isolates. Such integral studies have been rarely reported (Lee *et al.*, 2009; Thorsteinsdottir *et al.*, 2010).

Different from reports showing high resistance of *Enterobacteriaceae* to a wide range of antibiotics worldwide (Hendriksen *et al.*, 2008; Knezevic and Petrovic, 2008; Smet *et al.*, 2009; Tian *et al.*, 2009), herein were exclusively displayed high resistance rates to antibiotics routinely used for treating diarrhea and other diseases in Cuban piggeries during last years (i.e. tetracycline, ampicillin, sulphonamides, and kanamycin). Vieira *et al.* (2009) stated that there is an association between tetracycline treatment incidence rate in a herd and the probability of isolating a tetracycline resistant *E. coli* from the intestinal content of pigs. The resistance rate to gentamicin was low (4%) despite being frequently administered in Cuba. Except for chloramphenicol and nalidixic acid, for which 12% resistance rates were displayed, all isolates were susceptible to antibiotics less or practically never administered in Cuban piggeries (i.e. trimethoprim, trimethoprim:sulphamethoxazole, amoxicilline-clavulanic acid, ciprofloxacin, cefotaxime, cefazolin, and amikacin). Maynard *et al.* (2003) concluded that the genes behind phenotypic resistance are not static but are rather in a state of flux driven by various selection forces such as the use of specific antimicrobials, statement that may also justify the high susceptibility of *E. coli* to antibiotics less or practically never administered in Cuban piggeries found here.

In the near future, as recommended by Hendriksen *et al.* (2008) and Harada *et al.* (2008), data from antibiotic resistance surveillance in every geographic area or piggery have to be taken into account by the veterinary authorities when choosing antimicrobials for controlling swine colibacillosis in Cuba; but first, the specific identification and characterization of enteropathogens

have to be improved in the regional Veterinary Diagnostic Laboratories in order to avoid the uncontrolled antibiotic indication for treating diarrhea, which is mostly diagnosed by only clinical or macropathological examinations.

It is interesting that similar to this report, Maynard *et al.* (2003), Hariharan *et al.* (2004), Varga *et al.* (2008), and Akwar *et al.* (2008) found high resistance rates to tetracycline, ampicillin, and sulphonamide compounds, as well as high susceptibility to gentamicin and cephalosporins in porcine enteric *E. coli* from Canada. In Cuba, thousands of breeding stock pigs have been imported from Canada during last decade, and they could be the source of pathogens of non-obligatory declaration like *Enterobacteriaceae* carrying genes encoding drug resistance, but other epidemiological studies are needed to prove it. Previous information about antibiotic resistance of porcine *E. coli* from Cuba is not available (Blanco *et al.*, 2006).

The ERIC-PCR was applied for the first time to Cuban isolates of porcine pathogenic *E. coli* and was useful for their clonal and epidemiological discrimination. Among herds, F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup> isolates from piggeries E and C, as well as F18<sup>+</sup>/LT<sup>+</sup>/STb<sup>+</sup> isolates from piggeries E and F, were found to be clonally related. The frequent trading of breeding stock pigs among piggeries and other ways affecting piggery's biosafety could lead the spreading of pathogenic *E. coli* in Villa Clara province. Clonal relations were also found inside herds for STb<sup>+</sup> isolates in piggery A, for STa<sup>+</sup>/STb<sup>+</sup> isolates in piggeries D and B, for F18<sup>+</sup>/STx2e<sup>+</sup> isolates in piggery D, and for F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup> isolates in piggeries E and C. ERIC-PCR can be applied for identifying bacterial species, types, or isolates (Yuan *et al.*, 2010), and is an accurate and reliable technique for studying the molecular epidemiology of pathogenic *Enterobacteriaceae* (Dalla-Costa *et al.*, 1998; Manges *et al.*, 2004; Ramchandani *et al.*, 2005; Ibenyassine *et al.*, 2006; Jurkovič *et al.*, 2007) and other bacteria like *Pasteurella* (Saxena *et al.*, 2005). ERIC-PCR profiling of *E. coli* enabled also to identify contamination sites in a high capacity pork slaughter line (Namvar and Warriner, 2006).

In conclusion, the antibiotic therapy routinely employed in Cuba for controlling swine colibacillosis should be changed as most of *E. coli* tested herein were resistant to all commonly administered antibiotics, except to gentamicin. The introduction of other drugs for which ETEC showed high susceptibility (e.g. nalidixic acid, ciprofloxacin, gentamicin, or trimethoprim) appears advantageous; but first, it is necessary that the provincial Veterinary Diagnostic Laboratories improve the identification of pathogenic *E. coli* as well as the antibiotic resistance surveillance. The analysis on genetic diversity by ERIC-PCR demonstrated a clonal relationship among pathogenic *E. coli* carrying the same virulence factors and similar antibiotic resistance. The high genetic diversity of porcine enteropathogenic *E. coli* found in this survey was previously demonstrated in Cuba, when Blanco *et al.* (2006) described 21 distinct restriction patterns in 24 pathogenic *E. coli* by PFGE. As

both studies have been only performed in Central Cuba, the application of these epidemiological researches throughout the whole country might identify the *E. coli* clones which are most associated with clinical disease, and will contribute to the surveillance, prevention, and control of swine colibacillosis.

### **3.6 Acknowledgments**

This study was funded by an IUC program between VLIR-UOS (Belgium) and Universidad Central “Marta Abreu” de Las Villas (Cuba). DVM Miguel Angel Arce, DVM Einar Artiles, and Ir. René Cupull are specially thanked for their assistance during antibiotic resistance testing.



## Chapter 4

---

### *High prevalence of F4<sup>+</sup> and F18<sup>+</sup> E. coli in Cuban piggeries as determined by serological survey*

Based on:

de la Fé Rodríguez PY, Coddens A, Del Fava E, Cortiñas Abrahantes J, Shkedy Z, Maroto Martin LO, Cruz Muñoz E, Duchateau L, Cox E, Goddeeris BM, 2011. High prevalence of F4<sup>+</sup> and F18<sup>+</sup> *Escherichia coli* in Cuban piggeries as determined by serological survey. Tropical Animal Health and Production, 43:937-46.





#### 4.1 Abstract

Little information is available on the prevalence of swine enteropathogens in Cuba where diarrheic diseases are responsible for 31% and 37% of the total mortality during the neonatal and post-weaning periods. F4<sup>+</sup> and F18<sup>+</sup> ETEC and F18<sup>+</sup> VTEC induce diarrhea and edema disease in pigs, but their distribution has never been thoroughly studied in the Cuban swine population. Therefore, the present study estimated the prevalence of F4- and F18-specific antibodies in sera of 1044 6-months old gilts distributed in 34 piggeries spread over the Cuban territory. For the data analysis, which included the optical density of individual samples tested by ELISA, random-effects models and a mixture model in R (package “mixAK”; Komárek, 2009) were fitted. Low, moderate, and high levels of F4-specific antibodies were found in 67.6%, 26.8%, and 5.6% of the gilts, while 66.4% and 33.6% of gilts showed low and high levels of F18-specific antibodies. Hereby, we show that F4<sup>+</sup> and F18<sup>+</sup> *E. coli* are highly prevalent as potential enteropathogens in Cuban piggeries.

## 4.2 Introduction

F4<sup>+</sup> and F18<sup>+</sup> ETEC are common causes of diarrhea in suckling and recently weaned piglets worldwide. F18<sup>+</sup> VTEC causes edema disease, which is characterized by edema of the eyelids, ears, stomach, and colon as well as dyspnea, neurological symptoms, and sudden death (Oanh *et al.*, 2010). Zhang *et al.* (2007) implicated F4 and F18 as the major fimbrial antigen types among problems of ETEC and VTEC infections.

The pathogenesis of colibacillosis caused by F4<sup>+</sup> and F18<sup>+</sup> *E. coli* is well-known. The fimbriae mediate the adhesion of bacteria to specific receptors present on the surface of the enterocytes of the small intestine, leading to bacterial colonization and subsequent production of enterotoxins (LT, STa, and/or STb) which change the water and electrolyte balance across the mucosa leading to diarrhea. The edema disease is caused by the systemic vascular damage provoked by verocytotoxin (Stx2e; Fairbrother, 2006; Fairbrother and Gyles, 2006).

Swine colibacillosis causes losses due to mortality, costs of medication, and growth retardation, which have a significant negative impact on the economic feasibility of the swine industry (Lazo *et al.*, 2005; Blanco *et al.*, 2006; Tiels *et al.*, 2008). Housing and management conditions (e.g. hygiene, environment temperature, intake of maternal antibodies, and feeding), vaccination, and surveillance of antibiotic resistance are crucial in prevention of porcine neonatal and post-weaning diarrhea (Fairbrother *et al.*, 2005; Fairbrother, 2006).

In 2008, the average survival rate at weaning (number of weaned pigs/number of born pigs) on swine farms of the Cuban Company for Pork Production was as low as 70%. In Cuban piggeries, a specific etiologic diagnosis of the swine enteric pathogens is not always possible to perform. As a result, there is little information available on the prevalence of enteropathogens in Cuba where diarrheic diseases are responsible for 31% and 37% of the total mortality during the pre- and post-weaning periods, respectively (Cabrera and García, 2009).

F4- and F18-specific antibodies can be induced by parenteral vaccination, but also by an F4<sup>+</sup> and F18<sup>+</sup> ETEC/VTEC infection, respectively (Van den Broeck *et al.*, 1999a; Verdonck *et al.*, 2002). Vaccinations against F4 or F18 have not been applied in Cuba during the last decade; therefore, F4- or F18-specific serum antibodies in Cuban pigs will reflect the presence and spread of F4<sup>+</sup> ETEC or F18<sup>+</sup> ETEC/VTEC.


## 4.3 Materials and methods

### *Survey design*

This survey was performed on serum samples of replacement gilts (6 months old) originally collected by the Cuban Veterinary Diagnostic Laboratories for *Leptospira* and *Brucella* routine screening. Based on the experience of veterinary practitioners and a few reports of F4<sup>+</sup> and F18<sup>+</sup> *E.*

*coli* in Cuba, a prevalence between 60% and 90% of F4- and F18-specific antibodies was expected. Because a similar serosurvey has not been previously performed in Cuba, a 60% expected prevalence was selected as it gives a higher sample size ( $n = 1,025$  versus  $n = 385$  for the 90% expected prevalence). A 95% confidence interval and a low 3% desired absolute precision were fixed, resulting in a small confidence interval for theoretically infinite populations (Thrusfield, 1997; Naing *et al.*, 2006).

During the period between February and June 2008, 1,044 (19 more than planned) serum samples from farms of the Cuban Company for Pork Production (GRUPOR) were collected. In practice, they were distributed in 34 out of 94 piggeries of the company located in 10 out of 14 provinces all over the three Cuban regions (Fig. 1).



**Figure 1.** Map of Cuba showing the geographic location of the 34 piggeries (black dots) from where the prevalence of F4- and F18-specific antibodies were estimated. Roman numerals indicate the number of piggeries with similar location.

#### *Collection, transport, and treatment of blood serum samples*

Blood was collected from the orbital venous sinus (Fig. 2), and serum was obtained after 3 h incubation at environment temperatures (32-35°C) and stored at 4°C. Subsequently, centrifugation and heat inactivation were performed, followed by treatment with kaolin to remove nonspecific serum inhibitors. One volume of serum was mixed with four volumes of kaolin suspension [25% (w/v) in PBS (150 mM, pH 7.4)], and incubated at room temperature for 30 min. The serum was recovered after centrifugation at  $14,000\times g$  for 10 min and was stored at -20°C.



**Figure 2.** Puncture of the orbital venous sinus helped by a California needle in a pig, which is traditionally employed for blood collection or even for fluid-therapy in Cuba.

#### *Purification of F4 and F18 fimbriae*

The protocol was similar to the mechanical shearing described by Verdonck *et al.* (2002). F4 and F18 fimbriae were obtained from the *E. coli* strains IMM-01 (O149:K91/F4ac<sup>+</sup>/LT<sup>+</sup>/STb<sup>+</sup>) and 107/86 (O139:K12:H1/F18ab<sup>+</sup>/Stx2e<sup>+</sup>), respectively. Briefly, the *E. coli* strains were grown in TSB (Oxoid) at 37°C and 85 rpm overnight. Next, bacteria were collected by centrifugation (2851×*g*, 35 min, 4°C) and the pellet was resuspended and washed with PBS. Bacterial suspension was mixed on ice using an Ultra Turrax (IKA, Labortechnik) during 20 min. Afterwards, the bacterial cell debris was pelleted by centrifugation (26892×*g*, 40 min, 4°C, two times) and the fimbriae, contained in the supernatant, were precipitated with ammonium sulphate (at 40% saturation for F4 and at 20% saturation for F18, overnight, 4°C). The precipitated fimbrial suspension was centrifuged (1455×*g*, 30 min, 4°C), resuspended in PBS and dialyzed overnight (10 kDa cutoff) at 4°C against PBS. Finally, the quality of each fimbrial suspension was assessed by the Bicinchioninic acid reaction, 12% SDS-PAGE and Western Blot (Verdonk *et al.*, 2002).

#### *ELISA for detecting F4-specific immunoglobulins*

An indirect double-antibody ELISA was performed as previously described (Tiels *et al.*, 2008). Briefly, a solution of 1 µg/ml in PBS of IMM01 anti-F4 MAb (Van der Stede *et al.*, 2002) was applied as a coating to a 96-well MaxiSorp plate and incubated for 2 h at 37°C. The plates were blocked overnight with 0.2% (v/v) Tween® 80 in PBS at 4°C. Then, the F4 antigen was added at 5 µg/ml in dilution buffer [0.2% (v/v) Tween® 20 + 3% (w/v) BSA in PBS]. Next, the serum samples were applied in duplicates at a fixed dilution of 1/15, followed by the addition of anti-swine IgG (H+L) HRP-conjugated supplemented with 1% mouse serum. The presence of the bound conjugate was revealed by adding ABTS, and the OD was spectrophotometrically measured at 405 nm after incubation for 15 min at 37°C. The plates were washed three times [0.2% (v/v) Tween® 20 in PBS] between each

incubation step, except for the coating and blocking steps. The incubation condition of antigens and antibody solutions was for 1 h at 37°C.

#### *ELISA for detecting F18-specific immunoglobulins*

The main difference from the F4 ELISA was the coating with F18 fimbriae (2 µg/ml in 50 mM sodium bicarbonate pH 9.4; Tiels *et al.*, 2008), while the blocking, washing, incubation, and detection steps were similar.

The negative and positive control sera employed in both ELISAs were provided by the Laboratory of Immunology, Department of Virology, Parasitology and Immunology, Faculty of Veterinary Medicine, Ghent University, Belgium.

#### *Statistical analyses*

For data standardization, the average OD of the negative control serum per ELISA plate was subtracted from the OD value of each sample. In order to obtain a more suitable classification of individual results, we decided not to use the commonly used diagnostic OD cutoff point (i.e., mean OD + 3\*SD), but we modeled the homogenized data using a mixture of distribution.

It was assumed that pigs in the same location can be correlated; thus, three location variables (region, province, and piggery) were considered as grouping variables.

Let  $Y_{ijkl}$  be the vector of OD measurements, given the individual  $i$ , the region  $j$ , the province  $k$ , and the piggery  $l$ .

Firstly, we fit multilevel mixed-effects models, one for each data, using three levels of clustering, i.e. piggeries within provinces within regions, by means of SAS and R software, for the sake of comparison:

$$Y_{ijkl} = \mu + r_j + p_{jk} + f_{jkl} + \varepsilon_{ijkl}$$

, where the random effects  $r$ ,  $p$ , and  $f$  and the residual error  $\varepsilon$  are independently normally distributed with mean 0 and variance equal to  $\sigma_j^2$ ,  $\sigma_{jk}^2$ ,  $\sigma_{jkl}^2$  and  $\sigma_{ijkl}^2$ , respectively.

These models allow the mean OD estimation taking into account the additional variability due to the different clustering levels.

The correlation between F4 and F18 OD values was checked by the Pearson correlation coefficient with 95% confidence interval.

Secondly, in order to describe the overall distribution and to classify the individual OD values, we decided to fit the data using mixture models: the distribution of the data was fitted using a mixture of several Gaussian components, each of them representing a cluster of the data and having a

specific weight in the population. We fitted a series of Gaussian mixture models, increasing at each step the number of components up to 10. The analysis was done with the R package “mixAK” (Komárek, 2009), which estimates the mixture using Markov chain Monte Carlo (MCMC) methods. We used all the pigs pooled together, independently to their provenance: in this way, the classification of a pig in one of the components is only depending on its own OD value.

The choice of the optimal number of components was done using two selection criteria: the penalized expected deviance (PED; Plummer, 2008) and the difference in posterior deviances (Aitkin *et al.*, 2009). The PED penalizes the expected deviance  $\overline{D(\theta)}$  with the “optimism”  $p_{\text{opt}}$  because of using the same data twice, once for estimation and once for prediction: thus, the smaller the PED, the better. The approach of Aitkin (Aitkin, 2009) starts from the fact that models with growing numbers of parameters are automatically penalized by the increasing dispersion of the posteriors in their parameters. Thus, the selection between two models is given by the difference between the posterior distributions of the deviances:

$$\{D_{1,2}^{(m)} = D_1^{(m)} - D_2^{(m)} : m = 1, K, M\}$$

, where  $M$  is the length of the MCMC chain: we can derive from the posterior probability that Model 1 is better than Model 2,

$$\frac{1}{M} \sum_{m=1}^M I(D_{1,2}^{(m)} < 0)$$

, and derive 95% credible intervals for the difference in deviances.

#### 4.4 Results

##### *Exploratory data analysis: F4 OD data*

After data standardization, only two out of 1,044 samples resulted negative in the F4 ELISA, and the OD of the positives fluctuated from 0.017 to 1.887. The mean OD value of all negative control serum was 0.212 (n=56, SD=0.040). The analysis of these “raw” data shows a high positivity and differences between individual samples, but the proper description of the whole population is difficult to make, justifying the following analyses.

Next, the median F4 OD value and the interquartile range (IQR) per piggery were determined (Fig. 2). The median OD values were shown to be different between piggeries. Seven out of 34 piggeries had a high median OD value for F4-specific antibodies (above the third quartile for the entire sample, i.e., 1.01), whereas 18 out of 34 showed moderate levels (the median OD value was between the first

and third quartile, i.e., 0.53 and 1.01). Nine piggeries had little amount of F4-specific antibodies as the median OD value was below the first quartile.

Four piggeries (out of 7) with high median OD value for F4-specific antibodies are located in the Central province Ciego de Avila (E1-E4). Similar results were obtained for B2 (Matanzas, west region), and for G3 and G4 (Santiago de Cuba, east region).

The piggeries with moderate OD values were found to be spread over the country.

The lower median OD values were found in piggeries A2 (Havana City), C8 (Villa Clara), D3-D5 (Cienfuegos), F1-F3 (Las Tunas), and G2 (Santiago de Cuba). In most of them, the upper whisker, which contain OD values within  $1.5 \times \text{IQR}$ , was below the third quartile of the overall distribution, confirming the low level of F4-specific antibodies of sampled gilts.

#### *Exploratory data analysis: F18 OD data*

After data standardization, no negative values were found and the OD value of positives varied from 0.037 to 1.898. The mean OD value of all negative control serum was 0.106 ( $n=52$ ,  $SD=0.020$ ).

The distribution of F18 OD values between piggeries (Fig. 3) is more homogeneous compared to the F4 data as most piggeries (23 out of 34) had median F18 OD values between the first and the third quartiles of the overall distribution, which are equal to 0.51 and 1.01, respectively.


However, a high median OD value for F18-specific antibodies (above the third quartile, i.e., 1.01) was obtained in piggeries A1 (Havana City), C2 and C5 (Villa Clara), as well as E2 and E4 (Ciego de Avila).

Piggeries with the lowest median OD values are located in Matanzas (B1 and B2, west region), in Villa Clara (C1 and C8, central region) and in the eastern province Las Tunas (F2 and F3).


#### *Random-effects models*

Both multilevel models fitted in SAS and in R for F4 and F18 OD data confirm that most of the variance is residual or unexplained; there is some extra variation due to piggery, but very little due to province or region (Table 1). In particular, not all F4 variability is residual, and some heterogeneity can be ascribed to a smaller extent to the piggeries. Conversely, for F18, only the variance of the residual variability is significantly positive.





**Figure 2.** Box plot of the ELISA OD values for F4-specific antibodies between piggeries. The *boxes* contain OD values between the first and the third quartiles, and the *whiskers* contain OD values within  $1.5 \times \text{IQR}$ . The *horizontal dashed lines* indicate the first and the third quartiles of the overall distribution. The *line inside the boxes* represents the median OD value. In the *x-axis*, piggeries are organized by province from West to East Cuba: A1 and A2 (Havana City), A3 and A4 (Havana), B1 and B2 (Matanzas), C1-C8 (Villa Clara), C9 (Sancti Spiritus), D1-D5 (Cienfuegos), E1-E5 (Ciego de Avila), F1-F4 (Las Tunas), F5 (Holguín), and G1-G4 (Stgo de Cuba).



**Figure 3.** Box plot of the ELISA OD values for F18-specific antibodies between piggeries. The *boxes* contain OD values between the first and the third quartiles, and the *whiskers* contain OD values within  $1.5 \times \text{IQR}$ . The *horizontal dashed lines* indicate the first and the third quartiles of the overall distribution. The *line inside the boxes* represents the median OD value. In the *x-axis*, piggeries are organized by province from West to East Cuba: A1 and A2 (Havana City), A3 and A4 (Havana), B1 and B2 (Matanzas), C1-C8 (Villa Clara), C9 (Sancti Spiritus), D1-D5 (Cienfuegos), E1-E5 (Ciego de Avila), F1-F4 (Las Tunas), F5 (Holguín), and G1-G4 (Stgo de Cuba).

**Table 1.** Estimated mean ( $\mu$ ) and variance components (with standard errors) of the multilevel linear mixed model fixed for the F4 and F18 ELISA OD data using three levels of clustering (piggeries within provinces within regions).

Parameters	F4 OD data	F18 OD data
$\mu$	0.80 (0.22)	0.80 (0.034)
Region	0	0
Province	0.021 (0.017)	0
Piggery	0.047 (0.0030)	0.040 (0.044)
Residual	0.076 (0.00086)	0.10 (<0.0001)

The study of Pearson correlation coefficient ( $r = 0.19$ ) in R suggested that there is no correlation between F4 and F18 OD individual values, confirming the specificity of both ELISAs. Indeed, clear differences are observed when comparing the distribution of F4 and F18 OD values in the piggeries A1, B2, C5, C9, and E1 (Figs. 2 and 3).

#### *Mixture modeling: F4 OD data*

According to the PED, the best mixture has three components (PED=788), while the one with four components has a PED of 823. According to the difference in posterior deviances, the mixture with three components is much better than the mixture with two components, with  $D_{1,2}^{(m)} = -19$  and 95% credible interval (-29,-5) and posterior probability 0.99. Instead, we cannot confidently choose between the mixture with four components and with three, with  $D_{1,2}^{(m)} = 1$  and 95% credible interval (-17, 18) and posterior probability 0.57. Thus, we choose the mixture with three components, because it almost fits in the same way as the one with four components, being at the same time more parsimonious in terms of parameters.

Looking at the estimates (Table 2), we can label the components in the following way: the component 1 contains the low OD values, and the components 2 and 3 stand for the moderate and high OD values, respectively. Then, all the observations are rather assigned to a particular component, instead of being spread among all the population.

Figure 4 presents the histogram of the data over imposing the estimated mixture with three components. We can notice that the biggest components, the ones for low and moderate OD values, are gathering the serum levels from 0 up to 1.5, while after that value, the third component is taking place and gathers the highest OD values.

For separating the three components, the mixture OD cutoff points were estimated at their intersection: the OD cutoff point between the lowly and the moderately positive pigs was 0.910, while between the moderately and highly positives was 1.551.


Then, we can accurately determine how individual cases could be classified according to their ELISA signal as the cutoff OD values were obtained from the overall distribution (Table 3).

The prevalence of gilts with low, moderate, and high levels of F4-specific antibodies was 67.6%, 26.8%, and 5.6% respectively.

**Table 2.** Mixture parameters estimate of the fixed Gaussian models for the F4 and F18 ELISA OD data in which are based the definition of anti-F4 and anti-F18 OD groups.

OD data	Components	$\pi_k$	$\mu_k$	$\sigma_k$	OD groups
F4	1	0.599	0.614	0.224	Low
	2	0.349	1.036	0.283	Moderate
	3	0.052	1.683	0.117	High
F18	1	0.602	0.578	0.231	Low
	2	0.398	1.084	0.329	High

$\pi_k$ , estimate for the mixing weights or prior probability;  $\mu_k$ , mean;  $\sigma_k$ , standard deviation.



**Figure 4.** Histogram of the F4 ELISA OD data with the fixed three-component Gaussian mixture model (*thick curve*) overimposed, assessing the prevalence of F4-specific antibodies in Cuban swines. The *thin curves* show the three single components, and the *dashed lines* indicate the mixture OD cutoff points (0.910 and 1.551).

**Table 3.** Seroprevalence of F4-specific antibodies in Cuban piggeries according to the ELISA OD cutoff points calculated by the fixed three-component mixture model.

Piggeries	Number	Seroprevalence (%)		
		Low OD	Moderate OD	High OD
A1	17	71	29	0
A2	15	100	0	0
A3	76	91	9	0
A4	12	67	33	0
B1	29	100	0	0
B2	49	22	78	0
C1	31	77	19	3
C2	59	54	34	12
C3	73	44	52	4
C4	8	50	50	0
C5	14	93	7	0
C6	40	65	25	10
C7	28	79	21	0
C8	9	100	0	0
C9	26	35	35	31
D1	10	90	10	0
D2	9	89	11	0
D3	58	98	2	0
D4	8	100	0	0
D5	4	100	0	0
E1	48	8	50	42
E2	2	0	100	0
E3	8	25	63	13
E4	15	20	40	40
E5	27	100	0	0
F1	50	84	16	0
F2	52	87	13	0
F3	45	89	11	0
F4	60	63	35	2
F5	40	80	20	0
G1	30	83	17	0
G2	34	97	3	0
G3	46	22	65	13
G4	12	33	58	8
Total	1,044	67.6	26.8	5.6

Shaded lines represent the highly positive piggeries, where more than 50% of sows showed moderate or high OD values


### Mixture modeling: F18 OD data

According to the PED, the best mixture has two components (PED=812), while the second best has six components (PED=852). Instead, according to the difference in posterior deviances, it is quite difficult to choose the best mixture. The posterior probabilities  $P(\text{diff} < 0)$  increase as long as the number of components increases. For instance, if we compare the model with six and with two components, the posterior probability for the model with six groups is 0.70, with  $D_{1,2}^{(m)} = 5$  and 95% credible interval (-16, 27), while if we compare the models with three and with two components, the posterior probability for the model with three groups is 0.58, with  $D_{1,2}^{(m)} = 2$  and 95% credible interval (-16, 27). However, we notice that the mean of the difference is always positive and that the credible interval is more shifted toward the positive values. Given all this, we choose the mixture with two components, which is even indicated as the best one by the PED criterion.

Looking at the estimates (Table 2), the component 1 represents the group of gilts with the lower amount of F18-specific antibodies, whereas component 2 represents gilts with the higher amount of F18-specific antibodies.

Figure 5 presents the histogram of the data overimposing the estimated mixture with two components. The OD cutoff point determined by the intersection of both components was 0.894.

From the overall sample, we determined that 66.4% and 33.6% of gilts showed low and high OD values to F18-specific antibodies, respectively (Table 4).



**Figure 5.** Histogram of the F18 ELISA OD data with the fixed two-component Gaussian mixture model (*thick curve*) overimposed, assessing the prevalence of F18-specific antibodies in Cuban swines. The *thin curves* show the two single components, and the *dashed line* indicates the mixture OD cutoff point (0.894).

**Table 4.** Seroprevalence of F18-specific antibodies in Cuban piggeries according to the ELISA OD cutoff point calculated by the fixed two-component mixture model.

Piggeries	Number	Seroprevalence (%)	
		Low OD	High OD
A1	17	41	59
A2	15	87	13
A3	76	63	37
A4	12	50	50
B1	29	79	21
B2	49	96	4
C1	31	94	6
C2	59	34	66
C3	73	85	15
C4	8	25	75
C5	14	0	100
C6	40	65	35
C7	28	46	54
C8	9	100	0
C9	26	77	23
D1	10	50	50
D2	9	100	0
D3	58	57	43
D4	8	63	38
D5	4	50	50
E1	48	54	46
E2	2	0	100
E3	8	50	50
E4	15	13	87
E5	27	100	0
F1	50	64	36
F2	52	83	17
F3	45	80	20
F4	60	65	35
F5	40	70	30
G1	30	80	20
G2	34	85	15
G3	46	39	61
G4	12	50	50
Total	1,044	66.4	33.6

Shaded lines represent the highly positive piggeries, where more than 50% of sows showed high OD values

#### 4.5 Discussion

In this survey, the prevalence of F4- and F18-specific antibodies in sera of 1,044 Cuban replacement gilts was assessed. By ELISA and diverse statistical analyses, a high seroprevalence of specific antibodies against the F4 and F18 fimbriae was confirmed.

Spread over the whole country, 26.8% and 5.6% of gilts were moderately and highly positives for F4-specific antibodies, respectively, and 33.6% were highly positives for F18-specific antibodies. Since a commercial vaccine against F18 is not available, and since vaccination of sows against F4<sup>+</sup> ETEC has not been practiced anymore during the last 15 years in Cuba (Wong *et al.*, 1995), our results show that F4<sup>+</sup> and F18<sup>+</sup> *E. coli* are highly prevalent as potential enteropathogens in Cuban piggeries. An F4<sup>+</sup> or F18<sup>+</sup> ETEC/VTEC infection provokes induction of F4- and F18-specific serum antibodies (Verdonck *et al.*, 2002).

Serological surveys using ELISA showed its reliability for testing vaccines and for epidemiological studies regarding fimbriae carrying *E. coli*. In Belgium, a seroprevalence study performed on sows showed that in West-Vlaanderen province, 79% of non-vaccinated farms were seropositive for F4-specific antibodies, while in Vlaams-Brabant only 31% were seropositive (Van den Broeck *et al.*, 1999b). The herd seroprevalence of F18-specific antibodies was higher in open system Belgian piggeries (96.4%) than in the closed system ones (88.8%; Verdonck *et al.*, 2003).

Specific diagnosis of swine colibacillosis is limited and out-of-date in Cuba. By using slide agglutination and direct immunofluorescence, Pedroso and Talavera (1983) detected F4<sup>+</sup> and F5<sup>+</sup> *E. coli* in fecal samples of diarrheic and healthy piglets in Havana province. In 1998, applying a whole-bacterial cell ELISA, F4<sup>+</sup> *E. coli* were identified in the 2.8% of diarrheic piglets from farms where sows were vaccinated against F4 and F5 in Camagüey province (Fuentes *et al.*, 2001). Lazo *et al.* (2005) reported 43.8% morbidity and 12.8% mortality due to enteric colibacillosis clinically diagnosed in Villa Clara province, Cuba.

After more than a decade, this is the first study that refers to a high prevalence of F4<sup>+</sup> *E. coli* in Cuba, specifically in the central region, whereas Blanco *et al.* (2006) did not find F4<sup>+</sup> *E. coli* in diarrheic piglets from the same region. To our opinion, they used a small sample size, and differences between their bacteriological (direct) and our immunological (indirect) identification are obvious. However, the high seroprevalence of F18-specific antibodies reported in our survey, with 33.6% serum samples showing high OD values, correlates with their report of 61% occurrence of F18 fimbriae among *E. coli* isolates.

Recent surveys carried out in China (Cheng *et al.*, 2006), USA (Zhang *et al.*, 2007), Zimbabwe (Madoroba *et al.*, 2009), Brazil (Vidotto *et al.*, 2009), and Vietnam (Oanh *et al.*, 2010) showed virulent *E. coli* carrying genes encoding F4 or F18 fimbriae as commonly associated pathogens with piglet's diarrhea or edema disease.

As colibacillosis is a multifactorial disease, our findings could not be directly associated with disease. However, in Poland, Osek (1999) found a higher prevalence of virulent *E. coli* in diarrheic weaned pigs than in healthy pigs, while F4<sup>+</sup> *E. coli* were only present in the diarrheics (19.1% prevalence). In China, Cheng *et al.* (2005) found a 58.33% occurrence of F18 fimbriae in *E. coli* strains isolated from pigs with diarrhea or edema disease.

After analyzing the high positivity to F4- and F18-specific antibodies obtained in our survey and based on previous studies that have been done in Belgium, a high genetic susceptibility of the Cuban swines to colonization by F4<sup>+</sup> and F18<sup>+</sup> *E. coli* could be inferred. By the in vitro adhesion assay, the 80% of Belgian pigs were classified in the F4 susceptible phenotype A (F4acR<sup>+</sup>), and only 4% in the resistant phenotype E (Cox and Houvenaghel, 1987); it supports, in part, that in Flanders, 65% of non-vaccinated piggeries were seropositive for F4 (Van den Broeck *et al.*, 1999b). The 92% of Belgian pigs belonged to the FUT1M307G/G or FUT1M307G/A genotypes, both corresponding with sensitivity to F18<sup>+</sup> *E. coli* infections (Coddens *et al.*, 2007), which justified the high seroprevalence of F18-specific antibodies previously reported by Verdonck *et al.* (2003).

A low positive ELISA signal for F4- and F18-specific antibodies was obtained in 67.6% and 66.4% of the samples, respectively. Presumably, it could be the result of a weak bacterial infection determined by a good management and hygiene or by the prevalence of weak adhesion phenotypes. Yan *et al.* (2009) reported in China that the weak adhesion phenotypes of F4 (not yet described in Cuba) are present at a high frequency and mainly in crossbreds. As reported by Coddens *et al.* (2007) in F18 genetically susceptible piglets, weak and strong susceptible adhesion phenotypes were present in piglets aged from 1.5 to 23 weeks old.

The use of a vaccine against both F4 and F18 fimbrial adhesins could be one of many ways to control colibacillosis in Cuba. Tiels *et al.* (2008) recommended the FedF adhesin in combination with the F4 fimbriae as a good oral vaccine candidate, but more research on mucosal immunization is required.

In conclusion, this serological survey for F4- and F18-specific antibodies indicates that F4- and F18-fimbriated *E. coli* are widespread as potential etiologic agents of swine neonatal or post-weaning diarrhea and edema disease in Cuba. From the epidemiologic and zootechnic point of views, implementation of a national program for the prevention and treatment of swine colibacillosis appears advantageous.



#### **4.6 Acknowledgments**

This study was funded by an IUC program between VLIR-UOS and Universidad Central “Marta Abreu” de Las Villas. The authors thank the Cuban Company for Pork Production and the Cuban Institute of Veterinary Medicine for their support and help during the sera collection. Are specially thanked the veterinarians from the Serology Department of the Veterinary Diagnostic Laboratories located in Havana, Jovellanos (Matanzas), Santa Clara, Cienfuegos, Sancti Spiritus, Ciego de Avila, Las Tunas, Holguín, and Santiago de Cuba.

## Chapter 5

---

*Screening commercial pigs in Villa Clara province,  
Cuba, for mucin 4 polymorphisms and  
susceptibility/resistance to F18<sup>+</sup> E. coli*



### 5.1 Abstract

F4 and F18 fimbriae frequently mediate the adhesion of pathogenic *E. coli* to specific receptors on the gut epithelium of pigs. The phenotypic expression of susceptibility or resistance to adhesion of F18 to the surface of enterocytes has been linked to *Cfol* polymorphism in the *FUT1* gene (F18R). For F4, *Xba*I polymorphism in the porcine *mucin 4* gene has been linked with presence of the F4R, but not with absence. In order to know the frequency of genotypes and alleles determining susceptibility or resistance to F18 and *Xba*I polymorphism in the *mucin 4* gene in pigs raised in Villa Clara province, Cuba, genomic DNA of 90 F1 (Yorkshire x Landrace) x Duroc commercial pigs was tested by PCR-RFLP. Overall, the *Xba*I-resistant genotype was frequently detected (0.66), and the heterozygous or homozygous *Xba*I-digestible genotypes less frequently (0.31 and 0.03, respectively). These data compared with a previous survey reporting a high seroprevalence of F4-specific antibodies in sera of young sows in Cuba, confirm that *Xba*I polymorphism cannot be used as marker for F4R expression and supports the presence of other specific markers and or receptors. Pigs carrying the resistant genotype (AA) to F18<sup>+</sup> *E. coli* infections were not frequently found (0.13), and the susceptible (G) allele was highly frequent (0.73), coinciding with previous reports of a high prevalence of F18 fimbriae and antibodies against it in the Cuban swine herd.

## 5.2 Introduction

F4 and F18 fimbriae frequently mediate the adhesion of ETEC and F18 also of VTEC to specific receptors present on the surface of porcine enterocytes, leading to bacterial colonization and production of toxins, which provoke diarrhea (ETEC) or edema disease (VTEC; Fairbrother, 2006; Fairbrother and Gyles, 2006). Zhang *et al.* (2007) reported F4 and F18 as the major fimbrial antigens among *E. coli* isolates associated with diarrhea in young pigs in U.S.A.

DNA based-tests determining resistance or susceptibility of pigs to *E. coli* infections are important tools for breeding programs and for epidemiological studies undertaken to decrease the prevalence of swine colibacillosis in a herd. An *Xba*I polymorphism in intron 7 of the porcine *mucin 4* gene has been linked to the F4ab/ac fimbriae adhesive phenotype; by PCR-RFLP three genotypes of pigs regarding F4ab/ac mediated *E. coli* adhesion can be distinguished: resistant (RR), susceptible heterozygote (SR), and susceptible homozygote pigs (SS; Jørgensen *et al.*, 2004), however Rasschaert *et al.* (2007) demonstrated that the RR genotype also contained F4 susceptible pigs.

The F18R status of pigs is genetically determined (Bertschinger *et al.*, 1993), and susceptibility to *E. coli* colonization favored by F18 fimbriae appeared to be dependent on the activity of the *FUT1* gene, which encodes the alpha(1,2)-fucosyltransferase (Meijerink *et al.*, 1997; 2000). The discovery of the close linkage of a G→A polymorphism at nucleotide 307 on the *FUT1* gene and the F18R in porcine enterocytes has made the development possible of a PCR-RFLP using *Cfo*I enzyme digestion for identifying pigs resistant or susceptible to F18 adherence. Sequencing of the *FUT1* gene of pigs resistant (AA) to F18 revealed a transition G→A in both alleles on bp 307, resulting in an amino acid substitution at position 103 (Ala→Thr), different from the susceptible (GA or GG) pigs (Meijerink *et al.*, 1997).

Recently, a high seroprevalence of specific antibodies against F4 and F18 fimbriae was determined in the Cuban swine population (Chapter 4; de la Fé Rodríguez *et al.*, 2011). Furthermore, both fimbriae were most frequently identified on pathogenic *E. coli* isolated from suckling and weaned pigs with diarrhea in the Villa Clara province (Chapter 2). As a consequence, pigs raised in Cuba should have a high genetic susceptibility to these infections. Therefore we analyzed the receptor polymorphism for F18 fimbriae and determined *Xba*I polymorphism in the *mucin 4* gene of pigs raised in commercial farms located in the Villa Clara province.

## 5.3 Materials and methods

### *Survey design and collection of blood samples*

Blood was collected from the orbital venous sinus of 90 commercial pigs F1 (Yorkshire x Landrace) x Duroc equally distributed (15 per piggery) in six large continuously farrowing piggeries (namely

from A to F) located in Villa Clara province, Cuba. One ml of blood was gently mixed with 100 µl of 10% EDTA to prevent coagulation and was kept at -20°C after arrival to the laboratory. These pigs were also surveyed in Chapter 2 of this thesis.

#### *Extraction of genomic DNA*

Red blood cells were lysed by mixing 100 µl of blood with 1 ml of TE buffer (10 mM Tris-HCL + 1 mM EDTA, pH 7.5), this mixture was centrifuged at 3800xg for 20 sec, and the pellet was washed twice under the same conditions. Next, 2 µl of proteinase K (20 mg/ml, Invitrogen) diluted in 200 µl of K-buffer (50 mM KCL + 20 mM Tris-HCL + 2.5 mM MgCl<sub>2</sub> + 0.5% Tween® 20, pH 8.3) were added to the pellet, and incubation was performed at 56°C for 1 h in a warm water bath. Next, proteinase K was inactivated by incubation at 95°C for 10 min, centrifugation was performed at 3800xg for 20 sec, and the obtained supernatant was stored at -20°C, and used as template DNA for PCR analyses.

#### *DNA test for mucin 4 polymorphism*

PCR-RFLP was performed as described by Jørgensen *et al.* (2004). Briefly, the PCR reaction (20 µl) which allows the amplification of the *mucin 4* gene fragment contained 5 pmol of each primer (5'-GTGCCTTGGGTGAGAGGTTA-3'/5'-CACTCTGCCGTTCTCTTTCC-3'), 200 µM of each dNTPs (Roche Diagnostics GmbH), 2 mM of MgCl<sub>2</sub> (Promega), 0.5 U of SuperTaq (HT Biotechnology Ltd.), 2 µl of 10xSuperTaq buffer, and 5 µl of template DNA. The PCR amplifications were performed in a thermal cycler PTC-100™ (MJ Research Inc.) by incubation of the mixture at 95°C for 5 min for pre-denaturation, followed by 39 cycles of denaturation at 95°C for 15 sec, annealing at 65°C for 30 sec, and extension at 72°C for 1 min, with a final extension at 72°C for 7 min. Finally, 5.5 µl of the PCR product (367 bp) were digested with *Xba*I (Promega) for 2 h at 37°C in warm water bath. One allele (R) is resistant to *Xba*I digestion, whereas the S allele is digested into a 151 bp and a 216 bp fragment.

#### *DNA marker-based test for the F18R*

The PCR reaction (30 µl) for the amplification of the *FUT1* gene fragment contained 15 pmol of each primer (5'-CTTCCTGAACGTCTATCAAGACC-3'/5'-CTTCAGCCAGGGCTCCTTTAAG-3'), 200 µM of each dNTPs (Roche Diagnostics GmbH), 1.5 mM of MgCl<sub>2</sub> (Promega), 0.5 U of SuperTaq (HT Biotechnology Ltd.), 3 µl of 10xSuperTaq buffer, and 5 µl of template DNA. The PCR amplifications were performed by incubation of the mixture at 95°C for 3 min for pre-denaturation, followed by 35 cycles of denaturation at 95°C for 30 sec, annealing at 61°C for 30 sec, and extension at 72°C for 30 sec, the final extension was at 72°C for 10 min. In order to detect the G→A polymorphism at

nucleotide 307 of the *FUT1* gene described by Meijerink *et al.* (1997), ten µl of the PCR product (421 bp) were digested with *CfoI* (Promega) for 2 h at 37°C in a warm water bath, resulting in fragments of 328 and 93 bp for the resistant (A) allele, and fragments of 241, 93, and 87 bp for the susceptible (G) allele.


In every piggery and in the overall sample the genotypes frequency was determined dividing the number of every genotype by the number of tested pigs; the alleles frequency was determined dividing the number of every allele by the total number of alleles in every group.

#### 5.4 Results and discussion


The *mucin 4* and *FUT1* genes fragments were successfully amplified in all pigs, and the digestion with *XbaI* and *CfoI* enzymes, respectively, were reliable for detecting polymorphisms in both of them, allowing the detection of susceptible and resistant pig genotypes to *E. coli* colonization mediated by F4 or F18 fimbriae (Fig. 1, 2, and 3).

##### *Mucin 4 gene polymorphisms*


As shown in Table 1, pigs homozygous for indigestible alleles (RR) are highly frequent (0.66), Jørgensen *et al.* (2004) associated this status to resistance to *E. coli* adhesion mediated by F4ab/ac fimbriae. However, in our previous study we determined a high seroprevalence of F4-specific antibodies in gilts from Cuban piggeries (Chapter 4; de la Fé Rodríguez *et al.*, 2011). These findings support findings of Rasschaert *et al.* (2007) that the F4R should not only be linked to the activity of the *mucin 4* gene. Indeed, Rasschaert *et al.* (2007) found that F4ab/ac fimbriae could adhere to villous brush borders of pigs that carry only *XbaI*-indigestible alleles (RR), and reported the absence of the *XbaI* restriction site in the *mucin 4* gene fragment of 33.3% pigs positive for the adhesion of *E. coli* expressing F4ab/ac fimbriae. Although Erickson *et al.* (1992; 1994) proposed the intestinal mucin-like sialoglycoproteins (IMTGPs) as F4R, also other molecules have been proposed as receptors such as an intestinal transferrin (GP74; Grange and Mouricout, 1996), an intestinal glycosphingolipid (Grange *et al.*, 1999) and the porcine aminopeptidase-N (Rasschaert, 2008) have been proposed as F4R.



**Figure 1.** Agarose gel electrophoresis showing examples of the PCR product obtained from the amplification of the *FUT1* (421 bp; samples 1, 2, and 3) and *mucin 4* (367 bp; samples 4, 5, and 6) gene fragments.



**Figure 2.** Agarose gel showing examples of the product obtained after *XbaI* digestion of the *mucin 4* gene fragment. Pig 1 is homozygous for digestible alleles (SS). Pigs 3, 4, and 5 carry indigestible alleles (RR), and pig 2 is heterozygous (SR).



**Figure 3.** Agarose gel electrophoresis showing examples of the product obtained after *CfoI* digestion of the amplified *FUT1* gene fragment. Pigs 1 and 7 carry the GA susceptible genotype. Pigs 2, 3, 4, and 6 carry the GG susceptible genotype, and pig 5 carry the AA resistant genotype.



**Table 1.** The frequency of pigs carrying homozygous *mucin 4* gene for the presence of *Xba*I restriction site (SS), homozygous for its absence (RR), and heterozygous (SR) in the Villa Clara province, Cuba.

Piggery	Genotypes frequency of <i>Xba</i> I digestion patterns of the <i>mucin 4</i> gene			Alleles frequency	
				<i>Xba</i> I-indigestible	<i>Xba</i> I-digestible
	RR	SR	SS	R	S
A	0.80	0.20	0	0.90	0.10
B	0.60	0.40	0	0.80	0.20
C	0.93	0.07	0	0.97	0.03
D	0.40	0.53	0.07	0.67	0.33
E	0.73	0.20	0.07	0.83	0.17
F	0.47	0.47	0.07	0.70	0.30
Total	0.66	0.31	0.03	0.81	0.19

*FUT1* gene polymorphisms and *F18R*

Pigs carrying the resistant genotype (AA) to F18 were not frequently found (0.13; Table 2). Similar results were obtained by Shi *et al.* (2002) in 158 pigs of Yorkshire, Landrace, and Duroc breeds in China, where the frequency of the resistant genotype AA was 0.11 and the frequency of the susceptible genotypes GA and GG were 0.35 and 0.54, respectively. In 74 Belgian Landrace x Pietrain or Dutch Landrace an even lower frequency of the resistant genotype AA (0.08) was detected and the frequency of the susceptible genotypes GA and GG were 0.37 and 0.55, respectively (Coddens *et al.*, 2007).

**Table 2.** The frequency of F18 susceptible (GG, GA) and resistant (AA) genotypes identified by the *Cfo*I polymorphism in the *FUT1* gene of 90 pigs in the Villa Clara province, Cuba.

Piggery	Genotypes frequency			Alleles frequency	
	AA	GA	GG	A	G
A	0	0.40	0.60	0.20	0.80
B	0.07	0.27	0.67	0.20	0.80
C	0.27	0.20	0.53	0.37	0.63
D	0.13	0.13	0.73	0.20	0.80
E	0.13	0.13	0.73	0.20	0.80
F	0.20	0.47	0.33	0.43	0.57
Total	0.13	0.27	0.60	0.27	0.73

The high genetic susceptibility for adhesion of F18<sup>+</sup> *E. coli* in Cuban pigs in our study is in agreement with the high prevalence of F18-specific antibodies in young sows we reported previously (Chapter 4; de la Fé Rodríguez *et al.*, 2011) and with the high occurrence (61%) of the F18 coding gene in *E. coli* isolated from piglets with diarrhea in Central Cuba in a study of Blanco *et al.* (2006) and in ETEC and VTEC strains reported in Chapters 2 and 3 of this thesis.

The presence of the allele A determining F18 resistance at the frequency of 0.27 in commercial pigs is important for future breeding programs aiming the selection of F18-resistant breeding stocks in order to prevent diarrhea or edema disease in Cuba.

It is interesting that Chinese native pig breeds (Taihu, Huai) lacking genetic factors providing resistance to F18 fimbriae, have stronger resistance to post-weaning diarrhea and edema disease compared with western pig breeds due to a reduced growth rate of the Chinese pigs (Bao *et al.*, 2008).

At least for what F18<sup>+</sup> *E. coli* control is concerned, positive selection of Cuban pigs for the AA F18-resistant genotype could contribute to decrease swine diarrhea and prevent edema disease.

## **5.5 Acknowledgments**

We thank especially Dr. Annelies Coddens and Ir. Phillip Bellot from the Laboratory of Immunology, Faculty of Veterinary Medicine, Ghent University, for their assistance in detecting polymorphisms in the *mucin 4* and *FUT1* genes. This study was carried out in the frame of an IUC program between VLIR-UOS and Universidad Central “Marta Abreu” de Las Villas.



## Chapter 6

---

*General discussion and future perspectives*



### 6.1 General discussion

Cuban swine veterinary practitioners often complain about the high occurrence and yearly increased incidence of diarrhea in young pigs, as well as about the limitations for the specific identification of its etiology (e.g. enteropathogens or lack of food/water quality), which results in the application of insufficient prevention/control measures at national level like the common uncontrolled administration of antibiotics (Cabrera *et al.*, 2010). Diarrhea diagnosis in Cuban piggeries and provincial Veterinary Diagnostic Laboratories is frequently restricted to clinical and macropathological examinations (Cabrera and García, 2009), which are a tentative rather than a specific diagnostic tool (Elicker *et al.*, 2010). Indeed, the clinical picture of porcine neonatal or post-weaning diarrhea often do not point to a likely etiology because of the multifactorial nature of this disease and the variation in clinical signs and epidemiology that can be produced by the effect of one or more disease agents (Straw *et al.*, 2006). In 2008, the Cuban Institute for Swine Research reported gastroenteric diseases to cause 31% and 37% of the total piglet's mortality during the pre-weaning and post-weaning periods, respectively. Also, other causes of piglet's mortality related with diarrhea, like crushing or hypoglycemia, frequently occur in Cuban piggeries (Cabrera and García, 2009). Moreover, research, technology transfer, and laboratory protocols contributing to the epidemiological characterization of swine enteric infections have been limited during the last two decades and are out of date in Cuba (Pedroso and Talavera, 1983; Cabrera and García, 1985; Koudela *et al.*, 1989; NRAG, 1989; Fuentes *et al.*, 2001; Barrera *et al.*, 2005; Blanco *et al.*, 2006).

The epidemiology of any infectious syndrome is a wide matter to deal with, specifically when previous information is almost completely lacking. We had thus first to identify the pathogens currently associated with piglet's diarrhea in the Villa Clara province of Cuba. Then, pathogenic *E. coli*, which were the most frequent enteropathogen, were characterized for virulence factors, antibiotic susceptibility/resistance, and genetic profiles. Later, a seroprevalence study of antibodies against F4 and F18 fimbriae all over Cuba contributed to the epidemiological picture of *E. coli* commonly associated with piglet's diarrhea. Finally, the enteric receptor status for F18 fimbriae and the *Xba*I polymorphisms in the porcine *mucin 4* gene linked with presence, but not with absence of the F4R, of commercial pigs in Cuba were tested by DNA based-markers.

Failure of isolation and identification of certain enteropathogens actively involved in the pathogenesis of diarrhea in a swine herd could lead to the failure of a control program. Diseases such as TGE, coccidiosis, rotavirus, colibacillosis and clostridial enteritis are difficult to eliminate once they have become a problem in the herd; and they may still contribute to diarrhea due to failures in preventive programs (antibiotic resistance, vaccinations, sanitation, or management; Straw *et al.*, 2006). We observed that 64.4% and 42.2% of suckling piglets and weaned pigs with diarrhea, respectively, were infected with at least one of the following enteropathogens: ETEC, VTEC,  $\alpha$ -

toxigenic *C. perfringens*, *S. Newport*, TGEV, rotavirus A, *C. parvum* or *I. suis*, and their frequency ranged from 3.3% for VTEC to 25.6% for ETEC. Important enteropathogens like  $\beta$ -toxigenic *C. perfringens*, PEDV, *Eimeria* spp., and helminths were not detected, but we cannot confirm their absence in Cuba or even in Villa Clara as limited numbers of pigs during a limited period of time were surveyed. Swine helminths are common in Cuba as a tropical country: *Hyostrongylus rubidus*, *Strongyloides* sp., *Ascaris suum*, *Macracanthorhynchus hirudinaceus*, *Oesophagostomum* sp., *Trichuris suis*, and even *Fasciola hepatica* have recently been reported in pigs raised outdoors by private farmers in the Villa Clara province (de la Fé Rodríguez *et al.*, 2007).

Eight kinds of combined infections were identified in 25% of enteropathogen positive pigs. To our point of view, mixed infections have to be accurately identified to plan and implement an efficient disease control. Worldwide few controlled experiments have been designed to study the interaction between swine enteropathogens (Chapter 1). Lecce *et al.* (1982) and Enemark *et al.* (2003a) reported that a mixed infection rotavirus-ETEC or rotavirus-*Cryptosporidium*, respectively, causes a dramatic aggravation of diarrhea and clinical signs while mild clinical signs of illness were observed in piglets mono-infected with rotavirus, ETEC or *Cryptosporidium*. In Japan, Ushida *et al.* (2009) closely associated mixed infections with diarrhea as 42% of diarrheic weaned pigs had multi-infections, whereas such infections only occurred in 17% of weaned pigs with solid feces, and 33 % of diarrheic suckling piglets showed mixed infections, whereas they were absent in piglets without scour.

In this study we found that pathogenic *E. coli* were commonly associated with porcine pre- or post-weaning diarrhea. Seventeen clones of pathogenic *E. coli* isolated from diarrheic piglets carried genes for virulence factors F4, F5, F6, F18, F41, STa, STb, STx2e, or LT. F4<sup>+</sup> and F18<sup>+</sup> ETEC were the most frequently identified and they showed multi-resistance to most of antibiotics routinely administered in Cuban piggeries (i.e. tetracycline, ampicillin, sulphonamides, and kanamycin) in contrast to the F18<sup>+</sup>/STx2e<sup>+</sup> isolates (clonal group XI) which were only resistant to sulphonamides. These are important findings to manage in Cuban piggeries because of the fact that these pathogenic bacteria were isolated from diarrheic pigs, and they have been associated with clinical disease worldwide: in Poland, Osek (1999) isolated F4<sup>+</sup> *E. coli* only in diarrheic weaned pigs (19.1% prevalence), and in China, Cheng *et al.* (2005) reported a 58.33% occurrence of F18<sup>+</sup> *E. coli* in pigs suffering from diarrhea or edema disease. Boerlin *et al.* (2005) and Travis *et al.* (2006) reported linkages of antibiotic resistance and virulence genes on plasmids in porcine ETEC; consequently, uncontrolled administration of antibiotics, like is happening in Cuban piggeries, may contribute to the selection, persistence, and spreading of pathogenic *E. coli*. Therefore, the fact that ETEC showed susceptibility to several antibiotics less or never administered in Cuban piggeries (e.g. nalidixic acid, ciprofloxacin, gentamicin, trimethoprim, or cefazolin) has to be interpreted and implemented with caution. From our point of view, administration of new antibiotics in Cuban piggeries currently, will

only attenuate temporally the effects of pathogenic enterobacteria, and will favour the selection of resistant clones. In Cuban piggeries, improvements on the management of pregnant sows, a good attention to the parturition, sufficient intake of colostrum by newborn piglets, vaccination of young pigs and sows against enteropathogens, as well as improvements on housing and biosecurity will decrease diarrhea occurrence, and will decrease the demand of antibiotics.

In 29% and 38% of piggeries, 50% or more sows showed moderate or high levels of serum F4- and F18-specific antibodies, respectively. As vaccination against F4 has not been practiced for the last 15 years in Cuba (Wong *et al.*, 1995), and an F4<sup>+</sup> or F18<sup>+</sup> ETEC/VTEC infection induces F4- or F18-specific antibodies (Verdonck *et al.*, 2002), we can confirm that the F4<sup>+</sup> and F18<sup>+</sup> *E. coli* are highly prevalent as potential enteropathogens in Cuban piggeries. This is in contradiction with Blanco *et al.* (2006) who could not isolate F4<sup>+</sup> *E. coli* in Central Cuba; in addition, we reported multidrug-resistant F4<sup>+</sup> ETEC in the central province Villa Clara.

The high seroprevalence of F4- and F18-specific antibodies in sera of young sows in Cuba (Chapter 4) infer a high genetic susceptibility of pigs raised in Cuban piggeries for colonization by F4<sup>+</sup> and F18<sup>+</sup> *E. coli*.

Susceptibility to colonization by F18<sup>+</sup> *E. coli* is dependent on the glycosylation status influenced by the activity of the *FUT1* gene, which encodes the alpha(1,2)-fucosyltransferase (Meijerink *et al.*, 1997; 2000). Similar to reports by Shi *et al.* (2002) in Yorkshire, Landrace, and Duroc breeds in China and by Coddens *et al.* (2007) in Belgium, pigs carrying the resistant genotype (AA) were not frequently found (13%; Chapter 5). This finding is in agreement with the high prevalence of F18-specific antibodies previously reported, and with the high occurrence (61%) of the F18 fimbrial adhesin among pathogenic *E. coli* isolated from diarrheic piglets in Central Cuba (Blanco *et al.*, 2006). A genetic improvement of pigs raised in Cuba considering the AA resistant genotype to F18 mediated adhesion could be taken as a way to control swine diarrhea or edema disease.

The high seroprevalence of F4-specific antibodies, even though the resistant genotype described by Jørgensen *et al.* (2004) was highly present (66%) in commercial F1 (Yorkshire x Landrace) x Duroc pigs raised in Cuba supports that the polymorphism in *mucin 4* is not a reliable marker for F4ab/ac receptor expression as reported by Rasschaert *et al.* (2007) or that other receptors are involved. Indeed, other F4R candidates have been reported such as an intestinal transferrin (GP74; Grange and Mouricout, 1996), an intestinal glycosphingolipid (Grange *et al.*, 1999), and the porcine aminopeptidase-N (Rasschaert, 2008).

Diarrhea is a multifactorial disease influenced by management and environmental conditions, the interaction with the causative agents and host factors like immunity and genetic susceptibility



(Fairbrother, 2006; Fairbrother and Gyles, 2006). Apart from the stressful weaning, also housing and management aspects could play a role in the appearance of diarrhea outbreaks in Cuban piggeries:

- Big swine farms (> 500 sows) with an open system and all pig categories included are common, making health control difficult.
- A mixture of management failures such as administration of a not well-balanced ration to pregnant and lactating sows which affect their body condition, the presence of sows with an excess number of parturitions and incorrect care of parturition lead to a mixture of problems such as low weight at birth, and also low survival rate of newborn piglets due to mastitis-metritis-agalactia syndrome and or low milk production in sows (Lazo and Gutiérrez, 2011), insufficient intake of colostrum, loss of blood through the umbilical cord, and hypoglycemia.
- Open stables expose pigs to thermal stress by the combination of high environmental temperature and humidity. The lower survival rate (69.7%) of suckling piglets during 2008 was in September and October when, as usual, Cuba was affected by two potent hurricanes. Due to this tropical climate, power supply, water quality, feeding, housing, and sanitation are negatively impacted (Cabrera *et al.*, 2010).
- High density of insect vectors like flies inside installations. Recent surveys performed in Taiwan (Wang *et al.*, 2011) and the Czech Republic (Literak *et al.*, 2009) have implicated flies as vectors for transmission of antibiotic resistant enterobacteria in swine farms.
- Lack of good pre-starting feed (Cabrera *et al.*, 2010) and good feeding management before and after weaning.
- Lack of administration and inadequate control of the herd book.
- Another aspects directly attempt against biosecurity such as low quality of sanitation and disinfection (Cabrera *et al.*, 2010) and the fact that stables are open which make hygiene control difficult. Also, sometimes there is no delimitation of the "dirty" and "clean" areas inside the piggery and breeding of pigs by the personal of the piggery at their home is common which increases the risk of introduction of pathogens to the piggery, mostly when is not ruled the change of boots and clothing before entering the "clean" area. The common interchange of breeding stock pigs among farms and the housing of sows in large groups could also contribute to the spreading of enteropathogens.

## 6.2 Future perspectives

Most of the recent epidemiological data related to swine diarrhea in Cuba do not reach a casual diagnosis or are from studies done in the frame of scientific projects of educational institutions. In order to improve swine health in Cuba, the country has to reinforce the Regional Veterinary Diagnostic Laboratories, which are responsible for the analysis of diarrhea outbreaks, with an improvement of their infrastructure based on the experience and research results of Universities and other Institutions not directly linked with the National Institute of Veterinary Medicine like the National Centre for Animal Health and the Cuban Institute for Swine Research. The improvement of enteropathogen diagnosis at national level, as well as the implementation of efficient prevention and control programs for piglet diarrhea, particularly due to ETEC, might have a positive impact on the performance of young pigs in Cuba.

Due to the application of general laboratory protocols, current reports related with swine diarrhea in Cuba are not specific enough (Ricardo, 2008; Cabrera and Garcia, 2009; Cabrera *et al.*, 2010) so that non-infectious processes like alimentary diarrhea (e.g. non-infectious hyper-lactate or dyspeptic diarrhea; Ushida *et al.*, 2009) mask infectious diarrhea and vice versa. Specifically, we recommend to correctly implement the guidance NRAG (1989) of the Cuban Ministry of Agriculture which rules veterinary diagnostics, microbiological classification and assay procedures in the Veterinary Diagnostic Laboratories. Furthermore, laboratory protocols for the identification of enteropathogens need to be updated. For example, it might be favorable to include virotyping of *E. coli* and toxinotyping of *C. perfringens* by multiplex-PCR during epidemiological analysis and control of swine diarrhea in Cuba.

Pathogenic *E. coli* associated with diarrhea in young pigs appeared resistant to most antibiotics routinely employed in Cuban piggeries. We can infer that other pathogenic bacteria associated with diarrhea or other diseases, and even in other domestic animals or humans, are also resistant to antibiotics currently applied in Cuba. This can result in severe health risks due to outbreaks of multiresistant pathogens. We strongly recommend that Cuban veterinary authorities establish adequate antibiotic resistance surveillance, which is absent, all over the country. The logistic and administrative instances from the National Institute of Veterinary Medicine should consider that instead of cheap and indiscriminately employed antibiotics like tetracycline, a well managed antibiotic strategy together with good management practices of pigs can help to reduce antimicrobial resistance and disease in Cuban piggeries.

The reduction of the morbidity in young pigs due to colibacillosis from 8.3% in 1993 to 0.9% in 1995 in the Camagüey province, Cuba, coincided with the application of the national recombinant vaccine Vacoli, which contained the antigens K88 and K99 (Wong *et al.*, 1995). Nowadays, it might be necessary to re-introduce vaccination against pathogenic *E. coli* in Cuban piggeries. Results of this

thesis suggest that F4 and F18 fimbriae as well as enterotoxins are good candidates to be considered for this preventive strategy. Tiels *et al.* (2008) recommended FaeG adhesin (F4) in combination with FedF adhesin (F18) as a good oral vaccine candidate. Recently, Ruan *et al.* (2011) demonstrated that the fusion antigen FaeG-FedF-LT(192)A2:B elicits antibodies that neutralize LT toxin and inhibit the adherence of F4 and F18 fimbriated *E. coli*.

We think that the presence of the allele A determining F18 resistance at the frequency of 27% in Cuban swines could contribute to future breeding programs aimed at selection of F18-resistant breeding stock pigs in order to prevent diarrhea or edema disease. Selection and breeding of F4-resistant pigs also appear advantageous.

In conclusion, results of this thesis indicate that the economic performance of the Cuban swine industry might be favored by intervention strategies undertaken to improve enteropathogens' diagnosis and to better manage antibiotic use. Prevention programs of diarrhea should be reinforced by vaccination against *E. coli* and other enteropathogens, by breeding pigs resistant to F18<sup>+</sup> *E. coli*, and by improving housing and management in piggeries.

## Summary

Porcine pre- and post-weaning diarrhea are multi-factorial diseases that negatively impact the efficiency of swine production worldwide. Often, swine veterinarians complain about the scarce information on the epidemiology of diarrhea and about limitations for the specific identification of enteropathogens in Cuba. Therefore, insufficient control measures are undertaken which results in an increased incidence of swine diarrhea.

This thesis updates epidemiological knowledge concerning the infectious etiology of diarrhea in young pigs in Cuba through the differential identification of enteropathogens in neonatal, in suckling and in newly-weaned piglets. This epidemiological study identified *Escherichia coli* as a major enteropathogen. Therefore, a major focus was directed on *E. coli* infections in the rest of the thesis. First, the antibiotic resistance and the genetic relatedness of Cuban pathogenic *E. coli* isolates were determined. Then, the seroprevalence of antibodies against F4 and F18 fimbriae of *E. coli* was assessed. Finally, the genetic susceptibility of Cuban pigs for enteric colonization by F4<sup>+</sup> and F18<sup>+</sup> *E. coli* was analyzed.

The **first Chapter** is a review of literature focused on the swine production in the Cuban context and on the epidemiology of the main enteropathogens causing diarrhea in young pigs worldwide. The identification of different enteropathogens is also discussed and the epidemiological term “mixed types” which refers to the combined infectious etiology of diarrhea is introduced.

**Chapter 2** was undertaken to gain insights into the infectious etiology of porcine pre- and post-weaning diarrhea in Villa Clara province, Cuba. Intestinal contents of diarrheic pigs were tested by culture, microscopy, and immunological or DNA-based techniques. At least one enteropathogen was detected in 64.4% and in 42.2% of suckling and weaned pigs, respectively. Enterotoxigenic *E. coli* (ETEC) was significantly the most frequent pathogen, and most virotypes were either STa<sup>+</sup>/STb<sup>+</sup> or F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup>. The overall occurrence of the rest of the pathogens was 10% for transmissible gastroenteritis virus and *Cryptosporidium parvum*, 6.7% for rotavirus A and *Isospora suis*, 5.6% for  $\alpha$ -toxigenic *Clostridium perfringens*, 3.3% for verocytotoxigenic *E. coli* (VTEC), and 2.2% for *Salmonella enterica* subspecies *enterica* serotype Newport. Porcine epidemic diarrhea virus,  $\beta$ -toxigenic *C. perfringens*, *Eimeria* spp., and helminths were not identified. Twelve out of 48 enteropathogen positive piglets (25%) were infected with more than one pathogen and ETEC was present in 10 out of 12 mixed infections. These results demonstrate that several enteropathogens, either alone or as part

of a mixed infection, are associated with porcine pre- and post-weaning diarrhea in the Villa Clara province, Cuba.

Since *E. coli* is the most prevalent enteropathogen in pigs suffering from diarrhea in Cuba, we studied this pathogen in more detail in the next Chapters.

In **Chapter 3** we determined the antibiotic resistance profile and the genetic relatedness of pathogenic *E. coli* isolated from piglets suffering from diarrhea. The highest resistance rates were seen to antibiotics traditionally administered in Cuban piggeries: tetracycline (69%), ampicillin (54%), sulphonamide compounds (50%), and kanamycin (50%); 65% of isolates were multi-drug resistant. The ERIC-PCR revealed a high degree of polymorphism in the *E. coli* DNA sequences and relatedness among F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup> or F18<sup>+</sup>/LT<sup>+</sup>/STb<sup>+</sup> isolates from different piggeries, and among the STb<sup>+</sup>, STa<sup>+</sup>/STb<sup>+</sup>, F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup> or F18<sup>+</sup>/STx2e<sup>+</sup> ones from the same piggery. Genetically diverse as well as genetically related pathogenic *E. coli* highly susceptible to nalidixic acid, ciprofloxacin, gentamicin, amikacin, chloramphenicol, cephalosporins, amoxicilline-clavulanic acid, and trimethoprim are associated with piglet's diarrhea in the Villa Clara province, Cuba. The analysis on genetic diversity by ERIC-PCR demonstrated a clonal relationship among pathogenic *E. coli* carrying the same virulence factors and similar antibiotic resistance. The implementation of these epidemiological research data throughout Cuba can contribute to the surveillance, prevention, and control of swine colibacillosis.

Considering that F4- or F18-specific serum antibodies in Cuban pigs will reflect the presence and spread of F4<sup>+</sup> ETEC or F18<sup>+</sup> ETEC/VTEC, we determined in **Chapter 4** the prevalence of F4- and F18-specific antibodies in sera of 1044 gilts. For the data analysis random-effects models and a mixture model in R (package "mixAK"; Komárek, 2009) were fitted. Low, moderate, and high levels of F4-specific antibodies were found in 67.6%, 26.8%, and 5.6% of gilts, while 66.4% and 33.6% of them showed low and high levels of F18-specific antibodies, respectively. Hereby, we show that F4<sup>+</sup> and F18<sup>+</sup> *E. coli* are highly prevalent as potential enteropathogens in Cuban piggeries.

**Chapter 5** was carried out to firstly know the frequency of genotypes and alleles determining susceptibility or resistance to F18<sup>+</sup> *E. coli* infections. To investigate the susceptibility to F4<sup>+</sup> *E. coli* infections, the *Xba*I polymorphism in the porcine *mucin 4* gene was determined in pigs raised in Villa Clara province, Cuba. The *Xba*I-resistant genotype was frequently detected (0.66) and the susceptible heterozygote or homozygote genotypes were less frequent (0.31 and 0.03, respectively). Since we found that there is a high seroprevalence of F4-specific antibodies in sera of young sows in Cuba, we conclude that the *Xba*I polymorphism in *mucin 4* cannot be used as a reliable marker to determine genetic resistance to F4<sup>+</sup> *E. coli* infections.

Pigs carrying the F18-resistant genotype were not frequently found (0.13), while the heterozygous or homozygous susceptible genotypes occurred most frequently (0.27 and 0.60, respectively), coinciding with previous reports of a high prevalence of F18<sup>+</sup> *E. coli* and F18-specific antibodies in the Cuban swine herd. At least for F18<sup>+</sup> *E. coli* control, a genetic improvement of pigs considering the resistant genotype could be taken as a way to control swine diarrhea or edematous disease in Cuba. This strategy is not recommended for F4 since the *Xba*I polymorphism in the *mucin 4* gene does not appear to be a good marker for resistance to F4<sup>+</sup> *E. coli* infections.

Finally, the general discussion and future perspectives are presented in **Chapter 6**. A better production efficiency of the Cuban Company for pork production might be achieved by the improvement of surveillance, prevention, and control programs of infectious diarrhea in young pigs, particularly colibacillosis. Cuban veterinary authorities should acknowledge the importance of identification of enteropathogens and of surveillance of antibiotic resistance by the Veterinary Diagnostic Laboratories. In the short-term, vaccination against *E. coli*, the introduction of efficient antibiotics and improvement of management practices appear advantageous to control porcine pre- and post-weaning diarrhea in Cuba. In the long-term, breeding of F4- and F18-resistant pigs as well as investments in facilities will be necessary.



## Samenvatting

Diarree bij neonatale en pasgespeende biggen is een multifactoriële ziekte die economische schade berokkent aan varkenshouders wereldwijd. Voor Cubaanse dierenartsen is er weinig informatie beschikbaar over de epidemiologie van diarree bij jonge biggen en is de kennis over de identiteit van de enteropathogenen gelimiteerd. Daardoor kunnen onvoldoende controle maatregelen genomen worden wat resulteert in een verhoogd voorkomen van varkensdiarree.

Deze doctoraatsthesis geeft recente inzichten in de etiologie van diarree bij jonge biggen in Cuba. Hiertoe werd identificatie van de verschillende enteropathogenen aanwezig in faeces van biggen met diarree uitgevoerd. Omdat *Escherichia coli* de meest voorkomende pathogeen bleek, werd van deze *E. coli* stammen ook het antibioticumresistentieprofiel en de genetische verwantschap bepaald. Bovendien werd de seroprevalentie van specifieke antistoffen tegen de fimbriae van F4<sup>+</sup> *E. coli* en F18<sup>+</sup> *E. coli* -die gekend zijn als de meest voorkomende *E. coli* betrokken bij varkensdiarree- onderzocht en werd de genetische gevoeligheid van biggen voor F4<sup>+</sup> en F18<sup>+</sup> *E. coli* infecties bepaald.

In het **eerste Hoofdstuk** wordt een overzicht gegeven van recente literatuur omtrent de varkensproductie in Cuba en omtrent de epidemiologie van de belangrijkste enteropathogenen die diarree veroorzaken bij jonge biggen wereldwijd. Hier wordt ook de identificatie van de verschillende enteropathogenen besproken en wordt de epidemiologische term 'Mixed types' geïntroduceerd, die verwijst naar de aanwezigheid van verscheidene pathogenen in faeces van biggen die door diarree getroffen zijn.

Het onderzoek in **Hoofdstuk 2** werd ondernomen om inzicht te verwerven in de infectieuze oorzaken van diarree bij neonatale en pasgespeende varkens in de Villa Clara provincie in Cuba. Darminhoud van biggen met diarree werden getest door cultivatie, microscopie en immunologische of DNA-gebaseerde technieken. Ten minste 1 enteropathogeen werd gedetecteerd in respectievelijk 64.4% en 42.2% van de zuigende en gespeende biggen. Enterotoxigene *E. coli* (ETEC) was significant de meest voorkomende pathogeen en de meeste virotypes waren ofwel STa<sup>+</sup>/STb<sup>+</sup> of F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup>. Het algemeen voorkomen van de overige geteste pathogenen was 10% voor transmissibel gastroenteritis virus en *Cryptosporidium parvum*, 6.7% voor rotavirus A en *Isospora suis*, 5.6% voor α-toxigene *Clostridium perfringens*, 3.3% voor verotoxigene *E. coli* (VTEC) en 2.2% voor *Salmonella enterica* subspecies *enterica* serovar Newport. Het porcine epizoötische diarree virus, β-toxigene *C. perfringens*, *Eimeria* spp. en helminthen werden niet teruggevonden. Twaalf van de 48 enteropathogeen-positieve varkens (25%) werden geïnfecteerd met meer dan één



pathogeen en ETEC was aanwezig bij 10 van de 12 gemengde infecties. Deze resultaten tonen aan dat verschillende pathogenen, ofwel individueel ofwel als onderdeel van een gemengde infectie, geassocieerd kunnen worden met varkensdiarree in neonatale en pasgespeende biggen in de Villa Clara provincie in Cuba.

Gezien *E. coli* de meest voorkomende enteropathogeen is bij biggen met diarree in Cuba, werd deze pathogeen meer gedetailleerd bestudeerd in de volgende hoofdstukken.

In **Hoofdstuk 3** werd het antibioticumresistentieprofiel en de genetische verwantschap tussen pathogene *E. coli* isolaten van biggen die lijden aan diarree onderzocht. De hoogste graad van antibioticumresistentie werd gevonden bij antibiotica die traditioneel toegediend worden bij Cubaanse varkensbedrijven, namelijk tetracycline (69%), ampicilline (54%), sulfonamides (50%) en kanamycine (50%). Vijfenzestig procent van de isolaten waren resistent tegen meerdere antibiotica. De ERIC-PCR toonde aan dat er een hoge graad van polymorfismen is in de *E. coli* DNA sequenties en dat er een hoge graad van verwantschap is tussen F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup> of F18<sup>+</sup>/LT<sup>+</sup>/STb<sup>+</sup> isolaten van verschillende varkenshouderijen en tussen de STb<sup>+</sup>, STa<sup>+</sup>/STb<sup>+</sup>, F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup> of F18<sup>+</sup>/STx2e<sup>+</sup> isolaten. Zowel genetisch diverse als genetisch verwante pathogene *E. coli* die sterk gevoelig zijn voor nalidixinezuur, ciprofloxacine, gentamycine, amikacine, chloramphenicol, cephalosporines, amoxicilline-clavulaanzuur en trimethoprim zijn geassocieerd met diarree van biggen in de Villa Clara provincie in Cuba. De analyse van de genetische diversiteit met de ERIC-PCR toonde een klonale verwantschap aan tussen pathogene *E. coli* stammen die dezelfde virulentiefactoren dragen en hetzelfde antibioticumresistentieprofiel hebben. De implementatie van deze epidemiologische onderzoeksgegevens in heel Cuba kan bijdragen aan het epidemiologische surveillance, preventie en controle van varken colibacillose.

In acht genomen dat de aanwezigheid van F4- en F18-specifieke antistoffen bij Cubaanse varkens een weerspiegeling is van de aanwezigheid en verspreiding van F4<sup>+</sup> ETEC of F18<sup>+</sup> ETEC/VTEC, werd in **Hoofdstuk 4** het voorkomen van F4- en F18-specifieke antistoffen in sera van 1044 gelten onderzocht. Data analyse gebeurde aan de hand van een 'random effect model' en een 'mixture model' in R (package "mixAK"; Komárek, 2009). Lage, gemiddelde en hoge aantallen F4-specifieke antistoffen werden gevonden in respectievelijk 67.6%, 26.8% en 5.6% van de gelten, terwijl lage en hoge aantallen F18-specifieke antistoffen respectievelijk werden teruggevonden in 66.4% en 33.6% van de onderzochte gelten. Dit toont aan dat F4<sup>+</sup> *E. coli* en F18<sup>+</sup> *E. coli* een zeer veel voorkomende enteropathogeen is in Cubaanse varkensbedrijven.

In **Hoofdstuk 5** werd onderzocht wat het voorkomen is van de genotypes en de allelen die bepalen of biggen gevoelig of resistent zijn voor F18<sup>+</sup> *E. coli* infecties. Om de gevoeligheid voor F4<sup>+</sup> *E. coli* infecties te bepalen, werd het *Xba*I polymorfisme in het *mucine 4* gen van het varken onderzocht bij varkens gekweekt in de Villa Clara provincie in Cuba. Het *Xba*I resistente genotype werd het meest teruggevonden (0.66), terwijl de heterozygoot of homozygoot gevoelige genotypes minder frequent voorkwamen (respectievelijk 0.31 en 0.03). Gezien we een hoge seroprevalentie van F4-specifieke antistoffen in sera van jonge zeugen gevonden hebben in Cuba, kunnen we besluiten dat het *Xba*I polymorfisme in het *mucine 4* gen niet kan gebruikt worden als betrouwbare merker om genetische resistentie tegen F4<sup>+</sup> *E. coli* op te sporen.

Om de gevoeligheid voor F18<sup>+</sup> *E. coli* infecties te achterhalen, werd het *Cfo*I polymorfisme in het *FUT1* gen onderzocht. Varkens met het F18-resistente genotype werden niet frequent teruggevonden (0.13), terwijl de heterozygoot of homozygoot gevoelige genotypes het meest frequent voorkwamen (respectievelijk 0.27 en 0.60). Het hoge aantal F18<sup>+</sup> *E. coli* gevoelige biggen correleert met de hoge prevalentie van F18<sup>+</sup> *E. coli* en F18-specifieke antistoffen in de Cubaanse varkenspopulatie. Genetische selectie van biggen met het F18<sup>+</sup> *E. coli* resistente genotype zou dus in de toekomst een maatregel kunnen zijn om diarree en slingerziekte bij varkens in Cuba terug te dringen. Deze strategie wordt nog niet aanbevolen voor F4<sup>+</sup> *E. coli* gezien het *Xba*I polymorfisme in het *mucine 4* gen geen betrouwbare merker is om resistentie tegen F4<sup>+</sup> *E. coli* terug te vinden.

Tot slot worden de algemene discussie en de toekomstperspectieven weergegeven in **Hoofdstuk 6**. Er wordt voorgesteld dat een hogere productie-efficiëntie van het Cubaans Bedrijf voor varkensproductie kan bereikt worden door de verbetering van bewakings-, preventie- en controleprogramma's van infectieuze diarree, en voornamelijk van collibacillose bij jonge biggen. De Cubaanse overheid zou het belang moeten erkennen van de identificatie van de enteropathogenen en van het opsporen van antibioticumresistenties door de diagnostische laboratoria. Op korte termijn zou vaccinatie tegen *E. coli*, het invoeren van efficiënte antibiotica en de verbetering van management maatregelen zeer gunstig kunnen zijn om neonatale en speendiarree bij jonge biggen in Cuba in te perken. Op lange termijn zou selectie voor F4- en F18-resistente biggen en investeringen in de faciliteiten van de Cubaanse varkensbedrijven sterke voordelen kunnen opleveren om F4<sup>+</sup> en F18<sup>+</sup> *E. coli* infecties te bestrijden.



## Resumen

La diarrea pre- y post-destete de los cerdos son enfermedades multifactoriales que afectan negativamente la eficiencia de la producción en granjas porcinas a nivel mundial. A menudo los veterinarios del sector porcino de Cuba se quejan de la escasa información sobre la epidemiología de la diarrea y las limitaciones para la identificación específica de enteropatógenos. Por lo tanto, se llevan a cabo medidas de control insuficientes que resultan en un aumento de la incidencia de la diarrea porcina.

La presente tesis realiza una actualización del conocimiento epidemiológico sobre la etiología infecciosa de la diarrea en cerdos jóvenes de Cuba a través de la identificación diferencial de enteropatógenos en cerdos recién nacidos, lactantes, y recién destetados. Este estudio epidemiológico identificó a *Escherichia coli* como un enteropatógeno común por lo que la tesis se enfoca más en las infecciones por *E. coli*. En primer lugar se determinó la resistencia a los antibióticos y la relación genética de cepas cubanas de *E. coli* patógenas. Posteriormente se evaluó la seroprevalencia de anticuerpos contra las fimbrias F4 y F18 de *E. coli*. Por último se analizó la susceptibilidad genética de los cerdos de Cuba para la colonización entérica por *E. coli* F4<sup>+</sup> y F18<sup>+</sup>.

El **primer Capítulo** es una revisión bibliográfica dedicada a la producción porcina en el contexto cubano y a la epidemiología de los principales enteropatógenos causantes de diarrea en cerdos jóvenes en todo el mundo. También se discutió sobre la identificación diferencial de enteropatógenos y se introdujo el término epidemiológico "tipos mixtos" que se refiere a la etiología infecciosa combinada que la diarrea porcina puede presentar.

El **Capítulo 2** se llevó a cabo para obtener información sobre la etiología infecciosa de la diarrea porcina pre- y post-destete en la provincia de Villa Clara, Cuba. Fueron analizados contenidos intestinales de cerdos diarreicos por medio de cultivo, microscopía, y técnicas inmunológicas o basadas en el ADN. Por lo menos un enteropatógeno se detectó en el 64,4% y en el 42,2% de los lechones lactantes y destetados, respectivamente. *E. coli* enterotoxigénica (ETEC) fue significativamente el patógeno más frecuente, y la mayoría de los virotipos fueron STa<sup>+</sup>/STb<sup>+</sup> o F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup>. La ocurrencia global del resto de los patógenos fue del 10% para el virus de la gastroenteritis transmisible y *Cryptosporidium parvum*, 6,7% para rotavirus A e *Isospora suis*, 5,6% para *Clostridium perfringens*  $\alpha$ -toxigénico, 3,3% para *E. coli* verotoxigénica (VTEC), y 2,2% para *Salmonella enterica* subespecie *enterica* serovar Newport. El virus de la diarrea epidémica porcina, *C. perfringens*  $\beta$ -toxigénico, *Eimeria* spp., y helmintos no fueron identificados. Doce de los 48 lechones

positivos a enteropatógenos (25%) estaban infectados con más de un patógeno y ETEC estuvo presente en 10 de las 12 infecciones mixtas. Estos resultados demuestran que varios enteropatógenos, solos o como parte de una infección mixta, están asociados con la diarrea porcina pre- y post-destete en la provincia de Villa Clara, Cuba.

Dado que *E. coli* fue el enteropatógeno más frecuente de los cerdos diarreicos en Cuba, este patógeno se estudió más detalladamente en los capítulos siguientes.

En el **Capítulo 3** se determinó el perfil de resistencia a los antibióticos y la relación genética de *E. coli* patógenas aisladas de lechones que sufrían diarrea. Las mayores tasas de resistencia se mostraron ante antibióticos tradicionalmente administrados en granjas porcinas cubanas: tetraciclina (69%), ampicilina (54%), compuestos de la sulfonamida (50%), y kanamicina (50%); además el 65% de los aislados fueron resistentes a múltiples antibióticos. El ERIC-PCR reveló un alto grado de polimorfismo en las secuencias de ADN de *E. coli* así como relación entre los aislados F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup> o F18<sup>+</sup>/LT<sup>+</sup>/STb<sup>+</sup> detectados en granjas porcinas diferentes, y entre aislados STb<sup>+</sup>, STa<sup>+</sup>/STb<sup>+</sup>, F4<sup>+</sup>/STa<sup>+</sup>/STb<sup>+</sup> o F18<sup>+</sup>/Stx2e<sup>+</sup> procedentes de una misma cochiquera. *E. coli* patogénicas, genéticamente diversas o genéticamente relacionadas, así como altamente susceptibles al ácido nalidíxico, ciprofloxacina, gentamicina, amikacina, cloranfenicol, cefalosporinas, amoxicilina-ácido clavulánico y trimetoprima están asociadas con la diarrea de los lechones en la provincia de Villa Clara, Cuba. El análisis de la diversidad genética por medio de ERIC-PCR demostró relación clonal entre *E. coli* patógenas que portaban los mismos factores de virulencia y similar perfil de resistencia a los antibióticos. La implementación de los resultados de esta investigación epidemiológica en toda Cuba puede contribuir a la vigilancia, prevención y control de la colibacilosis porcina.

Teniendo en cuenta que los anticuerpos séricos específicos para F4 o F18 en los cerdos cubanos podrían reflejar la presencia y propagación de ETEC F4<sup>+</sup> o ETEC/VTEC F18<sup>+</sup>, en el **Capítulo 4** se determinó la prevalencia de anticuerpos específicos contra F4 y F18 en el suero de 1044 cerdas jóvenes. Para el análisis de datos se emplearon modelos de efectos aleatorios así como modelos de mezcla en R (paquete "mixAK"; Komárek, 2009). Niveles bajos, moderados y altos de anticuerpos específicos contra F4 se encontraron en el 67,6%, 26,8% y 5,6% de las cerdas, mientras que el 66,4% y 33,6% presentaron niveles bajos y altos de anticuerpos específicos contra F18, respectivamente. En este estudio se ha mostrado que *E. coli* F4<sup>+</sup> y F18<sup>+</sup> son muy frecuentes como enteropatógenos potenciales en cochiqueras cubanas.

El **Capítulo 5** se llevó a cabo para conocer en primer lugar la frecuencia de los genotipos y los alelos que determinan la susceptibilidad o resistencia a las infecciones por *E. coli* F18<sup>+</sup>. Para investigar la susceptibilidad a infecciones por *E. coli* F4<sup>+</sup> se determinó el polimorfismo *Xba*I del gen porcino *mucin 4* en cerdos criados en la provincia Villa Clara, Cuba. El genotipo resistente a la digestión por *Xba*I fue detectado frecuentemente (0,66), y los genotipos susceptibles heterocigoto u homocigoto fueron menos frecuentes (0,31 y 0,03, respectivamente). Dado que hemos encontrado que hay una alta seroprevalencia de anticuerpos específicos contra F4 en suero de cerdas jóvenes en Cuba, llegamos a la conclusión de que el polimorfismo *Xba*I en el gen *mucin 4* no puede ser utilizado como un marcador fiable para determinar la resistencia genética a infecciones por *E. coli* F4<sup>+</sup>. Los cerdos portadores del genotipo resistente a F18 no fueron encontrados con frecuencia (0,13), mientras que los genotipos susceptibles homocigoto o heterocigoto ocurrieron con mayor frecuencia (0,27 y 0,60, respectivamente), coincidiendo con informes anteriores de una alta prevalencia de *E. coli* F18<sup>+</sup> y de anticuerpos específicos contra F18 en el rebaño porcino de Cuba. Por lo menos para el control de *E. coli* F18<sup>+</sup> se podría considerar un mejoramiento genético de los cerdos teniendo en cuenta el genotipo resistente como una manera de controlar la diarrea porcina o la enfermedad edemática en Cuba. Esta estrategia no se recomienda para F4 debido a que el polimorfismo *Xba*I en el gen *mucin 4* no parece ser un buen marcador de resistencia a infecciones por *E. coli* F4<sup>+</sup>.

Por último, la discusión general y las perspectivas futuras se presentan en el **Capítulo 6**. Una mejor eficiencia productiva de la Compañía Cubana de producción de carne de cerdo podría lograrse mediante la mejora de los programas de vigilancia, prevención y control de la diarrea infecciosa en los cerdos jóvenes, en particular colibacilosis. Las autoridades veterinarias cubanas deberían incentivar la identificación de enteropatógenos y la vigilancia de resistencia a los antibióticos por los laboratorios de diagnóstico veterinario. En el corto plazo, la vacunación contra *E. coli*, la introducción de antibióticos eficaces y la mejora de las prácticas de gestión parecen ser ventajosos para controlar la diarrea porcina pre- y post-destete en Cuba. En el largo plazo, la cría de cerdos resistentes a F18 o F4, así como inversiones en instalaciones, será necesario.



## References

## A

- Abe M, Ito N, Morikawa S, Takasu M, Murase T, Kawashima T, Kawai Y, Kohara J, Sugiyama M, 2009.** Molecular epidemiology of rotaviruses among healthy calves in Japan: isolation of a novel bovine rotavirus bearing new P and G genotypes, *Virus Research*, **144**:250-7.
- Adesiyun AA, Kaminjolo JS, Ngeleka M, Mutani A, Borde G, Harewood W, Harper W, 2001.** A longitudinal study on enteropathogenic infections of livestock in Trinidad, *Revista da Sociedade Brasileira de Medicina Tropical*, **34**:29-35.
- Aitkin M, Liu CC, Chadwick T, 2009.** Bayesian model comparison and model averaging for small-area estimation, *Annals of Applied Statistics*, **3**:199-221.
- Akwar HT, Poppe C, Wilson J, Reid-Smith RJ, Dyck M, Waddington J, Shang D, McEwen SA, 2008.** Prevalence and patterns of antimicrobial resistance of fecal *Escherichia coli* among pigs on 47 farrow-to-finish farms with different in-feed medication policies in Ontario and British Columbia, *Canadian Journal of Veterinary Research*, **72**:195-201.
- Aliaga-Leyton A, Webster E, Friendship R, Dewey C, Vilaça K, Peregrine AS, 2011.** An observational study on the prevalence and impact of *Isospora suis* in suckling piglets in southwestern Ontario, and risk factors for shedding oocysts, *The Canadian Veterinary Journal*, **52**:184-8.
- Atii DJ, Ojeh CK, Durojaiye OA, 1990.** Detection of rotavirus antigen in diarrhoeic and non-diarrhoeic piglets in Nigeria, *Revue d'élevage et de médecine vétérinaire des pays tropicaux*, **42**:494-6.

## B

- Baba E, Gaafar SM, 1985.** Interfering effect of *Isospora suis* infection on *Salmonella typhimurium* infection in swine, *Veterinary Parasitology*, **17**:271-8.
- Baker AA, Davis E, Rehberger T, Rosener D, 2010.** Prevalence and diversity of toxigenic *Clostridium perfringens* and *Clostridium difficile* among swine herds in the midwest, *Applied and Environmental Microbiology*, **76**:2961-7.
- Baker DR, Billey LO, Francis DH, 1997.** Distribution of K88 *Escherichia coli*-adhesive and nonadhesive phenotypes among pigs of four breeds, *Veterinary Microbiology*, **54**:123-32.
- Ball JM, Tian P, Zeng CQ, Morris AP, Estes MK, 1996.** Age-dependent diarrhea induced by a rotaviral nonstructural glycoprotein, *Science*, **272**:101-4.
- Bao WB, Wu SL, Musa HH, Zhu GQ, Chen GH, 2008.** Genetic variation at the alpha-1-fucosyltransferase (FUT1) gene in Asian wild boar and Chinese and Western commercial pig breeds, *Journal of Animal Breeding and Genetics*, **125**:427-30.
- Barreiros MA, Alfieri AA, Alfieri AF, Médici KC, Leite JP, 2003.** An outbreak of diarrhoea in one-week-old piglets caused by group A rotavirus genotypes P[7],G3 and P[7],G5, *Veterinary Research Communications*, **27**:505-12.



- Barrera M, Díaz de Arce H, Acevedo AM, Cuello S, Rodríguez E, Vega A, Urquiaga R, Frías MT, 2005.** Transmissible gastroenteritis in Cuba: experimental reproduction of the disease and molecular characterization of the virus, *Spanish Journal of Agricultural Research*, **3:267-74**.
- Berberov EM, Zhou Y, Francis DH, Scott MA, Kachman SD, Moxley RA, 2004.** Relative importance of heat-labile enterotoxin in the causation of severe diarrheal disease in the gnotobiotic piglet model by a strain of enterotoxigenic *Escherichia coli* that produces multiple enterotoxins, *Infection and Immunity*, **72:3914-24**.
- Bert F, Johnson JR, Ouattara B, Leflon-Guibout V, Johnston B, Marcon E, Valla D, Moreau R, Nicolas-Chanoine MH, 2010.** Genetic diversity and virulence profile of *Escherichia coli* isolates causing spontaneous bacterial peritonitis and bacteremia in patients with cirrhosis, *Journal of Clinical Microbiology*, **48:2709-14**.
- Bertschinger HU, Stamm M, Vogeli P, 1993.** Inheritance of resistance to oedema disease in the pig: experiments with an *Escherichia coli* strain expressing fimbriae 107, *Veterinary Microbiology*, **35:79-89**.
- Bibbal D, Dupouy V, Prère MF, Toutain PL, Bousquet-Mélou A, 2009.** Relatedness of *Escherichia coli* Strains with Different Susceptibility Phenotypes Isolated from Swine Feces during Ampicillin Treatment, *Applied and Environmental Microbiology*, **75:2999-3006**.
- Bijlsma IG, de Nijs A, Frik JF, 1985.** The prevalence of different porcine phenotypes in the Netherlands concerning adherence of K88-positive *Escherichia coli* to intestinal epithelium (Abstract), *The Veterinary Quarterly*, **7:246-8**.
- Blake DP, Humphry RW, Scott KP, Hillman K, Fenlon DR, Low JC, 2003.** Influence of tetracycline exposure on tetracycline resistance and the carriage of tetracycline resistance genes within commensal *Escherichia coli* populations, *Journal of Applied Microbiology*, **94:1087-97**.
- Blanco M, Lazo L, Blanco JE, Dahbi G, Mora A, López C, González EA, Blanco J, 2006.** Serotypes, virulence genes, and PFGE patterns of enteropathogenic *Escherichia coli* isolated from Cuban pigs with diarrhea, *International Microbiology*, **9:53-60**.
- Boerlin P, Travis R, Gyles CL, Reid-Smith R, Lim H, Janecko N, Nicholson V, McEwen SA, Friendship R, Archambault M, 2005.** Antimicrobial resistance and virulence genes of *Escherichia coli* isolates from swine in Ontario, *Applied Environmental Microbiology*, **71:6753-61**.
- Brown IH, Paton DJ, 1991.** Serological studies of transmissible gastroenteritis in Great Britain, using a competitive ELISA, *The Veterinary Record*, **128:500-3**.
- Bruggeman M, Decostere A, Pasmans F, Haesebrouck F, Butaye P, 2008.** Sense and nonsense of determining the presence of *E. coli* in feces from diarrheic dogs, *Vlaams Diergeneeskundig Tijdschrift*, **78:177-81**.
- Bueschel DM, Jost BH, Billington SJ, Trinh HT, Songer JG, 2003.** Prevalence of cpb2, encoding beta2 toxin, in *Clostridium perfringens* field isolates: correlation of genotype with phenotype, *Veterinary Microbiology*, **94:121-9**.

**Bunting M, Lorant DE, Bryant AE, Zimmerman GA, McIntyre TM, Stevens DL, Prescott SM, 1997.** Alpha toxin from *Clostridium perfringens* induces proinflammatory changes in endothelial cells, *The Journal of Clinical Investigation*, **100**:565-74.

## C

**Cabrera LF, García MG, 1985.** Intestinal cryptosporidiosis of diarrhoeic piglets in our environment (Abstract), *Revista Cubana de Ciencias Veterinarias*, **16**:259-64.

**Cabrera Y, García A, 2009.** Incidencia de los trastornos gastrointestinales en la mortalidad en el sector especializado del GRUPOR en el año 2008, *Boletín Técnico Porcino*, **(11)**:11-2.

**Cabrera Y, García A, Martínez V, Melián Y, Naranjo R, 2010.** Incidencia de los trastornos gastrointestinales en la masa porcina cubana, *Proceedings of the IV Seminario Internacional Porcicultura Tropical, La Habana, Cuba*, pp. 374-8.

**Callebaut P, Correa I, Pensaert M, Jiménez G, Enjuanes L, 1988.** Antigenic differentiation between transmissible gastroenteritis virus of swine and a related porcine respiratory coronavirus, *Journal of General Virology*, **69**:1725-30.

**Cesaris L, Gillespie BE, Srinivasan V, Almeida RA, Zeconi A, Oliver SP, 2007.** Discriminating between strains of *Escherichia coli* using pulsed-field gel electrophoresis and BOX-PCR, *Foodborne Pathogens and Disease*, **4**:473-80.

**Chae C, Kim O, Choi C, Min K, Cho WS, Kim J, Tai JH, 2000.** Prevalence of porcine epidemic diarrhea virus and transmissible gastroenteritis virus infection in Korean pigs, *The Veterinary Record*, **147**:606-8.

**Chae C, Kwon D, Kim O, Min K, Cheon DS, Choi C, Kim B, Suh J, 1998.** Diarrhea in nursing piglets associated with coccidiosis: prevalence, microscopic lesions and coexisting microorganisms, *The Veterinary Record*, **143**:417-20.

**Chen F, Huang K, 2007.** Prevalence and phylogenetic analysis of *Cryptosporidium* in pigs in eastern China, *Zoonoses and Public Health*, **54**:393-400.

**Chen X, Gao S, Jiao X, Liu XF, 2004.** Prevalence of serogroups and virulence factors of *Escherichia coli* strains isolated from pigs with postweaning diarrhea in eastern China, *Veterinary Microbiology*, **103**:13-20.

**Cheng D, Sun H, Xu J, Gao S, 2005.** Prevalence of fimbrial colonization factors F18ab and F18ac in *Escherichia coli* isolates from weaned piglets with edema and/or diarrhea in China, *Veterinary Microbiology*, **110**:35-9.

**Cheng D, Sun H, Xu J, Gao S, 2006.** PCR detection of virulence factor genes in *Escherichia coli* isolates from weaned piglets with edema disease and/or diarrhea in China, *Veterinary Microbiology*, **115**:320-8.

**Chizhikov V, Wagner M, Ivshina A, Hoshino Y, Kapikian AZ, Chumakov K, 2002.** Detection and Genotyping of Human Group A Rotaviruses by Oligonucleotide Microarray Hybridization, *Journal of Clinical Microbiology*, **40**:2398-2407.

- Chiu CH, Su LH, Chu C, 2004.** *Salmonella enterica* serotype Choleraesuis: epidemiology, pathogenesis, clinical disease, and treatment, *Clinical Microbiology Reviews*, **17**:311-22.
- Choi BY, Sohn YS, Choi C, Chae C, 2003.** Lectin histochemistry for glycoconjugates in the small intestines of piglets naturally infected with *Isospora suis*, *The Journal of Veterinary Medical Science*, **65**:389-92.
- Choi C, Chae C, 1999.** Genotypic prevalence of F4 variants (ab, ac, and ad) in *Escherichia coli* isolated from diarrheic piglets in Korea, *Veterinary Microbiology*, **67**:307-10.
- Coddens A, Diswall M, Angström J, Breimer ME, Goddeeris B, Cox E, Teneberg S, 2009.** Recognition of blood group ABH type 1 determinants by the FedF adhesin of F18-fimbriated *Escherichia coli*, *The Journal of Biological Chemistry*, **284**:9713-26.
- Coddens A, Valis E, Benktander J, Ångström J, Breimer ME, Cox E, Teneberg S, 2011.** Erythrocyte and porcine intestinal glycosphingolipids recognized by F4 fimbriae of enterotoxigenic *Escherichia coli*, *PLoS One*, **6**:e23309.
- Coddens A, Verdonck F, Tiels P, Rasschaert K, Goddeeris BM, Cox E, 2007.** The age-dependent expression of the F18<sup>+</sup> *E. coli* receptor on porcine gut epithelial cells is positively correlated with the presence of histo-blood group antigens, *Veterinary Microbiology*, **122**:332-41.
- Collins JE, Bergeland ME, Bouley D, Ducommun AL, Francis DH, Yeske P, 1989.** Diarrhoea associated with *Clostridium perfringens* type A enterotoxin in neonatal pigs, *Journal of Veterinary Diagnostic Investigation*, **1**:351-3.
- Correa I, Gebauer F, Bullido MJ, Suñé C, Baay MF, Zwaagstra KA, Posthumus WP, Lenstra JA, Enjuanes L, 1990.** Localization of antigenic sites of the E2 glycoprotein of transmissible gastroenteritis coronavirus, *The Journal of General Virology*, **71**:271-9.
- Costinar L, Pascu C, Herman V, Sorescu D, Surpat A, 2010.** Epidemiological studies of enteropathogens (Rotavirus, *Escherichia coli*, *Coccidia*, *Balantidium coli*) in suckling piglets with diarrhea. In: E grosse Beilage and T Blaha (eds.), *Proceedings of the 2<sup>nd</sup> European Symposium on Porcine Health Management Congress, Hannover, Germany*, pp. 133.
- Cox E, Houvenaghel A, 1987.** In vitro adhesion of K88ab-, K88ac- and K88ad-positive *Escherichia coli* to intestinal villi, to buccal cells and to erythrocytes of weaned piglets, *Veterinary Microbiology*, **15**:201-7.
- Cox E, Pensaert MB, Callebaut P, 1993.** Intestinal protection against challenge with transmissible gastroenteritis virus of pigs immune after infection with the porcine respiratory coronavirus, *Vaccine*, **11**:267-72.
- Cruz EC, Salvarani FM, Alberafo IS, Silva RO, Lobato FF, Stancioli EF, Guedes RM, 2010.** Frequency of enteropathogens in diarrheic and non-diarrheic neonatal piglets. In: S D'Allaire and R Friendship (eds), *Proceedings of the 21<sup>st</sup> International Pig Veterinary Society Congress, Vancouver, Canada*, pp. 757.
- Cubero MJ, León L, Contreras A, Astorga R, Lanza I, Garcia A, 1993.** Transmissible gastroenteritis in pigs in south east Spain: prevalence and factors associated with infection, *The Veterinary Record*, **132**:238-41.

**Czanderlova L, Zizlavsky M, Kellnerova D, Gambota M, Odehnalova S, Odehnal J, 2010.** Prevalence of rotavirus and coronavirus in piglet production herds in the Czech Republic. In: S D'Allaire and R Friendship (eds), *Proceedings of the 21<sup>st</sup> International Pig Veterinary Society Congress, Vancouver, Canada*, pp. 815.

## D

**Dalla-Costa L, Irino K, Rodrigues J, Rivera ING, Trabulsi LR, 1998.** Characterization of diarrheagenic *Escherichia coli* clones by ribotyping and ERIC-PCR. *Journal of Medical Microbiology*, **47**:227-34.

**Damriyasa IM, Bauer C, 2006.** Prevalence and age-dependent occurrence of intestinal protozoan infections in suckling piglets, *Berliner und Munchener Tierarztliche Wochenschrift*, **119**:287-90.

**Das A, Mazumder Y, Dutta BK, Kumar A, Selvi S, 2009.** Diagnosis of acute diarrhea in pigs and piglets in Meghalaya, India, *Malaysian Journal of Microbiology*, **5**:38-44.

**Dauguschies A, Bialek R, Joachim A, Mundt HC, 2001.** Autofluorescence microscopy for the detection of nematode eggs and protozoa, in particular *Isospora suis*, in swine faeces, *Parasitology Research*, **87**:409-12.

**de la Fé Rodríguez PY, Brito E, Aguiar J, Rodríguez L, Hernández JA, 2007.** Estudio de la prevalencia de las endoparasitosis que afectan a los cerdos en el territorio de Cuba, *Revista Electrónica de Veterinaria REDVET*, **8**(4).

**de la Fé Rodríguez PY, Coddens A, Del Fava E, Cortiñas Abrahantes J, Shkedy Z, Maroto Martin LO, Cruz Muñoz E, Duchateau L, Cox E, Goddeeris BM, 2011.** High prevalence of F4<sup>+</sup> and F18<sup>+</sup> *Escherichia coli* in Cuban piggeries as determined by serological survey, *Tropical Animal Health and Production*, **43**:937-46.

**de la Fé Rodríguez PY, Ndemi Kiiru J, Maroto Martin LO, Cruz Muñoz E, Butaye P, Cox E, Goddeeris BM, 2012.** Characterization and clonal grouping of pathogenic *Escherichia coli* isolated from intestinal contents of diarrheic piglets in Villa Clara province, Cuba, according to their antibiotic resistance and ERIC-PCR profiles, *Veterinary Microbiology*, **154**:425-8.

**Delmas B, Gelfi J, Laude H, 1986.** Antigenic structure of transmissible gastroenteritis virus. II. Domains in the peplomer glycoprotein, *The Journal of General Virology*, **67**:1405-18.

**Delmas B, Gelfi J, L'Haridon R, Vogel LK, Sjöström H, Norén O, Laude H, 1992.** Amino peptidase N is a major receptor for the entero-pathogenic coronavirus TGEV, *Nature*, **357**:417-20.

**Dewulf J, Catry B, Timmerman T, Opsomer G, de Kruif A, Maes D, 2007.** Tetracycline-resistance in lactose-positive enteric coliforms originating from Belgian fattening pigs: degree of resistance, multiple resistance and risk factors, *Preventive Veterinary Medicine*, **78**:339-51.

**Dias RC, Marangoni DV, Smith SP, Alves EM, Pellegrino FL, Riley LW, Moreira BM, 2009.** Clonal composition of *Escherichia coli* causing community-acquired urinary tract infections in the State of Rio de Janeiro, Brazil, *Microbial Drug Resistance*, **15**:303-8.

- Djurickovic S, Thorsen J, Duncan JR, Roe CK, 1969.** Transmissible gastroenteritis of swine in Ontario, *Canadian Journal of Comparative Medicine*, **33:59-63**.
- Do TN, Cu PH, Nguyen HX, Au TX, Vu QN, Driesen SJ, Townsend KM, Chin JJ, Trott DJ, 2006.** Pathotypes and serogroups of enterotoxigenic *Escherichia coli* isolated from pre-weaning pigs in north Vietnam, *Journal of Medical Microbiology*, **55:93-9**.
- Driesen SJ, Carland PG, Fahy VA, 1993.** Studies on preweaning piglet diarrhea, *Australian Veterinary Journal*, **70:259-62**.
- Duan H, Chai T, Liu J, Zhang X, Qi C, Gao J, Wang Y, Cai Y, Miao Z, Yao M, Schlenker G, 2009.** Source identification of airborne *Escherichia coli* of swine house surroundings using ERIC-PCR and REP-PCR, *Environmental Research*, **109:511-7**.
- Dubreuil JD, 1997.** *Escherichia coli* STb enterotoxin, *Microbiology*, **143:1783-95**.

**E**

- Elicker S, Philadelphia D, Schilcher F, Fischer L, Weissenbock H, Sipos W, 2010.** Is pathomorphology sufficient for diagnosis of enteritic infections? In: S D'Allaire and R Friendship (eds), *Proceedings of the 21<sup>st</sup> International Pig Veterinary Society Congress, Vancouver, Canada*, **pp. 760**.
- Enemark HL, Ahrens P, Bille-Hansen V, Heegaard PM, Vigre H, Thamsborg SM, Lind P, 2003a.** *Cryptosporidium parvum*: infectivity and pathogenicity of the 'porcine' genotype, *Parasitology*, **126:407-16**.
- Enemark HL, Bille-Hansen V, Lind P, Heegaard PMH, Vigre H, Ahrens P, Thamsborg SM, 2003b.** Pathogenicity of *Cryptosporidium parvum*-evaluation of an animal model, *Veterinary Parasitology*, **113:35-57**.
- Erickson AK, Baker DR, Bosworth BT, Casey TA, Benfield DA, Francis DH, 1994.** Characterization of porcine intestinal receptors for the K88ac fimbrial adhesin of *Escherichia coli* as mucin-type sialoglycoproteins, *Infection and Immunity*, **62:5404-10**.
- Erickson AK, Willgohe JA, McFarland SY, Benfield DA, Francis DH, 1992.** Identification of two porcine brush border glycoproteins that bind the K88ac adhesin of *Escherichia coli* and correlation of these glycoproteins with the adhesive phenotype, *Infection and Immunity*, **60:983-8**.

**F**

- Fairbrother JM, 2006.** Neonatal *Escherichia coli* diarrhea. In: BE Straw, S D'Allaire, WL Mengeling and DJ Taylor (eds), *Diseases of Swine, 8th edn*, (Iowa State University Press, Ames), **pp. 641-9**.
- Fairbrother JM, Gyles CL, 2006.** Postweaning *Escherichia coli* Diarrhea and Edema Disease. In: BE Straw, S D'Allaire, WL Mengeling and DJ Taylor (eds), *Diseases of Swine, 8th edn*, (Iowa State University Press, Ames), **pp. 649-62**.

- Fairbrother JM, Nadeau E, Gyles CL, 2005.** *Escherichia coli* in postweaning diarrhea in pigs: an update on bacterial types, pathogenesis, and prevention strategies, *Animal Health Research Reviews*, **1:17-39**.
- Featherstone CA, Marshall JA, Giles M, Sayers AR, Pritchard GC, 2010.** *Cryptosporidium* species infection in pigs in East Anglia, *The Veterinary Record*, **166:51-2**.
- Francis DH, 1999.** Colibacillosis in pigs and its diagnosis, *Journal of Swine Health and Production*, **5:241-4**.
- Frydendahl K, 2002.** Prevalence of serogroups and virulence genes in *Escherichia coli* associated with postweaning diarrhea and edema disease in pigs and a comparison of diagnostic approaches, *Veterinary Microbiology*, **85:169-82**.
- Frydendahl K, Kåre Jensen T, Strodl Andersen J, Fredholm M, Evans G, 2003.** Association between the porcine *Escherichia coli* F18 receptor genotype and phenotype and susceptibility to colonisation and postweaning diarrhea caused by *E. coli* O138:F18, *Veterinary Microbiology*, **93:39-51**.
- Fuentes F, Campal AC, Casas S, Arteaga N, Castro MD, Barranco JA, León L, 2001.** Utilidad del sistema de diagnóstico rápido AuBIODOT-ECET en la detección de cepas de *Escherichia coli* enterotoxigénicas causantes de colibacilosis porcina, *Revista de Salud Animal*, **23:84-90**.
- G**
- Gaafar SM, Dugas S, Symensma R, 1973.** Resistance of pigs recovered from transmissible gastroenteritis against infection with *Ascaris suum* (Abstract), *American Journal of Veterinary Research*, **34:793-5**.
- Garmory HS, Chanter N, French NP, Bueschel D, Songer JG, Titball RW, 2000.** Occurrence of *Clostridium perfringens* beta2-toxin amongst animals, determined using genotyping and subtyping PCR assays, *Epidemiology and Infection*, **124:61-7**.
- Gibert M, Jolivet-Reynaud C, Popoff MR, 1997.** Beta2 toxin, a novel toxin produced by *Clostridium perfringens*, *Gene*, **203:65-73**.
- Ginter A, Williamson ED, Dessy F, Coppe P, Bullifent H, Howells A, Titball RW, 1996.** Molecular variation between the alpha-toxins from the type strain (NCTC 8237) and clinical isolates of *Clostridium perfringens* associated with disease in man and animals, *Microbiology*, **142:191-8**.
- González A, Almaguel R, Crespo A, Ly J, 2007.** Cuban management in pig meat production, *Revista Computadorizada de Producción Porcina*, **14:249-51**.
- Goyal SM, Rademacher RA, Pomeroy KA, 1987.** Comparison of electron microscopy with three commercial tests for the detection of rotavirus in animal feces, *Diagnostic Microbiology and Infectious Diseases*, **6:249-54**.
- Grange PA, Erickson AK, Levery SB, Francis DH, 1999.** Identification of an intestinal neutral glycosphingolipid as a phenotype-specific receptor for the K88ad fimbrial adhesin of *Escherichia coli*, *Infection and Immunity*, **67:165-72**.



- Grange PA, Mouricout MA, 1996.** Transferrin associated with the porcine intestinal mucosa is a receptor specific for K88ab fimbriae of *Escherichia coli*, *Infection and Immunity*, **64:606-10**.
- Griffith RW, Schwartz KJ, Meyerholz DK, 2006.** *Salmonella*. In: BE Straw, S D'Allaire, WL Mengeling and DJ Taylor (eds), *Diseases of Swine, 8th edn*, (Iowa State University Press, Ames), **pp. 739-54**.
- Greenberg HB, Estes MK, 2009.** Rotaviruses: from pathogenesis to vaccination, *Gastroenterology*, **136:1939-51**.
- GRUPOR, 2008.** Inform of the Epizootiology department of the Cuban Company for Pork Production (GRUPOR) presented in the frame of a Workshop on Swine Health and Production, Capitol, Havana city, Cuba.
- Guérin G, Duval-Iflah Y, Bonneau M, Bertaud M, Guillaume P, Ollivier L, 1993.** Evidence for linkage between K88ab, K88ac intestinal receptors to *Escherichia coli* and transferrin loci in pigs (Abstract), *Animal Genetics*, **24:393-6**.
- Gurtner C, Popescu F, Wyder M, Sutter E, Zeeh F, Frey J, von Schubert C, Posthaus H, 2010.** Rapid cytopathic effects of *Clostridium perfringens* beta-toxin on porcine endothelial cells, *Infection and Immunity*, **78:2966-73**.
- Guselle NJ, Appelbee AJ, Olson ME, 2003.** Biology of *Cryptosporidium parvum* in pigs: from weaning to market, *Veterinary Parasitology*, **113:7-18**.

## H

- Halaihel N, Masía RM, Fernández-Jiménez M, Ribes JM, Montava R, De Blas I, Gironés O, Alonso JL, Buesa J, 2010.** Enteric calicivirus and rotavirus infections in domestic pigs, *Epidemiology and Infection*, **138:542-8**.
- Hamnes IS, Gjerde BK, Forberg T, Robertson LJ, 2007.** Occurrence of *Cryptosporidium* and *Giardia* in suckling piglets in Norway, *Veterinary Parasitology*, **144:222-33**.
- Hansen GH, Delmas B, Besnardeau L, Vogel LK, Laude H, Sjöström H, Norén O, 1998.** The coronavirus transmissible gastroenteritis virus causes infection after receptor-mediated endocytosis and acid-dependent fusion with an intracellular compartment, *Journal of Virology*, **72:527-34**.
- Harada K, Asai T, Ozawa M, Kojima A, Takahashi T, 2008.** Farm-level impact of therapeutic antimicrobial use on antimicrobial-resistant populations of *Escherichia coli* isolates from pigs, *Microbial Drug Resistance*, **14:239-44**.
- Hariharan H, Coles M, Poole D, Page R, 2004.** Antibiotic resistance among enterotoxigenic *Escherichia coli* from piglets and calves with diarrhea, *Canadian Veterinary Journal*, **45:605-6**.
- Hariharan H, Coles M, Poole D, Page R, 2004.** Antibiotic resistance among enterotoxigenic *Escherichia coli* from piglets and calves with diarrhea, *Canadian Veterinary Journal*, **45:605-6**.
- Harleman JH, Meyer RC, 1984.** Life cycle of *Isospora suis* in gnotobiotic and conventionalized piglets, *Veterinary Parasitology*, **17:27-39**.

Hendriksen RS, Mevius DJ, Schroeter A, Teale C, Jouy E, Butaye P, Franco A, Utinane A, Amado A, Moreno M, Greko C, Stärk KDC, Berghold C, Myllyniemi AL, Hoszowski A, Sunde M, Aarestrup FM, 2008. Occurrence of antimicrobial resistance among bacterial pathogens and indicator bacteria in pigs in different European countries from year 2002 – 2004: the ARBAO-II study, *Acta Veterinaria Scandinavica*, **50**:19.

Hendriksen SW, van Leengoed LA, Roest HI, van Nes A, 2006. Neonatal diarrhoea in pigs: alpha- and beta2-toxin produced by *Clostridium perfringens* (Abstract), *Tijdschrift voor Diergeneeskunde*, **131**:910-3.

Hong TT, Linh NQ, Ogle B, Lindberg JE, 2006. Survey on the prevalence of diarrhea in pre-weaning piglets and on feeding systems as contributing risk factors in smallholdings in Central Vietnam, *Tropical Animal Health and Production*, **38**:397-405.

Hunter SE, Brown JE, Oyston PC, Sakurai J, Titball RW, 1993. Molecular genetic analysis of beta-toxin of *Clostridium perfringens* reveals sequence homology with alpha-toxin, gamma-toxin, and leukocidin of *Staphylococcus aureus*, *Infection and Immunity*, **61**:3958-65.

Hur J, Choi YY, Park JH, Jeon BW, Lee HS, Kim AR, Lee JH, 2011. Antimicrobial resistance, virulence-associated genes, and pulsed-field gel electrophoresis profiles of *Salmonella enterica* subsp. *enterica* serovar Typhimurium isolated from piglets with diarrhea in Korea, *Canadian Journal of Veterinary Research*, **75**:49-56.

## I

Ibenyassine K, AitMhand R, Karamoko Y, Cohen N, Nennaji MM, 2006. Use of repetitive DNA sequences to determine the persistence of enteropathogenic *Escherichia coli* in vegetables and in soil grown in fields treated with contaminated irrigation water, *Letters in Applied Microbiology*, **43**:528-33.

IMV, 2003. Informe preliminar sobre el síndrome gastroentérico del cerdo. Institute of Veterinary Medicine. Ministry of Agriculture, Havana, Cuba. April 2003.

Izumiyama S, Furukawa I, Kuroki T, Yamai S, Sugiyama H, Yagita K, Endo T, 2001. Prevalence of *Cryptosporidium parvum* infections in weaned piglets and fattening porkers in Kanagawa Prefecture, Japan, *Japanese Journal of Infectious Diseases*, **54**:23-6.

## J

Jabrane A, Elazhary Y, Talbot BG, Ethier R, Dubuc C, Assaf R, 1992. Porcine respiratory coronavirus in Quebec: Serological studies using a competitive inhibition enzyme-linked immunosorbent assay, *The Canadian Veterinary Journal*, **33**:727-33.

Janke BH, Morehouse LG, Solorzano RF, 1988. Single and mixed infections of neonatal pigs with rotaviruses and enteroviruses: clinical signs and microscopic lesions, *Canadian Journal of Veterinary Research*, **52**:364-9.



- Jensen GM, Frydendahl K, Svendsen O, Jørgensen CB, Cirera S, Fredholm M, Nielsen JP, Møller K, 2006. Experimental infection with *Escherichia coli* O149:F4ac in weaned piglets, *Veterinary Microbiology*, **115**:243-9.
- Jiménez G, Correa I, Melgosa MP, Bullido MJ, Enjuanes L, 1986. Critical epitopes in transmissible gastroenteritis virus neutralization, *Journal of Virology*, **60**:131-9.
- Johnson J, Buddle R, Reid S, Armson A, Ryan UM, 2008a. Prevalence of *Cryptosporidium* genotypes in pre and post-weaned pigs in Australia, *Experimental Parasitology*, **119**:418-21.
- Johnson J, Samarasinghe B, Buddle R, Armson A, Ryan U, 2008b. Molecular identification and prevalence of *Isospora* sp. in pigs in Western Australia using a PCR-RFLP assay, *Experimental Parasitology*, **120**:191-3.
- Jørgensen CB, Cirera S, Archibald AL, Anderson L, Fredholm M, Edfors-Lilja I, 2004. Porcine polymorphisms and methods for detecting them. *International application published under the patent cooperation treaty (PCT)*, PCT/DK2003/000807 or WO2004/048606-A2.
- Jung K, Kang BK, Lee CS, Song DS, 2008. Impact of porcine group A rotavirus co-infection on porcine epidemic diarrhea virus pathogenicity in piglets, *Research in Veterinary Science*, **84**:502-6.
- Jung K, Kim J, Kim O, Kim B, Chae C, 2003. Differentiation between porcine epidemic diarrhea virus and transmissible gastroenteritis virus in formalin-fixed paraffin-embedded tissues by multiplex RT-nested PCR and comparison with in situ hybridization, *Journal of Virological Methods*, **108**:41-7.
- Jurkovič D, Křižková L, Sojka M, Takáčová M, Dusinský R, Krajcovic J, 2007. Genetic diversity of *Enterococcus faecium* isolated from Bryndza cheese, *International Journal of Food Microbiology*, **116**:82-7.
- K**
- Kang JH, Kwon DH, Chung TW, Kim YD, Lee HG, Kim JW, Choe IS, Kim KW, Lim JS, Song EY, Kim CH, 2007. Development of a simple and rapid immunochromatographic strip test for diarrhea-causative porcine rotavirus in swine stool, *Journal of Virological Methods*, **146**:74-9.
- Karamon J, Ziomko I, Cencek T, 2007. Prevalence of *Isospora suis* and *Eimeria* spp. in suckling piglets and sows in Poland, *Veterinary Parasitology*, **147**:171-5.
- Katsuda K, Kohmoto M, Kawashima K, Tsunemitsu H, 2006. Frequency of enteropathogen detection in suckling and weaned pigs with diarrhea in Japan, *Journal of Veterinary Diagnostic Investigation*, **18**:350-4.
- Kavanagh OV, Ajami NJ, Cheng E, Ciarlet M, Guerrero RA, Zeng CQ, Crawford SE, Estes MK, 2010. Rotavirus enterotoxin NSP4 has mucosal adjuvant properties, *Vaccine*, **28**:3106-11.
- Kemeny LJ, 1976. Antibody response in pigs inoculated with transmissible gastroenteritis virus and cross reactions among ten isolates, *Canadian Journal of Comparative Medicine*, **40**:209-14.

- Khamrin P, Maneekarn N, Peerakome S, Chan-it W, Yagyu F, Okitsu S, Ushijima H, 2007.** Novel porcine rotavirus of genotype P[27] shares new phylogenetic lineage with G2 porcine rotavirus strain, *Virology*, **361:243-52**.
- Kim HJ, Park SI, Ha TP, Jeong YJ, Kim HH, Kwon HJ, Kang MI, Cho KO, Park SJ, 2010a.** Detection and genotyping of Korean porcine rotaviruses, *Veterinary Microbiology*, **144:274-86**.
- Kim L, Hayes J, Lewis P, Parwani AV, Chang KO, Saif LJ, 2000.** Molecular characterization and pathogenesis of transmissible gastroenteritis coronavirus (TGEV) and porcine respiratory coronavirus (PRCV) field isolates co-circulating in a swine herd, *Archives of Virology*, **145:1133-47**.
- Kim YJ, Kim JH, Hur J, Lee JH, 2010b.** Isolation of *Escherichia coli* from piglets in South Korea with diarrhea and characteristics of the virulence genes, *Canadian Journal of Veterinary Research*, **74:59-64**.
- Knezevic P, Petrovic O, 2008.** Antibiotic resistance of commensal *Escherichia coli* of food-producing animals from three Vojvodinian farms, Serbia, *International Journal of Antimicrobial Agents*, **31:360-3**.
- Komárek A, 2009.** A new R package for Bayesian estimation of multivariate normal mixtures allowing for selection of the number of components and interval-censored data, *Computational Statistics and Data Analysis*, **53:3932-47**.
- Koudela B, Gómez E, Abreu R, Vítovec J, 1989.** Porcine neonatal coccidiosis in Cuba, *Folia Parasitologica*, **36:31-2**.
- Kvác M, Hanzlíková D, Sak B, Kvetonová D, 2009.** Prevalence and age-related infection of *Cryptosporidium suis*, *C. muris* and *Cryptosporidium* pig genotype II in pigs on a farm complex in the Czech Republic, *Veterinary Parasitology*, **160:319-22**.
- Kwon D, Choi C, Jung T, Chung HK, Kim JP, Bae SS, Cho WS, Kim J, Chae C, 2002.** Genotypic prevalence of the fimbrial adhesins (F4, F5, F6, F41 and F18) and toxins (LT, STa, STb and STx2e) in *Escherichia coli* isolated from postweaning pigs with diarrhea or oedema disease in Korea, *The Veterinary Record*, **150:35-7**.
- L**
- Laguna E, 1998.** El cerdo Ibérico en la colonización y en el poblamiento porcino de América, *Solo Cerdo Ibérico*, **1(10)**.
- Laitat M, Ianosi V, Delleur V, Farnir F, Losson B, Gevaert D, Saegerman C, 2010.** Prevalence of *Isospora suis* in the South of Belgium. In: S D'Allaire and R Friendship (eds), *Proceedings of the 21<sup>st</sup> International Pig Veterinary Society Congress, Vancouver, Canada*, pp. **802**.
- Lazo L, Gutiérrez Y, 2011.** Estudio de factores de riesgo vinculados a la mortalidad neonatal en una granja porcina, Available in: <http://www.engormix.com/MA-porcicultura/sanidad/articulos/mortalidad-en-lechones-t3348/165-p0.htm>.

- Lazo L, Hernández L, Cruz E, Maroto LO, Manso MJ, Molina R, Rodríguez A, 2005.** Consideraciones epizootiológicas sobre el comportamiento de la colibacilosis entérica porcina en la provincia de Villa Clara, *Revista de Salud Animal*, **27:129-32**.
- Lecce JG, Balsbaugh RK, Clare DA, King MW, 1982.** Rotavirus and hemolytic enteropathogenic *Escherichia coli* in weanling diarrhea of pigs, *Journal of Clinical Microbiology*, **16:715-23**.
- Lee SI, Kang SG, Kang ML, Yoo SH, 2008.** Development of multiplex polymerase chain reaction assays for detecting enterotoxigenic *Escherichia coli* and their application to field isolates from piglets with diarrhea, *Journal of Veterinary Diagnostic Investigation*, **20:492-6**.
- Lee SI, Rayamahji N, Lee WJ, Cha SB, Shin MK, Roh YM, Yoo HS, 2009.** Genotypes, antibiogram, and pulsed-field gel electrophoresis profiles of *Escherichia coli* strains from piglets in Korea. *Journal of Veterinary Diagnostic Investigation*, **21:510-6**.
- Leten J, Smets K, Claerebout E, Mundt HC, Heesen H, Vercruysse J, 2002.** Isosporosis in suckling piglets in Flanders, *Vlaams Diergeneeskundig Tijdschrift*, **71:63-7**.
- Li BX, Ge JW, Li YJ, 2007.** Porcine aminopeptidase N is a functional receptor for the PEDV coronavirus, *Virology*, **365:166-72**.
- Literak I, Dolejska M, Rybarikova J, Cizek A, Strejckova P, Vyskocilova M, Friedman M, Klimes J, 2009.** Highly variable patterns of antimicrobial resistance in commensal *Escherichia coli* isolates from pigs, sympatric rodents, and flies, *Microbial Drug Resistance*, **15:229-37**.
- López L, Venteo, A, García M, Camuñas A, Ranz A, García J, Sarraseca J, Anaya C, Rueda P, 2009.** Antigen-capture blocking enzyme-linked immunosorbent assay based on a baculovirus recombinant antigen to differentiate Transmissible gastroenteritis virus from porcine respiratory coronavirus antibodies, *Journal of Veterinary Diagnostic Investigation*, **21:598-608**.
- Lundgren O, Peregrin AT, Persson K, Kordasti S, Uhnöo I, Svensson L, 2000.** Role of the enteric nervous system in the fluid and electrolyte secretion of rotavirus diarrhea, *Science*, **287:491-5**.

**M**

- Maddox-Hyttel C, Langkjaer RB, Enemark HL, Vigre H, 2006.** *Cryptosporidium* and *Giardia* in different age groups of Danish cattle and pigs--occurrence and management associated risk factors, *Veterinary Parasitology*, **141:48-59**.
- Madoroba E, Van Driessche E, De Greve H, Mast J, Ncube I, Read J, Beeckmans S, 2009.** Prevalence of enterotoxigenic *Escherichia coli* virulence genes from scouring piglets in Zimbabwe, *Tropical Animal Health and Production*, **41:1539-47**.
- Maidhof H, Guerra B, Abbas S, Elsheikha HM, Whittam TS, Beutin L, 2002.** A multiresistant clone of Shiga toxin-producing *Escherichia coli* O118:[H16] is spread in cattle and humans over different European countries, *Applied and Environmental Microbiology*, **68:5834-42**.
- Manges AR, Dietrich PS, Riley LW, 2004.** Multidrug-resistant *Escherichia coli* clonal groups causing community acquired pyelonephritis, *Clinical Infectious Diseases*, **38:329-34**.

- Martella V, Bányai K, Lorusso E, Bellacicco AL, Decaro N, Camero M, Bozzo G, Moschidou P, Arista S, Pezzotti G, Lavazza A, Buonavoglia C, 2007.** Prevalence of group C rotaviruses in weaning and post-weaning pigs with enteritis, *Veterinary Microbiology*, **123:26-33**.
- Martella V, Pratelli A, Greco G, Tempesta M, Ferrari M, Losio MN, Buonavoglia C, 2001.** Genomic characterization of porcine rotaviruses in Italy, *Clinical and Diagnostic Laboratory Immunology*, **8:129-32**.
- Matthijnsens J, Ciarlet M, Rahman M, Attoui H, Bányai K, Estes MK, Gentsch JR, Iturriza-Gómara M, Kirkwood CD, Martella V, Mertens PP, Nakagomi O, Patton JT, Ruggeri FM, Saif LJ, Santos N, Steyer A, Taniguchi K, Desselberger U, Van Ranst M, 2008.** Recommendations for the classification of group A rotaviruses using all 11 genomic RNA segments, *Archives of Virology*, **153:1621-9**.
- Mattion NM, Bellinzoni RC, Blackhall JO, La Torre JL, Scodeller EA, 1989.** Antigenic characterization of swine rotaviruses in Argentina, *Journal of Clinical Microbiology*, **27:795-8**.
- Maynard C, Fairbrother JM, Bekal S, Sanschagrin F, Levesque RC, Brousseau R, Masson L, Larivière S, Harel J, 2003.** Antimicrobial Resistance Genes in Enterotoxigenic *Escherichia coli* O149:K91 Isolates Obtained over a 23-Year Period from Pigs, *Antimicrobial Agents and Chemotherapy*, **47:3214-21**.
- McClane BA, 1996.** An overview of *Clostridium perfringens* enterotoxi, *Toxicon*, **34:1335-43**.
- Meijerink E, Fries R, Vögeli P, Masabanda J, Wigger G, Stricker C, Neuenschwander S, Bertschinger HU, Stranzinger G, 1997.** Two alpha(1,2) fucosyltransferase genes on porcine chromosome 6q11 are closely linked to the blood group inhibitor (S) and *Escherichia coli* F18 receptor (ECF18R) loci. *Mammalian Genome*, **8:736-41**.
- Meijerink E, Neuenschwander S, Fries R, Dinter A, Bertschinger HU, Stranzinger G, Vögeli P, 2000.** A DNA polymorphism influencing alpha(1,2)fucosyltransferase activity of the pig FUT1 enzyme determines susceptibility of small intestinal epithelium to *Escherichia coli* F18 adhesion, *Immunogenetics*, **52:129-36**.
- Melin L, Mattsson S, Katouli M, Wallgren P, 2004.** Development of post-weaning diarrhea in piglets. Relation to presence of *Escherichia coli* strains and rotavirus, *Journal of Veterinary Medicine B-Infectious Diseases and Veterinary Public Health*, **51:12-22**.
- Mišić Z, Katic-Radivojević S, Kulišić Z, 2003.** *Cryptosporidium* infections in nursing, weaning and post-weaned piglets and sows in the Belgrade district (Abstract), *Acta Veterinaria (Beograd)*, **5-6:361-6**.
- Miyazaki A, Fukuda M, Kuga K, Takagi M, Tsunemitsu H, 2010.** Prevalence of Antibodies against Transmissible Gastroenteritis Virus and Porcine Respiratory Coronavirus among Pigs in Six Regions in Japan, *The Journal of Veterinary Medical Science*, **72:943-6**.
- Moon HW, Hoffman LJ, Cornick NA, Booher SL, Bosworth BT, 1999.** Prevalences of some virulence genes among *Escherichia coli* isolates from swine presented to a diagnostic laboratory in Iowa, *Journal of Veterinary Diagnostic Investigation*, **6:557-60**.

- Moredo FA, Vigo GB, Cappuccio JA, Piñeyro P, Perfumo CJ, Giacoboni GI, 2007.** Antimicrobial resistance of *Escherichia coli* isolated from pigs in Argentina, *Revista Argentina de Microbiologia*, **39:227-9**.
- Morin M, Turgeon D, Jolette J, Robinson Y, Phaneuf JB, Sauvageau R, Beauregard M, Teuscher E, Higgins R, Lariviere S, 1983.** Neonatal diarrhea of pigs in Quebec: infectious causes of significant outbreaks, *Canadian Journal of Comparative Medicine*, **47:11-7**.
- Moyaert H, De Graef EM, Haesebrouck F, Decostere A, 2006.** Acquired antimicrobial resistance in the intestinal microbiota of diverse cat populations, *Research in Veterinary Science*, **81:1-7**.
- Mundt HC, Cohnen A, Dauschies A, Joachim A, Prosl H, Schmaschke R, Westphal B, 2005.** Occurrence of *Isospora suis* in Germany, Switzerland and Austria, *Journal of Veterinary Medicine Series B- Infectious Diseases and Veterinary Public Health*, **52:93-7**.
- Mundt HC, Joachi A, Becka M, Dauschies A, 2006.** *Isospora suis*: an experimental model for mammalian intestinal coccidiosis, *Parasitology Research*, **98:167-75**.
- Mundt HC, Joachim A, Dauschies A, Zimmermann M, 2003.** Population Biology Studies on *Isospora suis* in Piglets, *Parasitology Research*, **90:158-9**.
- N**
- Nagy B, Nagy G, Meder M, Mocsári E, 1996.** Enterotoxigenic *Escherichia coli*, rotavirus, porcine epidemic diarrhea virus, adenovirus and calici-like virus in porcine postweaning diarrhea in Hungary (Abstract), *Acta Veterinaria Hungarica*, **44:9-19**.
- Nagy J, Bilkei G, 2003.** Neonatal piglet losses associated with *Escherichia coli* and *Clostridium difficile* infection in a Slovakian outdoor production unit, *The Veterinary Journal*, **166:98-100**.
- Naing L, Winn T, Rusli BN, 2006.** Practical Issues in Calculating the Sample Size for Prevalence Studies, *Archives of Orofacial Sciences*, **1:9-14**.
- Namvar A, Warriner K, 2006.** Application of enterobacterial repetitive intergenic consensus-polymerase chain reaction to trace the fate of generic *Escherichia coli* within a high capacity pork slaughter line, *International Journal of Food Microbiology*, **108:155-63**.
- Nataro JP, Kaper JB, 1998.** Diarrheagenic *Escherichia coli*, *Clinical Microbiology Reviews*, **11:142-201**.
- Naylor CE, Eaton JT, Howells A, Justin N, Moss DS, Titball RW, Basak AK, 1998.** Structure of the key toxin in gas gangrene, *Nature Structural Biology*, **5:738-46**.
- Naylor RD, Martin PK, Barker LT, 1997.** Detection of *Clostridium perfringens* alpha toxin by enzyme-linked immunosorbent assay, *Research in Veterinary Science*, **63:101-2**.
- Nevarez AM, Ramírez R, Niño R, Rodríguez LE, Ramírez E, 1997.** Identificación de *Cryptosporidium* en cerdos con enteritis, *Veterinaria Mexico*, **28:231-4**.
- Niestrath M, Takla M, Joachim A, Dauschies A, 2002.** The role of *Isospora suis* as a pathogen in conventional piglet production in Germany, *Journal of Veterinary Medicine B- Infectious Diseases and Veterinary Public Health*, **49:176-80**.

**NRAG, 1989.** Norma Ramal del Ministerio de la Agricultura 1010: Diagnóstico Veterinario, Clasificación Microbiológica y Métodos de Ensayo (Protocol from the Agriculture Ministry 1010: Veterinary Diagnostics, Microbiologic Classification, and Assay Procedure).

**Nulsen MF, Mor MB, Lawton DE, 2008.** Antibiotic resistance among indicator bacteria isolated from healthy pigs in New Zealand, *New Zealand Veterinary Journal*, **56:29-35**.

**Nuñez A, McNeilly F, Perea A, Sanchez-Cordon PJ, Huerta B, Allan G, Carrasco L, 2003.** Coinfection by *Cryptosporidium parvum* and Porcine Circovirus Type 2 in Weaned Pigs, *Journal of Veterinary Medicine B- Infectious Diseases and Veterinary Public Health*, **50:255-8**.

## O

**Oanh TKN, Nguyen VK, Do TN, Goddeeris BM, De Greve H, 2010.** *Escherichia coli* strains causing edema disease in northern Vietnam share an identical verotoxin 2e, *Tropical Animal Health and Production*, **42:1797-804**.

**Oh JS, Song DS, Park BK, 2003.** Identification of a putative cellular receptor 150 kDa polypeptide for porcine epidemic diarrhea virus in porcine enterocytes, *Journal of Veterinary Science*, **4:269-75**.

**Okhuysen PC, Chappell CL, 2002.** *Cryptosporidium* virulence determinants-are we there yet?, *International Journal for Parasitology*, **32:517-25**.

**Olasz F, Fekete PZ, Blum-Oehler G, Boldogkoi Z, Nagy B, 2005.** Characterization of an F18<sup>+</sup> enterotoxigenic *Escherichia coli* strain from post weaning diarrhoea of swine, and of its conjugative virulence plasmid pTC, *FEMS Microbiology Letters*, **244:281-9**.

**ONE, 2011.** Oficina Nacional de Estadísticas (National Office for Statistics), Agricultura (Agriculture), Available in: <http://www.one.cu>.

**Osek J, 1999.** Prevalence of virulence factors of *Escherichia coli* strains isolated from diarrheic and healthy piglets after weaning, *Veterinary Microbiology*, **68:209-17**.

**Osek J, 2000.** Virulence factors and genetic relatedness of *Escherichia coli* strains isolated from pigs with post-weaning diarrhea, *Veterinary Microbiology*, **71:211-22**.

**Otten A, Takla M, Dauschies A, Rommel M, 1996.** The epizootiology and pathogenic significance of infections with *Isospora suis* in ten piglet production operations in Nordrhein-Westfalen, *Berliner und Münchener tierärztliche Wochenschrift*, **109:220-3**.

## P

**Parra GI, Vidales G, Gomez JA, Fernandez FM, Parreño V, Bok K, 2008.** Phylogenetic analysis of porcine rotavirus in Argentina: increasing diversity of G4 strains and evidence of interspecies transmission, *Veterinary Microbiology*, **126:243-50**.

**Pedroso M, Talavera A, 1983.** Inmunofluorescencia directa para el diagnóstico de *Escherichia coli* cepas K88 y K99 en cerdos diarreicos y no diarreicos, *Revista de Salud Animal*, **5:503-8**.



- Pensaert M, Callebaut P, Vergote J, 1986.** Isolation of a porcine respiratory, non-enteric coronavirus related to transmissible gastroenteritis, *Veterinary Quarterly*, **8:257-61**.
- Pensaert M, Cox E, van Deun K, Callebaut P, 1993.** A sero-epizootiological study of porcine respiratory coronavirus in Belgian swine, *The Veterinary Quarterly*, **15:16-20**.
- Petit L, Gibert M, Popoff MR, 1999.** *Clostridium perfringens*: toxinotype and genotype, *Trends in Microbiology*, **7:104-10**.
- Phillips RM, Westerman RB, 1991.** Enzyme immunofiltration assay for measurement of antibodies to transmissible gastroenteritis virus of swine: comparison with enzyme immunosorbence assay, serum neutralization, and indirect immunofluorescent antibody techniques, *Journal of Veterinary Diagnostic Investigation*, **3:346-8**.
- Plummer M, 2008.** Penalized loss functions for Bayesian model comparison, *Biostatistics*, **9:523-39**.
- Poppe C, Martin L, Muckle A, Archambault M, McEwen S, Weir E, 2006.** Characterization of antimicrobial resistance of *Salmonella* Newport isolated from animals, the environment, and animal food products in Canada, *The Canadian Journal of Veterinary Research*, **70:105-14**.
- Python P, Jörg H, Neuenschwander S, Hagger C, Stricker C, Bürgi E, Bertschinger HU, Stranzinger G, Vögeli P, 2002.** Fine-mapping of the intestinal receptor locus for enterotoxigenic *Escherichia coli* F4ac on porcine chromosome 13, *Animal Genetics*, **33:441-7**.

## Q

- Quilez J, Sanchez-Acedo C, Clavel A, del Cacho E, Lopez-Bernad F, 1996.** Prevalence of *Cryptosporidium* infections in pigs in Aragon (northeastern Spain), *Veterinary Parasitology*, **67:83-8**.

## R

- Rácz ML, Kroeft SS, Munford V, Caruzo TA, Durigon EL, Hayashi Y, Gouvea V, Palombo EA, 2000.** Molecular characterization of porcine rotaviruses from the southern region of Brazil: characterization of an atypical genotype G[9] strain, *Journal of Clinical Microbiology*, **38:2443-6**.
- Ramchandani M, Manges AR, DebRoy C, Smith SP, Johnson JR, Riley LW, 2005.** Possible animal origin of human-associated, multidrug-resistant, uropathogenic *Escherichia coli*, *Clinical Infectious Diseases*, **40:251-7**.
- Ramos AP, Stefanelli CC, Linhares RE, de Brito BG, Santos N, Gouvea V, de Cassia Lima R, Nozawa C, 2000.** The stability of porcine rotavirus in feces, *Veterinary Microbiology*, **71:1-8**.
- Rasschaert D, Duarte M, Laude H, 1990.** Porcine respiratory coronavirus differs from transmissible gastroenteritis virus by a few genomic deletions, *Journal of General Virology*, **71:2599-607**.
- Rasschaert K, 2008.** Identification of a novel F4 receptor involved in endocytosis and transcytosis. Chapter 4: Aminopeptidase N is a receptor for F4 *Escherichia coli*. PhD thesis. Faculty of Veterinary Medicine, Ghent University. ISBN 978-90-5864-130-4, **pp. 64-83**.

- Rasschaert K, Verdonck F , Goddeeris BM , Duchateau L, Cox E, 2007.** Screening of pigs resistant to F4 enterotoxigenic *Escherichia coli* (ETEC) infection, *Veterinary Microbiology*, **123:249-53**.
- Ricardo O, 2008.** Enfermedades entericas contagiosas del cerdo en Cuba, *Revista de la Asociación Cubana de Producción Animal (ACPA)*, **3:18-9**.
- Rico C, 2005.** Programa de mejora genética en Cuba. Su vinculación con el sector cooperativo y privado. VIII Encuentro de Nutrición y Producción de Animales Monogástricos. Available in: URL: [http://avpa.ula.ve/eventos/viii\\_encuentro\\_mono\\_gastricos/memorias/conferencia-2.pdf](http://avpa.ula.ve/eventos/viii_encuentro_mono_gastricos/memorias/conferencia-2.pdf).
- Roberts L, Walker EJ, 1982.** Field study of coccidial and rotaviral diarrhea in unweaned piglets, *The Veterinary Record*, **110:11-3**.
- Rodák L, Smíd B, Nevoránková Z, Smítalová R, Valíček L, 2004.** Verification of sensitivity and specificity of group A rotavirus detection in piglets faeces with monoclonal blocking ELISA methods, *Journal of Veterinary Medicine B- Infectious Diseases and Veterinary Public Health*, **51:160-5**.
- Rodák L, Smítalová R, Smíd B, Nevoránková Z, Valíček L, Pšikal I, 2009.** Comparing electron microscopy and a competitive blocking ELISA in the detection of rotaviruses in porcine faeces, *The Veterinary Journal*, **187:279-81**.
- Rosengren LB, Waldner CL, Reid-Smith RJ, 2009.** Associations between antimicrobial resistance phenotypes, antimicrobial resistance genes, and virulence genes of fecal *Escherichia coli* isolates from healthy grow-finish pigs, *Applied and Environmental Microbiology*, **75:1373-80**.
- Rousset E, Harel J, Dubreuil JD, 1998.** Sulfatide from the pig jejunum brush border epithelial cell surface is involved in binding of *Escherichia coli* enterotoxin b, *Infection and Immunity*, **66:5650-8**.
- Ruan X, Liu M, Casey TA, Zhang W, 2011.** A tripartite fusion, FaeG-FedF-LT(192)A2:B, of enterotoxigenic *Escherichia coli* (ETEC) elicits antibodies that neutralize cholera toxin, inhibit adherence of K88 (F4) and F18 fimbriae, and protect pigs against K88ac/heat-labile toxin infection, *Clinical and Vaccine Immunology*, **18:1593-9**.
- Ryan UM, Samarasinghe B, Read C, Buddle JR, Robertson ID, Thompson RC, 2003.** Identification of a novel *Cryptosporidium* genotype in pigs, *Applied and Environmental Microbiology*, **69:3970-4**.

## S

- Saif LJ, Sestak K, 2006.** Transmissible Gastroenteritis and Porcine Respiratory Coronavirus. In: BE Straw, S D'Allaire, WL Mengeling, DJ Taylor (eds), *Diseases of Swine. 8<sup>th</sup> edn*. Ames, (Iowa State University Press), pp. **489-516**.
- Saliki JT, Rodgers SJ, Eskew G, 1998.** Serosurvey of selected viral and bacterial diseases in wild swine from Oklahoma, *Journal of Wildlife Diseases*, **34:834-8**.
- Samarasinghe B, Johnson J, Ryan U, 2008.** Phylogenetic analysis of *Cystoisospora* species at the rRNA ITS1 locus and development of a PCR-RFLP assay, *Experimental Parasitology*, **118:592-5**.



- Sanford SE, 1987.** Enteric cryptosporidial infection in pigs: 184 cases (1981-1985), *Journal of the American Veterinary Medical Association*, **190:695-8**.
- Santos N, Hoshino Y, 2005.** Global distribution of rotavirus serotypes/genotypes and its implication for the development and implementation of an effective rotavirus vaccine, *Reviews in Medical Virology*, **15:29-56**.
- Savic B, Pavlovic I, Ivetic V, Zutic M, Kureljusic B, 2010.** Prevalence of isosporidial infection in piglets with clinical signs of enteropathy. In: S D'Allaire and R Friendship (eds), *Proceedings of the 21<sup>st</sup> International Pig Veterinary Society Congress, Vancouver, Canada*, **pp. 800**.
- Saxena MK, Kumar AA, Chaudhari P, Shivachandra SB, Singh VP, Sharma B, 2005.** Ribotyping of Indian isolates of *Pasteurella multocida* based on 16S and 23S rRNA genes, *Veterinary Research Communications*, **29:527-35**.
- Schwegmann-Wessels C, Herrler G, 2006.** Transmissible gastroenteritis virus infection: a vanishing specter, *Deutsche Tierärztliche Wochenschrift*, **113:157-9**.
- Schwegmann-Wessels C, Zimmer G, Laude H, Enjuanes L, Herrler G, 2002.** Binding of transmissible gastroenteritis coronavirus to cell surface sialoglycoproteins, *The Journal of Virology*, **76:6037-43**.
- Sears CL, Kaper JB, 1996.** Enteric bacterial toxins: mechanisms of action and linkage to intestinal secretion, *Microbiological Reviews*, **60:167-215**.
- Sedlak K, Bartova E, Machova J, 2008.** Antibodies to selected viral disease agents in wild boars from the Czech Republic, *Journal of Wildlife Diseases*, **44:777-80**.
- Sellwood R, Gibbons RA, Jones GW, Rutter JM, 1975.** Adhesion of enteropathogenic *Escherichia coli* to pig intestinal brush borders: the existence of two pig phenotypes, *Journal of Medical Microbiology*, **8:405-11**.
- Sestak K, Zhou Z, Shoup DI, Saif LJ, 1999.** Evaluation of the baculovirus-expressed S glycoprotein of transmissible gastroenteritis virus (TGEV) as antigen in a competition ELISA to differentiate porcine respiratory coronavirus from TGEV antibodies in pigs, *Journal of Veterinary Diagnostic Investigation*, **11:205-14**.
- Shi QS, Xie XM, Liu XC, Huang SQ, He CQ, 2002.** Experimental results on enterotoxigenic *E.coli* F18 receptor genotypes, *Yi Chuan*, **24:656-8**.
- Skampardonis V, Sotiraki S, Kostoulas P, Leontides L, 2010.** Effect of toltrazuril treatment in nursing piglets naturally infected with *Isospora suis*, *Veterinary Parasitology*, **172:46-52**.
- Smet A, Martel A, Persoons D, Dewulf J, Heyndrickx M, Cloeckaert A, Praud K, Claeys G, Catry B, Herman L, Haesebrouck F, Butaye P, 2009.** Comparative analysis of extended-spectrum-beta-lactamase-carrying plasmids from different members of *Enterobacteriaceae* isolated from poultry, pigs and humans: evidence for a shared beta-lactam resistance gene pool?, *Journal of Antimicrobial Chemotherapy*, **63:1286-300**.
- Smith MG, Jordan D, Chapman TA, Chin JJ, Barton MD, Do TN, Fahy VA, Fairbrother JM, Trott DJ, 2010.** Antimicrobial resistance and virulence gene profiles in multi-drug resistant

- enterotoxigenic *Escherichia coli* isolated from pigs with post-weaning diarrhea, *Veterinary Microbiology*, **145:299-307**.
- Song DS, Kang BK, Oh JS, Ha GW, Yang JS, Moon HJ, Jang YS, Park BK, 2006.** Multiplex reverse transcription-PCR for rapid differential detection of porcine epidemic diarrhea virus, transmissible gastroenteritis virus, and porcine group A rotavirus, *Journal of Veterinary Diagnostic Investigation*, **18:278-81**.
- Songer JG, 1996.** Clostridial enteric diseases of domestic animals, *Clinical Microbiology Reviews*, **9:216-34**.
- Songer JG, Meer RR, 1996.** Genotyping of *Clostridium perfringens* by polymerase chain reaction is a useful adjunct to diagnosis of clostridial enteric disease in animals, *Anaerobe*, **2:197-203**.
- Songer JG, Uzal FA, 2005.** Clostridial enteric infections in pigs, *Journal of Veterinary Diagnostic Investigation*, **17:528-36**.
- Spaan W, Cavanagh D, Horzinek MC, 1988.** Coronaviruses: structure and genome expression, *Journal of General Virology*, **69:2939-52**.
- Steel RB, Torres-Medina A, 1984.** Effects of environmental and dietary factors on human rotavirus infection in gnotobiotic piglets, *Infection and Immunity*, **43:906-11**.
- Steyer A, Poljsak-Prijatelj M, Barlic-Maganja D, Marin J, 2008.** Human, porcine and bovine rotaviruses in Slovenia: evidence of interspecies transmission and genome reassortment, *Journal of General Virology*, **89:1690-8**.
- Straw BE, Dewey CE, Wilson MR, 2006.** Differential diagnosis of disease. In: BE Straw, S D'Allaire, WL Mengeling and DJ Taylor (eds), *Diseases of Swine, 8th edn*, (Iowa State University Press, Ames), pp. **244-8**.
- Suárez-Luengas L, Clavel A, Quílez, J, Goñi-Cepero MP, Torres E, Sánchez-Acedo C, del Cacho E, 2007.** Molecular characterization of *Cryptosporidium* isolates from pigs in Zaragoza (northeastern Spain), *Veterinary Parasitology*, **148:231-5**.
- Svensmark B, Askaa J, Wolstrup C, Nielsen K, 1989a.** Epidemiological studies of piglet diarrhea in intensively managed Danish sow herds. IV. Pathogenicity of porcine rotavirus, *Acta Veterinaria Scandinavica*, **30:71-6**.
- Svensmark B, Nielsen K, Dalsgaard K, Willeberg P, 1989b.** Epidemiological studies of piglet diarrhea in intensively managed Danish sow herds. III. Rotavirus infection, *Acta Veterinaria Scandinavica*, **30:63-70**.

**T**

- Teneberg S, Hirst TR, Angström J, Karlsson KA, 1994.** Comparison of the glycolipid-binding specificities of cholera toxin and porcine *Escherichia coli* heat-labile enterotoxin: identification of a receptor-active non-ganglioside glycolipid for the heat-labile toxin in infant rabbit small intestine (Abstract), *Glycoconjugate Journal*, **11:533-40**.

- Thomson JR, 2006.** Diseases of the digestive system. In: BE Straw, S D’Allaire, WL Mengeling, DJ Taylor (eds.), *Diseases of Swine*. 8<sup>th</sup> edn. Ames, (Iowa State University Press), pp. 244-8.
- Thorsteinsdottir TR, Haraldsson G, Fridriksdottir V, Kristinsson KG, Gunnarsson E, 2010.** Prevalence and Genetic Relatedness of Antimicrobial-Resistant *Escherichia coli* Isolated From Animals, Foods and Humans in Iceland, *Zoonoses and Public Health*, **57:189-96**.
- Thrusfield M, 1997.** Veterinary Epidemiology, (Blackwell Science Ltd., Oxford), pp. 182-4.
- Tian GB, Wang HN, Zou LK, Tang JN, Zhao YW, Ye MY, Tang JY, Zhang Y, Zhang AY, Yang X, Xu CW, Fu YJ, 2009.** Detection of CTX-M-15, CTX-M-22, and SHV-2 Extended-Spectrum b-Lactamases (ESBLs) in *Escherichia coli* fecal-sample isolates from pig farms in China, *Foodborne Pathogens and Disease*, **6:297-304**.
- Tiels P, Verdonck F, Coddens A, Goddeeris B, Cox E, 2008.** The excretion of F18<sup>+</sup> *E. coli* is reduced after oral immunization of pigs with a FedF and F4 fimbriae conjugate, *Vaccine*, **26:2154-63**.
- Torres JM, Sánchez C, Suñé C, Smerdou C, Prevec L, Graham F, Enjuanes L, 1995.** Induction of antibodies protecting against transmissible gastroenteritis coronavirus (TGEV) by recombinant adenovirus expressing TGEV spike protein, *Virology*, **213:503-16**.
- Travis R, Gyles CL, Reid-Smith R, Poppe C, McEwen SA, Friendship R, Janecko N, Boerlin P, 2006.** Chloramphenicol and kanamycin resistance among porcine *Escherichia coli* in Ontario, *Journal of Antimicrobial Chemotherapy*, **58:173-7**.

## U

- Ursu K, Kisfali P, Rigo D, Ivanics E, Erdelyi K, Dan A, Meleg B, Martella V, Banyai K, 2009.** Molecular analysis of the VP7 gene of pheasant rotaviruses identifies a new genotype, designated G23, *Archives of Virology*, **154:1365-9**.
- Ushida K, Kishimoto A, Piao SJ, Itoh M, Shiga A, Nakanishi N, Tsukahara T, 2009.** An epidemiological survey on pigs showing symptoms of infectious enteric diseases and dyspepsia in Japan, *Animal Science Journal*, **80:556-61**.
- Uysal HK, Boral O, Metiner K, Ilgaz A, 2009.** Investigation of intestinal parasites in pig feces that are also human pathogens, *Turkiye Parazitologii Dergisi*, **33:218-21**.

## V

- Vaandrager AB, van der Wiel E, Hom ML, Luthjens LH, de Jonge HR, 1994.** Heat-stable enterotoxin receptor/guanylyl cyclase C is an oligomer consisting of functionally distinct subunits, which are non-covalently linked in the intestine, *The Journal of Biological Chemistry*, **269:16409-15**.
- Van den Broeck W, Cox E, Goddeeris BM, 1999a.** Induction of immune responses in pigs following oral administration of purified F4 fimbriae, *Vaccine*, **17:2020-9**.
- Van den Broeck W, Cox E, Goddeeris BM, 1999b.** Seroprevalence of F4<sup>+</sup> enterotoxigenic *Escherichia coli* in regions with different pig farm densities, *Veterinary Microbiology*, **69:207-16**.

- Van der Stede Y, Cox E, Goddeeris BM, 2002.** Antigen dose modulates the immunoglobulin isotype responses of pigs against intramuscularly administered F4-fimbriae, *Veterinary Immunology and Immunopathology*, **88:209-16**.
- Varga C, Rajic A, McFall ME, Reid-Smith RJ, Deckert AE, Pearl DL, Avery BP, Checkley SL, McEwen SA, 2008.** Comparison of antimicrobial resistance in generic *Escherichia coli* and *Salmonella* spp. cultured from identical fecal samples in finishing swine, *Canadian Journal of Veterinary Research*, **72:181-7**.
- Vengust G, Valencak Z, Bidovec A, 2006.** A serological survey of selected pathogens in wild boar in Slovenia, *Journal of Veterinary Medicine B- Infectious Diseases and Veterinary Public Health*, **53:24-7**.
- Verdonck F, Cox E, Ampe B, Goddeeris BM, 2003.** Open status of pig-breeding farms is associated with slightly higher seroprevalence of F18<sup>+</sup> *Escherichia coli* in northern Belgium, *Preventive Veterinary Medicine*, **60:133-41**.
- Verdonck F, Cox E, van Gog K, Van der Stede Y, Duchateau L, Deprez P, Goddeeris BM, 2002.** Different kinetic of antibody responses following infection of newly weaned pigs with an F4 enterotoxigenic *Escherichia coli* strain or an F18 verotoxigenic *Escherichia coli* strain, *Vaccine*, **20:2995-3004**.
- Versalovic J, Koeuth T, Lupski JR, 1991.** Distribution of repetitive DNA sequences in eubacteria and application to fingerprinting of bacterial genomes, *Nucleic Acids Research*, **19:6823-31**.
- Vidotto MC, de Lima NCS, Fritzen JTT, de Freitas JC, Venâncio EJ, Ono MA, 2009.** Frequency of virulence genes in *Escherichia coli* strains isolated from piglets with diarrhea in the North Parana State, Brazil, *Brazilian Journal of Microbiology*, **40:199-204**.
- Vieira AR, Houe H, Wegener HC, Danilo MA, Wong LF, Emborg HD, 2009.** Association between tetracycline consumption and tetracycline resistance in *Escherichia coli* from healthy Danish slaughter pigs, *Foodborne Pathogens and Disease*, **6:99-109**.
- Vítovec J, Hamadejová K, Landová L, Kvác M, Kvetonová D, Sak B, 2006.** Prevalence and pathogenicity of *Cryptosporidium suis* in pre- and post-weaned pigs, *Journal of Veterinary Medicine B- Infectious Diseases and Veterinary Public Health*, **53:239-43**.
- Vítovec J, Koudela B, 1990.** Double alteration of the small intestine in conventional and gnotobiotic piglets experimentally infected with the coccidium *Isospora suis* (Apicomplexa, Eimeriidae), *Folia Parasitologica*, **37:21-33**.
- Vu-Khac H, Holoda E, Pilipcinec E, Blanco M, Blanco JE, Dahbi G, Mora A, López C, González EA, Blanco J, 2007.** Serotypes, virulence genes, intimin types and PFGE profiles of *Escherichia coli* isolated from piglets with diarrhea in Slovakia, *The Veterinary Journal*, **174:176-87**.
- Vu-Khac H, Holoda E, Pilipcinec E, Blanco M, Blanco JE, Mora A, Dahbi G, López C, González EA, Blanco J, 2006.** Serotypes, virulence genes, and PFGE profiles of *Escherichia coli* isolated from pigs with postweaning diarrhea in Slovakia, *BMC Veterinary Research*, **2(10):1-8**.

## W

- Wade WF, Gaafar SM, 1981.** Effects of Salmonellosis on subsequent infections with *Ascaris suum* in swine, *Veterinary Parasitology*, **8:309-17**.
- Wang XM, Jiang HX, Liao XP, Liu JH, Zhang WJ, Zhang H, Jiang ZG, Lü DH, Xiang R, Liu YH, 2010.** Antimicrobial resistance, virulence genes, and phylogenetic background in *Escherichia coli* isolates from diseased pigs, *FEMS Microbiology Letters*, **306:15-21**.
- Wang YC, Chang YC, Chuang HL, Chiu CC, Yeh KS, Chang CC, Hsuan SL, Lin WH, Chen TH, 2011.** Transmission of Salmonella between swine farms by the housefly (*Musca domestica*), *Journal of Food Protection*, **74:1012-6**.
- Wattiau P, Van Hesse M, Schlicker C, Vander Veken H, Imberechts H, 2008.** Comparison of classical serotyping and PremiTest assay for routine identification of common *Salmonella enterica* serovars, *Journal of Clinical Microbiology*, **46:4037-40**.
- Weng YB, Hu YJ, Li Y, Li BS, Lin RQ, Xie DH, Gasser RB, Zhu XQ, 2005.** Survey of intestinal parasites in pigs from intensive farms in Guangdong Province, People's Republic of China, *Veterinary Parasitology*, **127:333-6**.
- Wieler LH, Ilieff A, Herbst W, Bauer C, Vieler E, Bauerfeind R, Failing K, Klös H, Wengert D, Baljer G, Zahner H, 2001.** Prevalence of enteropathogens in suckling and weaned piglets with diarrhea in southern Germany, *Journal of Veterinary Medicine B- Infectious Diseases and Veterinary Public Health*, **48:151-9**.
- Williams R, Esterhuysen JJ, Robinson JT, 1994.** Pseudorabies and transmissible gastroenteritis: a serological survey in South Africa, *The Onderstepoort Journal of Veterinary Research*, **61:67-70**.
- Wong I, Moreno M, Molerio M, Valderrama S, Juglar M, Horrach M, Bover E, Borroto A, Basulto R, Calzada L, Hernandez R, Herrera L, de la Fuente J, 1995.** Immunity and protection elicited by a recombinant vaccine against enterotoxigenic *E. coli*, *Biotechnología Aplicada*, **12:9-15**.
- Woods RD, Pirtle EC, Sacks JM, Gibbs EP, 1990.** Serologic survey for transmissible gastroenteritis virus neutralizing antibodies in selected feral and domestic swine sera in the southern United States, *Journal of Wildlife Diseases*, **26:420-2**.

## X

- Xiao L, Escalante L, Yang C, Sulaiman I, Escalante AA, Montali RJ, Fayer R, Lal AA, 1999.** Phylogenetic analysis of *Cryptosporidium* parasites based on the small-subunit rRNA gene locus, *Applied Environmental Microbiology*, **65:1578-83**.

## Y

- Yaeger M, Funk N, Hoffman L, 2002.** A survey of agents associated with neonatal diarrhea in Iowa swine including *Clostridium difficile* and porcine reproductive and respiratory syndrome virus, *Journal of Veterinary Diagnostic Investigation*, **14:281-7**.

- Yan X, Huang X, Ren J, Zou Z, Yang S, Ouyang J, Zeng W, Yang B, Xiao S, Huang L, 2009.** Distribution of *Escherichia coli* F4 adhesion phenotypes in pigs of 15 Chinese and Western breeds and a White DurocxErhualian intercross, *Journal of Medical Microbiology*, **58:1112-7**.
- Yu JR, Seo M, 2004.** Infection status of pigs with *Cryptosporidium parvum*, *Korean Journal of Parasitology*, **42:45-7**.
- Yuan L, Saif LJ, 2002.** Induction of mucosal immune responses and protection against enteric viruses: rotavirus infection of gnotobiotic pigs as a model, *Veterinary Immunology and Immunopathology*, **87:147-60**.
- Yuan L, Stevenson GW, Saif LJ, 2006.** Rotavirus and reovirus. In: BE Straw, S D'Allaire, WL Mengeling, DJ Taylor (eds.), *Diseases of Swine*. 8<sup>th</sup> edn. Ames, (Iowa State University Press), pp. **435-54**.
- Yuan W, Chai TJ, Miao ZM, 2010.** ERIC-PCR identification of the spread of airborne *Escherichia coli* in pig houses, *Science of the Total Environment*, **408:1446-50**.
- Z**
- Zeng J, Deng G, Wang J, Zhou J, Liu X, Xie Q, Wang Y, 2011.** Potential protective immunogenicity of recombinant *Clostridium perfringens*  $\alpha$ - $\beta$ 2- $\beta$ 1 fusion toxin in mice, sows and cows, *Vaccine*, **29:5459-66**.
- Zhang W, Zhao M, Ruesch L, Omot A, Francis D, 2007.** Prevalence of virulence genes in *Escherichia coli* strains recently isolated from young pigs with diarrhea in the US, *Veterinary Microbiology*, **123:145-52**.
- Zhao S, Qiayumi S, Friedman S, Singh R, Foley SL, White DG, McDermott PF, Donkar T, Bolin C, Munro S, Baron EJ, Walker RD, 2003.** Characterization of *Salmonella enterica* serotype Newport isolated from humans and food animals, *Journal of Clinical Microbiology*, **41:5366-71**.
- Zijlstra RT, McCracken BA, Odle J, Donovan SM, Gelberg HB, Petschow BW, Zuckermann FA, Gaskins HR, 1999.** Malnutrition modifies pig small intestinal inflammatory responses to rotavirus, *Journal of Nutrition*, **129:838-43**.
- Zintl A, Neville D, Maguire D, Fanning S, Mulcahy G, Smith HV, De Waal T, 2007.** Prevalence of *Cryptosporidium* species in intensively farmed pigs in Ireland, *Parasitology*, **134:1575-82**.





**Curriculum vitae**

Pedro Yoelvys de la Fé Rodríguez was born on the 21<sup>st</sup> of March 1978 in Santa Clara, Villa Clara, Cuba. He coursed the high school at the Pre-University Institute of Exact Sciences from 1992 to 1995 in Santa Clara. Pedro graduated Doctor in Veterinary Medicine (DVM, awarded with Gold diploma) at Universidad Central "Marta Abreu" de Las Villas (UCLV), Cuba on July 2001, and finished the Master in Parasitology at Institute of Tropical Medicine "Pedro Kourí", Havana city, Cuba on March 2004.

**Professional experience**

From September 2001, he has been teaching at the Faculty of Agriculture and Animal Sciences, UCLV, Cuba. Teaching assistant of the undergraduate courses Veterinary Practice, Biological Agents (Parasitology-Microbiology-Immunology), Veterinary Epidemiology, and Zoonotic Diseases as member of the academic staff. Postgraduate lecturer in the master program Advanced Animal Health and in the specialized course of Veterinary Clinics. He has supported the Cuban company for pork production and the provincial Veterinary Diagnostic Laboratory on aspects related with animal health. Head of the discipline Preventive Veterinary Medicine at UCLV.

In April 2007, he began a joint (sandwich) PhD in the field of "Swine diarrheic diseases" at the Department of Veterinary Medicine and Zootechnics, Faculty of Agriculture and Animal Sciences, UCLV, Cuba and the Laboratory of Immunology, Department of Virology, Parasitology, and Immunology, Faculty of Veterinary Medicine, Ghent University, Belgium.

**Publications in peer-reviewed journals**

Publications related with the PhD project in top 50% journals of Veterinary Sciences

- de la Fé Rodríguez PY, Coddens A, Del Fava E, Cortiñas Abrahantes J, Shkedy Z, Maroto Martin LO, Cruz Muñoz E, Duchateau L, Cox E, Goddeeris BM, 2011. High prevalence of F4<sup>+</sup> and F18<sup>+</sup> *Escherichia coli* in Cuban piggeries as determined by serological survey, Tropical Animal Health and Production, 43:937-46.
- de la Fé Rodríguez PY, Ndemi Kiiru J, Maroto Martin LO, Cruz Muñoz E, Butaye P, Cox E, Goddeeris BM, 2012. Characterization and clonal grouping of pathogenic *Escherichia coli* isolated from intestinal contents of diarrheic piglets in Villa Clara province, Cuba, according to their antibiotic resistance and ERIC-PCR profiles, Veterinary Microbiology, 154:425-8.

Submitted:

- de la Fé Rodríguez PY, Maroto Martin LO, Cruz Muñoz E, Imberechts H, Butaye P, Goddeeris BM, Cox E, 2011. Several enteropathogens are associated with porcine pre-weaning and post-weaning diarrhea in Villa Clara province, Cuba, Tropical Animal Health and Production (manuscript number TROP-2977).

Other publications

- Brito E, Hernández M, de la Fé Rodríguez PY, Silverio E, 2010. Prevalence and economic losses due to *Fasciola hepatica* in bovine slaughterhouses from three provinces located in Central Cuba (in Spanish). Revista Electrónica de Veterinaria REDVET 2(4). ISSN 1695-7504. Available in: <http://www.veterinaria.org/revistas/redvet/n040410/041006.pdf>



- de la Fé Rodríguez PY, Brito E, Aguiar J, Rodríguez L, Hernández JA, 2007. Prevalence of swine endoparasitosis in Cuba (in Spanish). REDVET 8(4). Available in: <http://www.veterinaria.org/revistas/redvet/n040407/040703.pdf>
- de la Fé Rodríguez PY, Brito E, Rodríguez L, Quiñones MC, 2006. Relevant aspects about the life cycle of *Taenia saginata* in Villa Clara province (in Spanish). REDVET 7(4). Available in: <http://www.veterinaria.org/revistas/redvet/n040406/040607.pdf>
- de la Fé Rodríguez PY, Duménigo BE, Brito E, Aguiar J, 2006. *Toxocara canis* and *Larva Migrans Visceralis* syndrome (in Spanish). REDVET 7(4). Available in: <http://www.veterinaria.org/revistas/redvet/n040406/040612.pdf>
- de la Fé Rodríguez PY, Rodríguez CM, Brito E, Simón E, Aguiar J, Rodríguez L, 2004. *Diphyllbothrium* spp. ¿an extinguished bothriocephalum in Villa Clara? (In Spanish). REDVET 5(8). Available in: <http://www.veterinaria.org/revistas/redvet/n080804.html>
- Gutierrez G, Garcia R, Cuesta M, Salado J, de la Fé Rodríguez PY, Aguiar J, González R, Camacho C, Avello E, Jimenez R, Herrada N, 2004. Analysis of the students promotion in the first years of the Veterinary Medicine career after the implementation of the study plans A, B, C, and C' during the last 26 academic years (in Spanish). Available in: <http://www.monografias.com/trabajos15/carrera-veterinaria/carrera-veterinaria.html>

#### Abstracts related with the PhD project in International Conferences proceedings

- de la Fé Rodríguez PY, Del Fava E, Cortiñas Abrahantes J, Shkedy Z, Maroto Martin LO, Cruz Muñoz E, Duchateau L, Cox E, Coddens A, Goddeeris BM, 2010. Seroprevalence of F4<sup>+</sup> and F18<sup>+</sup> *Escherichia coli* in Cuban young sows. In: D'Allaire S and Friendship R (Eds.), Proceedings of the 21<sup>st</sup> International Pig Veterinary Society (IPVS) Congress. Vancouver, British Columbia, Canada, pp. 764.
- de la Fé Rodríguez PY, Maroto Martin LO, Cruz Muñoz E, Imberechts H, Butaye P, Goddeeris BM, Cox E, 2010. Pathogens associated with pre and post-weaning porcine diarrhea in Cuba. In: grosse Beilage E and Blaha T (Eds.), Proceedings of the 2<sup>nd</sup> European Symposium on Porcine Health Management. Hannover, Germany, pp. 134.

#### Abstracts related with the PhD project in International Conferences held in Cuba

- Gonzalez Y, de la Fé Rodríguez PY, Cruz Muñoz E, 2011. Study of the seroprevalence of antibodies against the *Escherichia coli* fimbrial antigens F4 and F18 in pigs from Cuba. VII International Congress of Veterinary Sciences. Havana city, Cuba.
- Aguiar J, de la Fé Rodríguez PY, Cruz Muñoz E, Gonzalez A, Rodríguez A, Quiñónez R, Castillo JC, del Valle A, Valdez G, Rodríguez L, 2011. Pathogens associated with the neonatal and post-weaning diarrheic syndrome in pigs from Villa Clara. VII International Congress of Veterinary Sciences. Havana city, Cuba.
- Cruz Muñoz E, de la Fé Rodríguez PY, Gonzalez A, Maroto Martin LO, Rodríguez A, Gonzalez Y, Vera A, Aguiar J, 2009. Main pathologies in neonatal diarrheic piglets raised under tropical conditions. IV Conference about Sustainability and Agriculture Development. Faculty of Agriculture and Animal Sciences, UCLV, Santa Clara, Cuba.

### Attendance to other International Conferences

- Gutiérrez G, Aguiar J, Delgado M, Cruz E, Lazo L, de la Fé Rodríguez PY, Gonzalez R, García R, Pérez I, Salado J, Herrada N, 2002. Study of rotaviruses, *E. coli* K99 and *Cryptosporidium* spp. in diarrheic calves from the central provinces of Cuba. Proceedings of the XVIII Latin-American Congress of Veterinary Sciences. Havana city, Cuba.
- de la Fé Rodríguez PY, Olazabal E, Suarez YE, 2002. Incidence of fasciolosis in bovine and sheep from Villa Clara province. International Veterinary Congress-IICVV. II Veterinarians Virtual Congress. Málaga, Spain.
- de la Fé Rodríguez PY, Rodríguez L, Quiñonez MC, Brito E, 2002. Studying the epidemiology of *Taenia saginata-Cysticercus bovis* (sin. *C. inermis*) in Villa Clara province from 1998 to 2001. XVI Latin-American Congress of Microbiology. VI Cuban Congress of Microbiology and Parasitology. III Cuban Congress of Tropical Medicine. Institute of Tropical Medicine "Pedro Kourí", Havana city, Cuba.
- de la Fé Rodríguez PY, 2001. Bovine fasciolosis in Villa Clara province. I Conference about Sustainability and Agriculture Development, Symposium of Animal Health and Production. Faculty of Agriculture and Animal Sciences, UCLV, Santa Clara, Cuba.



## Word of thanks

It is a pleasure for me to thank people and institutions that supported me during the PhD project.

Prof. Bruno Goddeeris and Prof. Eric Cox, Prof. Eduardo Cruz and Prof. Luis Maroto, I was honoured to have you as promoters. Thanks of your meritorious advices we were able to make a good contribution to the Veterinary Medicine in Cuba and to finish this PhD project. All knowledge I got from you will guide me to advance on my professional and personal live. Thanks.

Members of the examination committee, I appreciate your time, advices and insightful suggestions, they contributed to improve the manuscript and my way of thinking. Prof. Piet Deprez, Prof. Luc Duchateau, Prof. Patrick Butaye, Prof. Edilbert Van Driessche, Prof. Eduardo Cruz and Prof. Hans Nauwynck: thanks very much.

Dr. Annelies Coddens, Prof. Ziv Shkedy, Prof. Hein Imberechts, Prof. Jose Cortinas Abrahantes, Ir. Emanuele del Fava, Dr. John Ndemi Kiiru, Prof. Patrick Butaye and Prof. Luc Duchateau: thanks for your help and scientific advice.

Special thanks go to the members of the Laboratory of Immunology at the Faculty of Veterinary Medicine of UGent, with them I have shared a nice time abroad and they have contributed to my results in different ways. Dear all your names written here as they are written on my heart will never permit that all you pass away from my mind: Gosia, Annelies, Maryam, Bakr, Ut, Philippe, Joanna, Anastasia, Evelien, Yu, Edith, Kris, Kristine, Petra, Vesna, Michaela, Delfien, Eva, Kim, Simon, Rudy, Denise, Griet, Sarah, Ann, Celine, Thary, Maria, Korneel, Pieter, Tine, Maaïke, Dirk, Mieke, Frank, Herman, Jochen, Cliff, Hanne, Nina, Mathias.

Colleagues at the Faculty of Plant and Animal Sciences and at the Department of Veterinary Medicine and Zootechnics of Universidad Central "Marta Abreu" de Las Villas: thanks for your confidence and support. I would like to specially thank Cary, Ania, Leopoldina, Javier, Alberto, Miguel, Eida, Alcides, Raciél, Salado, Einar, Andres, Tony, Cupull, Yanetsy, Osvaldo, Elio, Marianela.

Friends in Cuba Yoandys, Yake, William, Ernesto, Yannay, Yailen, Roberto, Mari, Daimi, Caridad, Omar, Izquierdo, Cira, Pedrin, Lester, Livan, Dania, Maikel: thanks for always being pending of my progress and all your best wishes for me. Karina and Alvaro thanks for being present always.

I gratefully acknowledge all institutions that made my PhD work possible. This PhD project started and ended thanks of the Belgian cooperation materialized through an Interuniversity Cooperation Program between VLIR-UOS and Universidad Central "Marta Abreu" de Las Villas, Cuba, and a Short Research Stay scholarship from VLIR-UOS. I would like to acknowledge the University of Ghent, the Cuban Company for Pork Production, and the Cuban Institute of Veterinary Medicine.

Last, but not least, I thank my family. Mom and Dad, thanks for the education you gave me and for your unconditional support to my plans. My sister Nani, my brother in law Alfredo and my nephews Alfreto and Abraham: you are inspiration for me. Tere, Jesu and Yake you are very special also.

Geidys thanks for your love which has always being on my side, in difficult and happy moments. Leslie de la Caridad, my daughter, you are going to be a month old and we have not met yet little girl!!!, sorry for that, and I have to confess you was the powerful inspiration to finish my PhD. I look forward to come back home to enjoy you.

A todos, **MUCHAS GRACIAS.**



